

Electronic Appendix 1. – Vascular plants endemic and subendemic to the Carpathians and their subunits. Apomictic taxa are marked (rows with a grey background; these taxa were omitted in the ecological analyses). If all subspecies of particular species are endemic, we mentioned the species also separately in the table (rows with an orange background; these species were not included in the statistical evaluations).

Abbreviations: AC – Apuseni Carpathians, EC – Eastern Carpathians, SC – Southern Carpathians, Tr – Transylvanian Basin, WC – Western Carpathians; A – Austria, CZ – Czech Republic, HU – Hungary, PL – Poland, RO – Romania, SK – Slovakia, SRB – Serbia, UA – Ukraine. Subunits with only scattered/sporadic occurrence of the evaluated taxon are listed in the end separated by a slash, e.g. EC (PL, UA, RO), SC (RO), AC (RO)/WC (PL).

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Achillea oxyloba</i> subsp. <i>schurii</i> (Sch. Bip.) Heimerl (Syn.: <i>A. schurii</i> Sch. Bip.; <i>Ptarmica tenuifolia</i> (Schur) Schur)	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Morariu & Beldie 1976; Oprea 2005; Ciocârlan 2009; Počynok & Prokopiv 2010; Sârbru et al. 2013; Ziman & Derbak 2013
<i>Aconitum bucovinense</i> Zapał. (Syn.: <i>A. callibotryon</i> subsp. <i>bucovinense</i> (Zapał.) Grinț.; <i>A. firmum</i> subsp. <i>bucovinense</i> (Zapał.) Graebn. et P. Graebn.)	EC (PL, UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Starmühler 1998a, b, 2000; Mitka 2001, 2002, 2003, 2012; Ilnicki & Mitka 2009; Boroń et al. 2011; Novikoff & Mitka 2011a, b
<i>Aconitum degenii</i> Gayer subsp. <i>degenii</i> ¹ (Syn.: <i>A. paniculatum</i> subsp. <i>degenii</i> (Degen) Graebn.)	WC (SK), EC (PL, UA, RO), SC (RO), AC (RO)	West-East-South-Apuseni-Carpathian (pan-Carpathian) endemic	Mucher 1993; Starmühler 1997, 1998a, b, 2000; Mitka 2001, 2003; Ilnicki & Mitka 2011; Novikoff & Mitka 2011a, b; Novikov 2013a; Eliáš jr. et al. 2015
<i>Aconitum firmum</i> Rchb.	WC (CZ, SK, PL), EC (UA, RO), SC (RO), AC (RO)	West-East-South-Apuseni-Carpathian (pan-Carpathian) subendemic	Starmühler 1997, 1998b, 2000; Mitka 2001, 2003; Starmühler & Mitka 2001; Mihok et al. 2005; Novikoff & Mitka 2011a, b
<i>Aconitum firmum</i> Rchb. subsp. <i>firmum</i> (Syn.: <i>A. napellus</i> subsp. <i>firmum</i> (Rchb.) Gayer; <i>A. firmum</i> subsp. <i>palmatifidum</i> (Rchb.) Beldie)	WC (SK, PL), EC (UA, RO), SC (RO), AC (RO)	West-East-South-Apuseni-Carpathian (pan-Carpathian) endemic	Skalický 1990; Starmühler 1997, 1998b, 2000; Mitka 2001, 2003; Starmühler & Mitka 2001; Mitka et al. 2007; Novikoff & Mitka 2011a
<i>Aconitum firmum</i> subsp. <i>fissurae</i> Nyár. ^{2,3} (Syn.: <i>A. hunyadense</i> Degen; <i>A. romanicum</i> Wol.)	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian subendemic	Starmühler 1996a, 1997, 1998b, 1999, 2000; Mitka 2001, 2002; Starmühler & Mitka 2001; Ilnicki & Mitka 2009; Novikoff & Mitka 2011a
<i>Aconitum firmum</i> subsp. <i>maninense</i> (Skalický) Starm. (Syn.: <i>A. firmum</i> var. <i>maninense</i> Skalický)	WC (SK, PL)	West-Carpathian endemic	Starmühler & Mitka 2001; Mitka 2003; Mitka et al. 2007; Ilnicki & Mitka 2009
<i>Aconitum firmum</i> subsp. <i>moravicum</i> Skalický	WC (CZ, SK, PL)	West-Carpathian endemic	Skalický 1990; Mitka 2003; Mitka et al. 2007; Ilnicki & Mitka 2009
<i>Aconitum firmum</i> subsp. <i>skerisorae</i> (Gayer) Starm. ⁴ (Syn.: <i>A. skerisorae</i> Gayer; <i>A. napellus</i> subsp. <i>skerisorae</i> (Gayer) Seitz)	AC (RO)	Apuseni-Carpathian endemic	Starmühler 2000; Mitka 2003
<i>Aconitum lasianthum</i> (Rchb.) Simonk. ⁵ (Syn.: <i>A. vulparia</i> subsp. <i>lasianthum</i> (Rchb.) Ciocârlan)	SC (RO)/EC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Grințescu 1953; Morariu & Beldie 1976; Starmühler 1999; Oprea 2005
<i>Aconitum lasiocarpum</i> (Rchb.) Gayer (Syn.: <i>A. paniculatum</i> subsp. <i>lasiocarpum</i> (Rchb.) Soó; <i>A. toxicum</i> subsp. <i>lasiocarpum</i> (Rchb.) Grinț.)	WC (SK, PL), EC (SK, PL, UA, RO), †SC (RO)	West-East-Carpathian subendemic	Joachimik et al. 1999; Mitka & Starmühler 2000; Mitka 2001, 2003; Ilnicki & Mitka 2011; Novikoff & Mitka 2011a, b
<i>Aconitum lasiocarpum</i> (Rchb.) Gayer subsp. <i>lasiocarpum</i> ⁶	EC (SK, PL, UA, RO)	East-Carpathian endemic	Mitka & Starmühler 2000; Mitka 2001, 2003; Ilnicki & Mitka 2011; Novikoff & Mitka 2011a, b
<i>Aconitum lasiocarpum</i> subsp. <i>kotulae</i> (Pawl.) Starm. et Mitka ^{7,8} (Syn.: <i>A. variegatum</i> subsp. <i>kotulae</i> Pawl.)	WC (SK, PL), EC (SK, PL, UA, RO), †SC (RO)	West-East-Carpathian subendemic	Mitka & Starmühler 2000; Mitka 2001, 2003; Ilnicki & Mitka 2011; Novikoff & Mitka 2011a, b

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Aconitum moldavicum</i> Hacq. (Syn.: <i>A. lycoctonum</i> subsp. <i>moldavicum</i> (Hacq.) Jalas)	WC (SK, PL, HU), EC (SK, PL, UA, RO), SC (RO), AC (RO)	West-East-South-Apuseni-Carpathian (pan-Carpathian) subendemic	Mitka 2003; Mihok et al. 2005; Novikoff & Mitka 2011a, b; Mitka et al. 2013
<i>Aconitum moldavicum</i> Hacq. subsp. <i>moldavicum</i> ⁹	WC (SK, PL, HU), EC (SK, PL, UA, RO), SC (RO), AC (RO)	West-East-South-Apuseni-Carpathian (pan-Carpathian) subendemic	Mitka 2003, 2008; Novikoff & Mitka 2011a, b
<i>Aconitum moldavicum</i> subsp. <i>hosteanum</i> (Schur) Graebn. et P. Graebn. ¹⁰ (Syn.: <i>A. hosteanum</i> Schur; <i>A. moldavicum</i> subsp. <i>hosteanum</i> (Schur) Beldie, nom. illeg.)	EC (PL, UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian subendemic	Mitka 2003, 2008; Čornej 2011; Novikoff & Mitka 2011a, b
<i>Aconitum toxicum</i> Rchb. subsp. <i>toxicum</i> ¹¹	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian subendemic	Mucher 1993; Mitka 2001; Ilnicki & Mitka 2011
<i>Aconitum toxicum</i> subsp. <i>bucegiense</i> (Nyár.) Mucher	EC (RO), SC (RO)	East-South-Carpathian endemic	Mucher 1993; Mitka 2001; Starmühler & Mitka 2001
<i>Aconitum toxicum</i> subsp. <i>crispulum</i> (Nyár.) Mucher	SC (RO)	South-Carpathian endemic	Mucher 1993; Starmühler & Mitka 2001
<i>Alchemilla acrostegia</i> Plocek	WC (SK)	West-Carpathian endemic (Západné Tatry Mts: Červené vrchy massif)	Plocek 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla aequidens</i> Pawł.	WC (SK, PL)	West-Carpathian endemic	Plocek 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla amauroptera</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla amblyoides</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla amicorum</i> Pawł.	WC (SK, PL)	West-Carpathian endemic (Západné Tatry Mts: Červené vrchy massif)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla anceps</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla animosa</i> Plocek	WC (SK)	West-Carpathian endemic	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla aspera</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla babiogorensis</i> Pawł.	WC (PL), EC (UA)	West-East-Carpathian endemic	Volgin & Syčák 1989a; Plocek 1992; Syčák 2002, 2011
<i>Alchemilla bogumili</i> Pawłus	WC (SK, PL)	West-Carpathian endemic	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla boleslai</i> Pawł. ¹²	WC (SK, PL)	West-Carpathian endemic	Plocek 1983, 1992; Kliment 1999; Kurtto et al. 2007; Negrean 2011
<i>Alchemilla brachycodon</i> Plocek	WC (SK)	West-Carpathian endemic (Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla braun-blanquetii</i> Pawł.	WC (PL), EC (UA)	West-East-Carpathian endemic	Plocek 1992; Syčák 2002, 2011; Kurtto et al. 2007
<i>Alchemilla bucovinensis</i> Sytschak ¹³	EC (UA, RO)	East-Carpathian endemic	Syčák 1992, 2002, 2011; Kurtto et al. 2007
<i>Alchemilla calviflora</i> Plocek	WC (PL)	West-Carpathian endemic (Tatry Zachodnie Mts: Czerwone Wierchy massif)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla chalarodesma</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1990; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla chilitricha</i> Plocek	WC (SK)	West-Carpathian endemic (Západné Tatry Mts: Sivý vrch Mt. group)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla contractilis</i> (Plocek) Fröhner	WC (SK), SC (RO)	West-South-Carpathian endemic	Plocek 1992; Kurtto et al. 2007; Negrean 2011
<i>Alchemilla crassa</i> Plocek	WC (SK)	West-Carpathian endemic	Plocek 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla curtischista</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla czywczyensis</i> Pawł. ¹⁴	EC (UA, RO)	East-Carpathian subendemic	Pawłowski 1952; Fröhner 1986; Syčák 2002, 2011; Kurtto et al. 2007, 2009

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Alchemilla decurrens</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla delitescens</i> Plocek	WC (SK)	West-Carpathian endemic (Oravské Beskydy Mts)	Plocek 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla deyllii</i> Plocek ex Soják	EC (UA)	East-Carpathian endemic	Soják 1983b; Syčák 2002, 2011; Kurtto et al. 2007
<i>Alchemilla dolichotoma</i> Plocek	EC (RO), SC (RO)	East-South-Carpathian endemic	Plocek 1985; Kurtto et al. 2007
<i>Alchemilla dostalii</i> Plocek	WC (SK)	West-Carpathian endemic (Západné Tatry Mts: Mt. Osobitá)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla echinogloba</i> Plocek	WC (SK)	West-Carpathian endemic (Oravské Beskydy Mts: Mt. Pilsko)	Plocek 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla erythropodoides</i> Pawł.	WC (SK)	West-Carpathian endemic (Tatry Mts)	Fröhner 1968, 1975; Kurtto et al. 2007
<i>Alchemilla eugenii</i> Pawł. ¹⁵	WC (PL, ?SK)	West-Carpathian endemic (Tatry Zachodnie Mts: Czerwone Wierchy massif)	Pawłowski 1952; Plocek 1992; Kurtto et al. 2007
<i>Alchemilla exaperta</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla fusoides</i> Plocek	WC (SK)	West-Carpathian endemic (Oravské Beskydy Mts: Mt. Pilsko)	Plocek 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla giewontica</i> Pawł.	WC (PL)	West-Carpathian endemic (Tatry Zachodnie Mts: Mt. Giewont)	Plocek 1992; Kurtto et al. 2007; Mirek & Piękoś-Mirkowa 2010
<i>Alchemilla grandiceps</i> Plocek	WC (SK)	West-Carpathian endemic (Oravské Beskydy Mts: Mt. Babia hora)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla gruneica</i> Plocek ¹⁶	WC (CZ, SK)	West-Carpathian subendemic	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla gymnopoda</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla hoverlensis</i> Pawlus et Lovelius	EC (UA)	East-Carpathian endemic (Čornohora Mts)	Pawlus 1988; Syčák 1992, 2011; Kurtto et al. 2007
<i>Alchemilla hyperptycha</i> Plocek	WC (SK)	West-Carpathian endemic (Vysoké Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla isodonta</i> Plocek	WC (SK)	West-Carpathian endemic (Nízke Beskydy Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla jasiewiczii</i> Pawł.	WC (PL)	West-Carpathian endemic (Tatry Wysokie Mts)	Plocek 1992; Kurtto et al. 2007; Mirek & Piękoś-Mirkowa 2010
<i>Alchemilla kornasiana</i> Pawł.	WC (PL), EC (UA)	West-East-Carpathian endemic	Plocek 1992; Syčák 2002, 2011; Kurtto et al. 2007
<i>Alchemilla kosiarensis</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla kulczyńskii</i> Pawł.	WC (SK, PL)	West-Carpathian endemic (Západné Tatry Mts: Červené vrchy massif)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla ladislai</i> Pawł. ¹⁷	WC (SK, PL)	West-Carpathian endemic (Západné Tatry Mts: Červené vrchy massif)	Plocek 1992; Syčák 2002, 2011; Kurtto et al. 2007
<i>Alchemilla laevipes</i> Plocek	WC (SK)	West-Carpathian endemic (Oravské Beskydy Mts: Mt. Babia hora)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla laxa</i> Plocek ¹⁸	WC (SK)	West-Carpathian endemic (Oravské Beskydy Mts: Mt. Pilsko)	Plocek 1990, 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla longidens</i> Plocek	WC (SK)	West-Carpathian endemic (Oravské Beskydy Mts: Mt. Babia hora)	Plocek 1990, 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla lorata</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla loxotropa</i> Plocek	WC (SK)	West-Carpathian endemic (Oravské Beskydy Mts: Mt. Babia hora)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla ludovitiana</i> Plocek	WC (SK)	West-Carpathian endemic (Branisko Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla marginata</i> Plocek ¹⁹	WC (SK)	West-Carpathian endemic (Vysoké Tatry Mts)	Plocek 1992; Kurtto et al. 2007

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Alchemilla megalodonta</i> Plocek	WC (SK)	West-Carpathian endemic (Vysoké Tatry Mts)	Plocek 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla microsphaerica</i> Fröhner	WC (SK, PL)	West-Carpathian endemic	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla monocophila</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla multiloba</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla obesa</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla oculimarina</i> Pawł.	WC (SK, PL)	West-Carpathian endemic (Vysoké Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla patens</i> Plocek	WC (SK)	West-Carpathian endemic (Nízke Tatry Mts)	Plocek 1992; Šípošová et al. 2004b; Kurtto et al. 2007
<i>Alchemilla polonica</i> Pawł. (Syn.: <i>A. pseudincisa</i> var. <i>polonica</i> (Pawł.) Plocek)	WC (SK, PL)	West-Carpathian endemic	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla pseudincisa</i> Pawł.	WC (SK, PL), EC (UA)	West-East-Carpathian endemic	Volgin & Syčák 1989b; Plocek 1992; Syčák 2002, 2011; Kurtto et al. 2007
<i>Alchemilla pseudothmari</i> Pawł.	WC (PL)	West-Carpathian endemic (Tatry Zachodnie Mts: Czerwone Wierchy massif)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla reversantha</i> Plocek (Syn.: <i>A. inversa</i> Plocek non Juz., nom. illeg.)	WC (SK)	West-Carpathian endemic (Veľká Fatra Mts)	Plocek 1992; Šípošová et al. 2004a; Kurtto et al. 2007
<i>Alchemilla rhodobasis</i> Plocek	WC (SK)	West-Carpathian endemic (Západné Tatry Mts: Sivý vrch Mt. group)	Plocek 1986, 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla rhodocycla</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla sejuncta</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla sericoneuroides</i> Pawł.	WC (SK, PL)	West-Carpathian endemic (Tatry Mts)	Plocek 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla smaragdina</i> Plocek	WC (SK)	West-Carpathian endemic	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla smytniensis</i> Pawł.	WC (PL), EC (UA)	West-East Carpathian endemic	Syčák 1992, 2011; Kurtto et al. 2007
<i>Alchemilla sojakii</i> Plocek	WC (SK)	West-Carpathian endemic (Krivánska Fatra Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla sokolowskii</i> Pawł. ²⁰	WC (PL)	West-Carpathian endemic (Tatry Zachodnie Mts: Czerwone Wierchy massif)	Plocek 1992; Kurtto et al. 2007; Mirek & Piękoś-Mirkowa 2010
<i>Alchemilla stanislaae</i> Pawł.	WC (SK, PL)	West-Carpathian endemic (Tatry Mts, Nízke Tatry Mts) ²¹	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla stenoleuca</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla suavis</i> Plocek ²²	WC (CZ, SK)	West-Carpathian endemic	Plocek 1973, 1992; Syčák 1992, 2011; Kurtto et al. 2007
<i>Alchemilla subsessilis</i> Plocek	WC (SK)	West-Carpathian endemic (Oravské Beskydy Mts: Mt. Pilsko)	Plocek 1992; Kliment 1999; Kurtto et al. 2007
<i>Alchemilla superata</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla szaferi</i> Pawł.	EC (UA)	East-Carpathian endemic	Syčák 2002, 2011; Kurtto et al. 2007
<i>Alchemilla tacikii</i> Plocek	WC (SK, PL)	West-Carpathian endemic (Vysoké Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla thaumasia</i> Plocek	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla turkulensis</i> Pawł.	EC (PL, UA)	East-Carpathian endemic	Syčák 2002, 2011; Kurtto et al. 2007
<i>Alchemilla versipiloides</i> Pawł. ²³	WC (PL)	West-Carpathian endemic (Tatry Zachodnie Mts: Czerwone Wierchy massif)	Plocek 1992; Kurtto et al. 2007; Piękoś-Mirkowa & Mirek 2009; Mirek & Piękoś-Mirkowa 2010
<i>Alchemilla virginea</i> Plocek	WC (SK)	West-Carpathian endemic (Krivánska Fatra Mts)	Plocek 1992; Kurtto et al. 2007

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Alchemilla wallischii</i> Pawł.	WC (SK, PL)	West-Carpathian endemic (Tatry Mts, Nízke Tatry Mts)	Plocek 1992; Kurtto et al. 2007
<i>Alchemilla zapalowiczii</i> Pawł. ²⁴	WC (†SK), EC (UA)	East-Carpathian subendemic	Walters & Pawłowski 1968; Plocek 1992; Kurtto et al. 2007, 2009; Negrean 2011
<i>Alchemilla zmudae</i> Pawł. ²⁵	WC (PL)	West-Carpathian endemic (Tatry Mts)	Plocek 1992; Kurtto et al. 2007; Mirek & Piękoś-Mirkowa 2009, 2010
<i>Allium fussii</i> A. Kern. ²⁶ (Syn.: <i>A. fuscum</i> subsp. <i>fussii</i> (A. Kern.) Ciocârlan; <i>A. paniculatum</i> subsp. <i>fussii</i> (A. Kern.) Ciocârlan, nom. illeg.)	EC (RO), SC (RO)	East-South-Carpathian endemic	Brullo et al. 1996; Mráz 2005a; Oprea 2005; Ciocârlan 2009; Šafařová et al. 2011
<i>Alopecurus pratensis</i> subsp. <i>laguriformis</i> (Schur) Tzvelev (Syn.: <i>A. laguriformis</i> Schur)	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Morariu & Beldie 1976; Oprea 2005; Čornej 2011; Tasenkevich 2011
<i>Andryala laevitomentosa</i> (Nyár. ex Sennikov) P. D. Sell ex Greuter ^{27, 28} (Syn.: <i>Pietrosia laevitomentosa</i> Nyár. ex Sennikov)	EC (RO)	East-Carpathian endemic (Bistrița Mts: Mt. Pietrosul Broștenilor)	Morariu & Beldie 1976; Beldie 1979; Dihoru & Pârvu 1987; Sennikov 1999a; Greuter 2003; Oprea 2005, 2007; Negrea & Pricop 2009a, b; Ferreira et al. 2015
<i>Antennaria carpatica</i> (Wahlenb.) Bluff et Fingerh. subsp. <i>carpatica</i> ²⁹	WC (SK, PL), EC (UA); †RO (EC, SC)	West-East-Carpathian endemic	Chrtěk & Pouzar 1985; Oltean et al. 1994; Kliment 1999; Oprea 2005; Ciocârlan 2009; Sârbu et al. 2013; Ziman & Derbak 2013
<i>Anthemis cretica</i> subsp. <i>pyrethriiformis</i> (Schur) Govaerts ^{30, 31} (Syn.: <i>A. carpatica</i> subsp. <i>pyrethriiformis</i> (Schur) Prodan)	SC (RO)/EC (RO)	East-South-Carpathian endemic	Morariu & Beldie 1976; Beldie 1979; Oprea 2005; Sârbu et al. 2013
<i>Anthemis kitaibelii</i> Spreng. (Syn.: <i>A. cretica</i> subsp. <i>kitaibelii</i> (Spreng.) Ciocârlan)	SC (RO), AC (RO)	South-Apuseni-Carpathian endemic	Oprea 2005; Hurdu et al. 2012a, b
<i>Aquilegia nigricans</i> subsp. <i>subscaposa</i> (Borbás) Soó (Syn.: <i>A. subscaposa</i> Borbás; <i>A. vulgaris</i> subsp. <i>subscaposa</i> (Borbás) Borza)	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Morariu & Beldie 1976; Oprea 2005; Dihoru & Negrean 2009
<i>Aquilegia transsilvanica</i> Schur ³²	SC (RO)/EC (RO)	East-South-Carpathian endemic	Oprea 2005; Kobiv 2012b
<i>Arabidopsis halleri</i> subsp. <i>tatrica</i> (Pawł.) Kolník (Syn.: <i>Cardaminopsis halleri</i> subsp. <i>tatrica</i> (Pawł.) Dostál ex Měsíček)	WC (SK, PL)	West-Carpathian endemic	Kolník & Marhold 2006
<i>Arabidopsis neglecta</i> (Schult.) O’Kane et Al-Shehbaz ³³ (Syn.: <i>Cardaminopsis neglecta</i> (Schult.) Hayek)	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Jalas & Suominen 1994; Kliment 1999; Měsíček 2002; Oprea 2005
<i>Arenaria tenella</i> Kit. (Syn.: <i>A. ciliata</i> subsp. <i>tenella</i> (Kit.) Braun-Blanq.)	WC (SK, PL)	West-Carpathian endemic (Tatry Mts)	Perný & Michalková 2012
<i>Armeria maritima</i> subsp. <i>barcensis</i> (Simonk.) P. Silva (Syn.: <i>A. barcensis</i> Simonk.)	EC (RO)	East-Carpathian endemic	Beldie 1979; Dihoru & Pârvu 1987; Ciocârlan 2009
<i>Armeria pocutica</i> Pawł. ³⁴	EC (?†UA, RO)	East-Carpathian endemic (Maramureș Mts)	Pawłowski 1962; Ciocârlan 1988, 2009; Kahalo & Syčák 2009; Negrean 2011
<i>Asperula carpatica</i> Morariu ³⁵	EC (RO)/SC (RO)	East-South-Carpathian endemic	Negrean 2011
<i>Astragalus australis</i> subsp. <i>krajinae</i> (Domin) Domin ³⁶ (Syn.: <i>A. krajinae</i> Domin)	EC (UA)	East-Carpathian endemic	Domin 1931a, 1935; Witkowski et al. 2003; Ziman 2009; Čornej 2011
<i>Astragalus exscapus</i> subsp. <i>transsilvanicus</i> (Schur) Nyár. (Syn.: <i>A. exscapus</i> var. <i>transsilvanicus</i> (Schur) Gams)	Tr (RO)	endemic to Transylvanian Basin	Roman et al. 1996; Dihoru & Negrean 2009

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Astragalus peterfii</i> Jáv. ³⁷	Tr (RO)	endemic to Transylvanian Basin	Roman et al. 1996; Şuteu et al. 2003; Dihoru & Negrean 2009; Bartha 2012
<i>Astragalus pseudopurpureus</i> Guşul.	EC (RO)	East-Carpathian endemic (Hășmaş Mts)	Ciocârlan 2009; Dihoru & Negrean 2009
<i>Astragalus roemerii</i> Simonk. ³⁸	EC (RO)/AC (RO)	East-Apuseni-Carpathian endemic	Váczy & Beldie 1976; Dihoru & Negrean 2009; Bartha & Bartók 2013
<i>Athamanta turbith</i> subsp. <i>hungarica</i> (Borbás) Tutin ³⁹ (Syn.: <i>A. hungarica</i> Borbás)	SC (RO, SRB)	South-Carpathian endemic	Stevanović et al. 1991; Popescu et al. 2003; Hurdu et al. 2012a, b; Sârbu et al. 2013
<i>Aubrieta columnae</i> subsp. <i>platycarpa</i> (Ciocârlan) Ciocârlan ⁴⁰ (Syn.: <i>A. intermedia</i> subsp. <i>falcata</i> Ciocârlan)	SC (RO)	South-Carpathian endemic (Piatra Craiului Mts)	Beldie & Váczy 1976; Oprea 2005; Ciocârlan 2006, 2009; Negrean 2011
<i>Barbarea lepuznica</i> Nyár. ⁴¹ (Syn.: <i>B. vulgaris</i> subsp. <i>lepuznica</i> (Nyár.) Soó)	SC (RO)	South-Carpathian subendemic	Dihoru & Pârvu 1987; Ciocârlan 2009; Dihoru & Negrean 2009; Strajeru & Stevanović 2013
<i>Bromus monocladus</i> Domin ⁴² (Syn.: <i>B. pannonicus</i> subsp. <i>monocladus</i> (Domin) P. M. Sm.; <i>Bromopsis pannonica</i> subsp. <i>monoclada</i> (Domin) Holub)	WC (SK)	West-Carpathian endemic	Kliment 1999; Dúbravková 2014 in litt.; Somlyay 2014 in litt.
<i>Campanula carpatica</i> Jacq.	WC (SK), EC (UA, RO), SC (RO)/AC	West-East-South-Apuseni-Carpathian (pan-Carpathian) endemic	Kliment 1999; Goliašová et al. 2008
<i>Campanula crassipes</i> Heuff.	SC (RO, SRB)	South-Carpathian endemic	Hurdu et al. 2012a, b
<i>Campanula glomerata</i> subsp. <i>subcapitata</i> (Popov) Fed. (Syn.: <i>C. subcapitata</i> Popov)	EC (SK, UA)	East-Carpathian endemic	Kliment 1999; Ziman et al. 2006; Goliašová et al. 2008
<i>Campanula kladniana</i> (Schur) Witasek ⁴³ (Syn.: <i>C. rotundifolia</i> subsp. <i>kladniana</i> (Schur) Tacik)	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Čopyk 1976; Tasenkevych 2003b; Oprea 2005; Ziman et al. 2006
<i>Campanula serrata</i> (Kit.) Hendrych (Syn.: <i>C. napuligera</i> Schur)	WC (SK, PL), EC (UA, RO), SC (RO), AC	West-East-South-Apuseni-Carpathian (pan-Carpathian) endemic	Kliment 1999; Goliašová et al. 2008
<i>Campanula tatrae</i> Borbás ⁴⁴ (Syn.: <i>C. polymorpha</i> (Witasek) Prain et al.; <i>C. rotundifolia</i> subsp. <i>polymorpha</i> (Witasek) Tacik)	WC (SK, PL), EC (UA, RO), SC (RO), AC (RO)	West-East-South-Apuseni-Carpathian (pan-Carpathian) endemic	Kliment 1999; Goliašová et al. 2008
<i>Campanula xylocarpa</i> Kovanda	WC (SK, HU)	West-Carpathian endemic	Kliment 1999; Goliašová et al. 2008
<i>Cardamine glanduligera</i> O. Schwarz ⁴⁵ (Syn.: <i>Dentaria glandulosa</i> Waldst. et Kit. ex Willd.)	WC (CZ, SK, PL, HU), EC (SK, PL, UA, RO), SC (RO), AC (RO), Tr (RO)	subendemic to Western, Eastern, Southern & Apuseni Carpathians and Transylvanian Basin (pan-Carpathian subendemic)	Jalas & Suominen 1994; Roman et al. 1996; Kliment 1999; Marhold & Kochjarová 2002
<i>Carduus kernerii</i> Simonk. subsp. <i>kernerii</i> ⁴⁶ (Syn.: <i>C. transsilvanicus</i> A. Kern.)	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Franco 1976; Čopyk 1976; Oprea 2005; Čornej 2011; Hurdu et al. 2012a, b; Ziman & Derbak 2013
<i>Carduus kernerii</i> subsp. <i>lobulatiformis</i> (Csüros et Nyár.) Soó ⁴⁷ (Syn.: <i>C. lobulatiformis</i> Csüros et Nyár.)	SC (RO)	South-Carpathian endemic	Dihoru & Pârvu 1987; Ciocârlan 2009; Dihoru & Negrean 2009; Hurdu et al. 2012a
<i>Carduus lobulatus</i> Borbás	WC (SK, PL)	West-Carpathian endemic	Kliment 1999; Šípošová et al. 2004a; Zarzycki 2008
<i>Centaurea reichenbachii</i> DC. ⁴⁸ (Syn.: <i>C. reichenbachii</i> Schur ex Hayek)	AC (RO)	Apuseni-Carpathians endemic	Dostál 1976; Ochsmann 2000; Hurdu et al. 2012a; Ciocârlan 2009; Sârbu et al. 2013
<i>Centaurea rodensis</i> Simonk. ^{49, 50, 51} (Syn.: <i>C. carpatica</i> (Porcius) Porcius, nom. illeg.; <i>C. rarauensis</i> Prodan)	EC (?UA, RO)	East-Carpathian endemic (Rodna Mts)	Beldie 1979; Tasenkevich 2011; Hurdu et al. 2012a, b; Koutecký 2013

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Centaurea simonkaiana</i> Hayek (Syn.: <i>C. trichocephala</i> subsp. <i>simonkaiana</i> (Hayek) Dostál)	AC (RO)	Apuseni-Carpathian endemic	Dihoru & Parvu 1987; Dihoru & Negrean 2009
<i>Centaurea triniifolia</i> Heuff. ⁵²	SC (RO, SRB)	South-Carpathian endemic	Dostál 1976; Ochsmann 2000; Hurdu et al. 2012a
<i>Cephalaria radiata</i> Griseb. et Schenk ⁵³	Tr (RO)/EC (RO), SC (RO), AC (RO)	subendemic to Transylvanian Basin	Prodan 1961; Dihoru & Pârvu 1987; Oprea 2005
<i>Cephalaria uralensis</i> subsp. <i>multifida</i> (Roman) Roman et Beldie ⁵⁴	SC (RO)	South-Carpathian endemic (Almajului Mts: Porțile de Fier Gorge)	Beldie & Váczy 1976; Dihoru & Pârvu 1987; Ciocârlan 2011; Sârbu et al. 2013
<i>Cerastium arvense</i> subsp. <i>lerchenfeldianum</i> (Schur) Asch. et Graebn. ^{55, 56} (Syn.: <i>C. lerchenfeldianum</i> Schur)	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Oprea 2005; Negrean & Oltean 1989; Hurdu et al. 2012b
<i>Cerastium tatrae</i> Borbás (Syn.: <i>C. glandulosum</i> (Kit.) Jáv. non Schur, nom. illeg.; <i>C. arvense</i> subsp. <i>glandulosum</i> (Kit.) Soó)	WC (SK, PL)	West-Carpathian endemic (Tatry Mts)	Kliment 1999; Letz & Michalková 2012
<i>Cerastium transsilvanicum</i> Schur	EC (RO), SC (RO)	East-South-Carpathian endemic	Čopyk 1976; Morariu & Beldie 1976; J alas & Suominen 1983; Oprea 2005; Ciocârlan 2009
<i>Chrysosplenium alpinum</i> Schur	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Beldie 1977; Oprea 2005; Kobiv 2010; Čornej 2011; Sârbu et al. 2013
<i>Clinopodium pulegium</i> (Rochel) Bräuchler ⁵⁷ (Syn.: <i>Micromeria pulegium</i> (Rochel) Benth., nom. illeg.)	SC (RO, SRB)	South-Carpathian subendemic	Diklić & Nikolić 1986a; Bogosavljević et al. 2007; Hurdu et al. 2012b; Đug et al. 2013; Slavkovska et al. 2013
<i>Cochlearia borzaeana</i> (Coman et Nyár.) Pobed.	EC (RO)	East-Carpathian endemic	Kochjarová et al. 2006; Dihoru & Negrean 2009
<i>Cochlearia tatrae</i> Borbás	WC (SK, PL)	West-Carpathian endemic (Tatry Mts)	Kliment 1999; Kochjarová & Valachovič 2002
<i>Cota tinctoria</i> subsp. <i>fussii</i> (Griseb. et Schenk) Oberpr. et Greuter (Syn.: <i>Anthemis tinctoria</i> subsp. <i>fussii</i> (Griseb. et Schenk) Beldie)	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathians endemic	Dihoru & Pârvu 1987; Oprea 2005
<i>Crocus banaticus</i> J. Gay ⁵⁸	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian subendemic	Mihály & Komendar 1993; Oprea 2005; Myhaľ 2009; Hurdu et al. 2012b
<i>Crocus discolor</i> G. Reuss (Syn.: <i>C. scepusiensis</i> (Rehman et Woł.) Borbás)	WC (SK, PL)	West-Carpathian endemic	Májovský et al. 1991; Kliment 1999
<i>Cyanus dominii</i> (Dostál) Holub	WC (SK)	West-Carpathian endemic	Olšovská et al. 2011, 2012; Olšovská & Löser 2013
<i>Cyanus dominii</i> (Dostál) Holub subsp. <i>dominii</i> (Syn.: <i>Cyanus triumfetti</i> subsp. <i>dominii</i> (Dostál) Dostál; <i>Centaurea triumfetti</i> subsp. <i>dominii</i> Dostál)	WC (SK)	West-Carpathian endemic	Olšovská et al. 2011, 2012; Olšovská & Löser 2013
<i>Cyanus dominii</i> subsp. <i>slovenicus</i> (Dostál) Olšovská (Syn.: <i>Centaurea triumfetti</i> var. <i>slovenica</i> Dostál)	WC (SK)	West-Carpathian endemic	Olšovská et al. 2011, 2012; Olšovská & Löser 2013
<i>Cyanus dominii</i> subsp. <i>sokolensis</i> (Pawł.) Olšovská (Syn.: <i>Centaurea axillaris</i> var. <i>sokolensis</i> Pawł.)	WC (SK)	West-Carpathian endemic	Olšovská et al. 2011, 2012; Olšovská & Löser 2013
<i>Cyanus maramarosiensis</i> (Jáv.) Dostál ⁵⁹ (Syn.: <i>Centaurea maramarosiensis</i> (Jáv.) Czerep.; <i>C. mollis</i> subsp. <i>maramarosiensis</i> (Jáv.) Soó)	EC (?SK, UA, RO)	East-Carpathian endemic	Kliment 1999; Oprea 2005; Ciocârlan 2009; Sârbu et al. 2013
<i>Cyanus mollis</i> (Waldst. et Kit.) J. Presl et C. Presl ⁶⁰ (Syn.: <i>Centaurea mollis</i> Waldst. et Kit.; <i>Cyanus montanus</i> subsp. <i>mollis</i> (Waldst. et Kit.) Soják)	WC (CZ, SK, PL, HU), EC (UA, RO), SC (RO), AC (RO), Tr (RO)	subendemic to Western, Eastern, Southern & Apuseni Carpathians and Transylvanian Basin (pan-Carpathian subendemic)	Kliment 1999; Oprea 2005; Olšovská & Löser 2013

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Cyanus pinnatifidus</i> (Schur) Holub (Syn.: <i>Centaurea pinnatifida</i> Schur; <i>C. triumfetti</i> subsp. <i>pinnatifida</i> (Schur) Dostál)	EC (RO), SC (RO), AC	East-South-Apuseni-Carpathian endemic	Oprea 2005; Olšovská et al. 2012
<i>Cyanus pinnatifidus</i> (Schur) Holub subsp. <i>pinnatifidus</i> (Syn.: <i>Centaurea pinnatifida</i> Schur subsp. <i>pinnatifida</i>)	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Oprea 2005
<i>Cyanus pinnatifidus</i> subsp. <i>sooanus</i> (Borhidi) Greuter ⁶¹ (Syn.: <i>Centaurea pinnatifida</i> subsp. <i>sooana</i> (Borhidi) Soó)	EC (RO)	East-Carpathian endemic (Ceahlău Mts)	Borhidi 1957; Oprea 2005
<i>Cyclamen purpurascens</i> subsp. <i>immaculatum</i> (Hrabětová) Halda et Soják ⁶² (Syn.: <i>C. fatrense</i> Halda et Soják)	WC (SK)	West-Carpathian endemic (Veľká Fatra Mts, Nízke Tatry Mts)	Kliment 1999; Kanka et al. 2008; Turis 2009; Slovák et al. 2012; Kučera et al. 2013
<i>Dactylorhiza cordigera</i> subsp. <i>siculorum</i> (Soó) Soó ⁶³ (Syn.: <i>Orchis cordigera</i> subsp. <i>siculorum</i> Soó)	EC (RO), AC (RO)	East-Apuseni-Carpathian endemic	Soó 1967, 1980b; Popescu & Sanda 1998; Oprea 2005; Ciocârlan 2009; Hurdu et al. 2012a
<i>Dactylorhiza maculata</i> subsp. <i>schurii</i> (Klinge) Soó ⁶⁴ (Syn.: <i>Orchis maculata</i> var. <i>schurii</i> (Klinge) Paucă et Beldie)	EC (UA, RO), SC (RO), AC (RO), Tr (RO)	endemic to Eastern, Southern & Apuseni Carpathians and Transylvanian Basin	Soó 1967; Paucă & Beldie 1972; Čornej 2011; Hurdu 2014 in litt.
<i>Daphne arbuscula</i> Čelak.	WC (SK)	West-Carpathian endemic (Muránska planina Mts)	Erdelská & Turis 1996
<i>Delphinium elatum</i> subsp. <i>nacladense</i> (Zapał.) Holub ⁶⁵ (Syn.: <i>D. nacladense</i> Zapał.)	EC (PL, UA, RO), SC (RO)	East-South-Carpathian endemic	Čopyk 1976; Starmühler 1996b; Oprea 2005; Kobiv et al. 2007a; Chorney et al. 2008; Mitka et al. 2008; Novikov 2013b
<i>Delphinium oxysepalum</i> Borbás et Pax	WC (SK, PL)	West-Carpathian endemic	Kliment 1999
<i>Delphinium simonkaianum</i> Pawł.	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Morariu & Beldie 1976; Beldie 1977; Dihoru & Pârnu 1987; Oprea 2005
<i>Dianthus callizonus</i> Schott et Kotschy	SC (RO)	South-Carpathian endemic (Piatra Craiului Mts)	Morariu & Beldie 1976; Coldea 1997; Ciocârlan 2009; Dihoru & Negrean 2009; Sârbu et al. 2013
<i>Dianthus carthusianorum</i> subsp. <i>tenuifolius</i> (Schur) Hegi ⁶⁶ (Syn.: <i>D. tenuifolius</i> Schur)	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Kuz'mina 2004; Fedorončuk & Čornej 2005; Oprea 2005; Čornej 2011
<i>Dianthus giganteus</i> subsp. <i>banaticus</i> (Heuff.) Tutin (Syn.: <i>D. banaticus</i> (Heuff.) Borbás)	SC (RO, SRB)	South-Carpathian endemic	Dihoru & Pârnu 1987; Oprea 2005; Hurdu et al. 2012a, b
<i>Dianthus glacialis</i> subsp. <i>gelidus</i> (Schott, Nyman et Kotschy) Tutin (Syn.: <i>D. gelidus</i> Schott., Nyman et Kotschy)	EC (RO), SC (RO)	East-South-Carpathian endemic	Dihoru & Pârnu 1987; Oprea 2005; Hurdu et al. 2012a
<i>Dianthus henteri</i> Heuff. ex Griseb. et Schenk	SC (RO)	South-Carpathian endemic	Dihoru & Pârnu 1987; Popescu et al. 2003; Oprea 2005; Ciocârlan 2009; Sârbu et al. 2013
<i>Dianthus nitidus</i> Waldst. et Kit. (Syn.: <i>D. nitidus</i> Waldst. et Kit. subsp. <i>nitidus</i>)	WC (SK, †PL)	West-Carpathian endemic	Kliment 1999; Kmeťová 2012
<i>Dianthus praecox</i> Willd. ex Spreng. subsp. <i>praecox</i> ^{67, 68} (Syn.: <i>D. plumarius</i> subsp. <i>praecox</i> (Willd. ex Spreng.) Domin)	WC (SK, PL)	West-Carpathian endemic	Kmeťová 1985, 2012
<i>Dianthus praecox</i> subsp. <i>lummitzeri</i> (Wiesb.) Kmeťová ⁶⁹ (Syn.: <i>D. lummitzeri</i> Wiesb.; <i>D. plumarius</i> subsp. <i>lummitzeri</i> (Wiesb.) Domin)	WC (A, CZ, SK)	West-Carpathian endemic	Baksay 1972; Kmeťová 1985, 2012; Somogyi et al. 2012
<i>Dianthus praecox</i> subsp. <i>pseudopraecox</i> (Novák) Kmeťová ex Dostál ^{70, 71} (Syn.: <i>D. hungaricus</i> subsp. <i>pseudopraecox</i> (Novák) Kmeťová ex Futák)	WC (SK, HU)	West-Carpathian endemic	Kmeťová 1985, 2012
<i>Dianthus spiculifolius</i> Schur ⁷² (Syn.: <i>D. kitaibelii</i> subsp. <i>spiculifolius</i> (Schur) Novák; <i>D. petraeus</i> subsp. <i>spiculifolius</i> (Schur) Ciocârlan)	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Morariu & Beldie 1976; Negrean & Oltean 1989; Fedorončuk & Diduch 2002b; Oprea 2005; Holobiuc et al. 2009

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Doronicum carpaticum</i> (Griseb. et Schenk) Nyman ⁷³	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Álvarez Fernández 2003; Pachswöll 2013
<i>Draba doreri</i> Heuff. ⁷⁴	SC (RO)	South-Carpathian endemic	Sârbru & Lupu 1989; Ion 2012; Catană et al. 2013
<i>Draba haynaldii</i> Stur	EC (RO), SC (RO)	East-South-Carpathian endemic	Beldie 1979; Oprea 2005; Dihoru & Negrean 2009
<i>Draba kotschyi</i> Stur ⁷⁵	EC (RO), SC (RO)	East-South-Carpathian endemic	Jalas et al. 1996; Oprea 2005; Ciocârlan 2009
<i>Draba lasiocarpa</i> subsp. <i>klasterskyi</i> (Chrtek) Chrtek	WC (SK)	West-Carpathian endemic (Slovenský kras Mts)	Peniašteková & Kliment 2002
<i>Draba simonkaiana</i> Jáv. ⁷⁶ (Syn.: <i>D. stellata</i> subsp. <i>simonkaiana</i> (Jáv.) Ciocârlan)	SC (RO)	South-Carpathian endemic	Dihoru & Pârvu 1987; Oprea 2005; Ciocârlan 2009; Dihoru & Negrean 2009; Sârbru et al. 2013
<i>Erigeron hungaricus</i> (Vierh.) Pawł. ⁷⁷ (Syn.: <i>E. nanus</i> Schur non Nutt., nom. illeg.)	WC (SK, PL), EC (RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999; Oprea 2005; Mirek & Piękoś-Mirkowa 2008b; Ciocârlan 2009
<i>Eritrichium jankae</i> Simonk. ⁷⁸ (Syn.: <i>E. nanum</i> subsp. <i>jankae</i> (Simonk.) Jáv.)	EC (RO), SC (RO)	East-South-Carpathian endemic	Oprea 2005; Ciocârlan 2009; Șuteu 2012; Sârbru et al. 2013
<i>Erysimum hungaricum</i> Zapal. ^{79, 80}	EC (RO)	East-Carpathian endemic (Maramureș Mts: Mt. Lostun Mic)	Kobiv et al. 2007a; Kobiv 2010; Sârbru et al. 2013
<i>Erysimum pieninicum</i> (Zapal.) Pawł.	WC (PL)	West-Carpathian endemic (Pieniny Mts)	Maciejewska-Rutkowska et al. 2007; Korzeniak 2008
<i>Erysimum wahlenbergii</i> (Asch. et Engl.) Borbás	WC (SK, PL)	West-Carpathians endemic	Kliment 1999; Michalková 2002
<i>Erysimum witmannii</i> Zaw. ⁸¹	WC (SK, PL, HU), EC (UA, RO), SC (RO)/?†AC	West-East-South-Carpathian endemic	Kliment 1999; Michalková 2002
<i>Erysimum witmannii</i> Zaw. subsp. <i>witmannii</i> ^{82, 83, 84} (Syn.: <i>E. baumgartenianum</i> Schur)	WC (SK, PL), EC (RO), SC (RO)/?†AC (RO)	West-East-South-Carpathian endemic	Nyárády 1955; Borza 1964; Tomšovic 1988; Kliment 1999; Ančev & Polatschek 2006
<i>Erysimum witmannii</i> subsp. <i>pallidiflorum</i> (Szépl. ex Jáv.) Soó (Syn.: <i>E. pallidiflorum</i> Szépl. ex Jáv.)	WC (SK, HU)	West-Carpathian endemic	Michalková 1999, 2002
<i>Erysimum witmannii</i> subsp. <i>transsilvanicum</i> (Schur) P. W. Ball ^{85, 86} (Syn.: <i>E. transsilvanicum</i> Schur; <i>E. czetzianum</i> Schur)	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Nyárády 1955; Popescu et al. 2003; Kobiv 2010; Oprea & Sîrbu 2012, 2013; Vojtkó et al. 2012
<i>Euphorbia carpatica</i> Wol. ^{87, 88} (Syn.: <i>Tithymalus carpaticus</i> (Wol.) Á. Löve et D. Löve; <i>T. jasiewiczii</i> Chrtek et Křisa)	EC (UA, RO)	East-Carpathian endemic	Čopyk 1976; Mirek et al. 2002; Dihoru & Negrean 2009; Čornej 2011
<i>Euphorbia sojakii</i> (Chrtek et Křisa) Dubovik (Syn.: <i>E. austriaca</i> subsp. <i>sojakii</i> Chrtek et Křisa; <i>Tithymalus sojakii</i> (Chrtek et Křisa) Holub)	EC (SK, PL, UA)	East-Carpathian endemic	Kliment 1999
<i>Euphrasia exaristata</i> Smejkal ⁸⁹	WC (SK, PL)	West-Carpathian endemic (Západné Tatry Mts: Červené vrchy massif)	Králik 1997; Staszkievicz 2009
<i>Euphrasia slovaca</i> (Yeo) Holub subsp. <i>slovaca</i> (Syn.: <i>E. arctica</i> subsp. <i>slovaca</i> Yeo)	WC (CZ, SK, PL), EC (UA, RO)	West-East-Carpathian endemic	Králik 1997; Kliment 1999; Oprea 2005; Čornej 2011; Sârbru et al. 2013
<i>Euphrasia stipitata</i> Smejkal	WC (SK)	West-Carpathian endemic (Krivánska Fatra Mts)	Králik 1997; Kliment 1999
<i>Euphrasia tatrae</i> Wettst. ⁹⁰ (Syn.: <i>E. minima</i> subsp. <i>tatrae</i> (Wettst.) Hayek; <i>E. minima</i> var. <i>tatrae</i> (Wettst.) Pawł.)	WC (SK, PL), EC (UA, RO), ?SC (RO)	West-East-Carpathian endemic	Smejkal 1963; Mihoková & Mikoláš 1994; Smejkal & Čefovský 1999
<i>Ferula sadlerana</i> Ledeb. ^{91, 92}	WC (SK, HU), AC (RO)	West-Apuseni-Carpathian subendemic	Lendvay & Kalapos 2014
<i>Festuca amethystina</i> subsp. <i>orientalis</i> Krajina ⁹³ (Syn.: <i>F. inarmata</i> Schur)	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Cvelev 1972, 1974, 1976; Prokudin et al. 1977; Oprea 2005; Sârbru et al. 2013

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Festuca bucegiensis</i> Markgr.-Dann.	SC (RO)	South-Carpathian endemic	Oprea 2005; Ciocârlan 2009; Sârbu et al. 2013
<i>Festuca carpatica</i> F. Dietr. (Syn.: <i>F. pseudolaxa</i> Schur; <i>Leucopoa carpatica</i> (F. Dietr.) H. Scholz)	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999
<i>Festuca gautieri</i> subsp. <i>lutea</i> (Hack.) Foggi et Signorini (Syn.: <i>F. gautieri</i> subsp. <i>lutea</i> (Hack.) Ciocârlan; <i>F. scoparia</i> subsp. <i>lutea</i> (Hack.) Beldie)	EC (RO)	East-Carpathian endemic (Hășmaș Mts)	Morariu & Beldie 1976; Oprea 2005; Sârbu et al. 2013
<i>Festuca nitida</i> subsp. <i>flaccida</i> (Schur) Markgr.-Dann.	EC (RO), SC (RO)	East-South-Carpathian endemic	Oprea 2005; Ciocârlan 2009; Sârbu et al. 2013
<i>Festuca pachyphylla</i> Degen ex Nyár. (Syn.: <i>F. rupicola</i> subsp. <i>pachyphylla</i> (Degen ex Nyár.) Beldie; <i>F. stricta</i> subsp. <i>rumelica</i> Foggi et Petrova)	SC (RO)	South-Carpathian endemic	Markgraf-Dannenberg 1980; Ciocârlan 2009; Hurdu et al. 2012a
<i>Festuca porcii</i> Hack.	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Beldie 1972; Čopyk 1976; Oprea 2005; Kobiv et al. 2007a
<i>Festuca saxatilis</i> Schur ⁹⁴ (Syn.: <i>F. rupicola</i> subsp. <i>saxatilis</i> (Schur) Rauschert; <i>F. rupicola</i> subsp. <i>saxatilis</i> (Schur) Beldie, nom. illeg.; <i>F. stricta</i> subsp. <i>saxatilis</i> (Host) Foggi et Signorini)	EC (SK, UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Kliment 1999; Oprea 2005
<i>Festuca tatrae</i> (Czakó) Degen ⁹⁵ (Syn.: <i>F. amethystina</i> subsp. <i>tatrae</i> (Czakó) Soó)	WC (SK, PL)	West-Carpathian endemic	Kliment 1999; Ciocârlan 2009; Čornej 2011
<i>Festuca versicolor</i> Tausch subsp. <i>versicolor</i> ⁹⁶	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian subendemic	Kliment 1999; Oprea 2005
<i>Festuca versicolor</i> subsp. <i>dominii</i> Krajina	EC (RO)	East-Carpathian endemic (Rodna Mts)	Dihoru & Pârnu 1987; Oprea 2005; Ciocârlan 2009
<i>Galium abaujense</i> Borbás	WC (SK, HU)	West-Carpathian endemic	Kliment 1999
<i>Galium album</i> subsp. <i>suberectum</i> (Klokov) Michalk. ^{97, 98} (Syn.: <i>G. suberectum</i> Klokov)	WC (SK), EC (UA)	West-East-Carpathian endemic	Klokov 1961; Michalková 1993; Gynda 2004
<i>Galium baillonii</i> D. Brândză	SC (RO)	South-Carpathian endemic	Dihoru & Pârnu 1987; Popescu et al. 2003; Oprea 2005; Ciocârlan 2009
<i>Galium kitaibelianum</i> Schult. et Schult. f. ⁹⁹	EC (RO), SC (RO), AC	East-South-Apuseni-Carpathian subendemic	Oprea 2005; Hurdu et al. 2012b
<i>Galium transcarpaticum</i> Stojko et Tasenk.	EC (UA)	East-Carpathian endemic	Tasenkevych 2003b; Gynda 2004; Kobiv 2010; Čornej 2011
<i>Genista tinctoria</i> subsp. <i>oligosperma</i> (Andrae) Jáv. ¹⁰⁰ (Syn.: <i>G. oligosperma</i> (Andrae) Simonk.; <i>G. rupestris</i> Schur)	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Čopyk 1976; Morariu & Beldie 1976; Oprea 2005; Ciocârlan 2009; Čornej 2011
<i>Gentiana cruciata</i> subsp. <i>phlogifolia</i> (Schott et Kotschy) Tutin ¹⁰¹ (Syn.: <i>G. phlogifolia</i> Schott et Kotschy)	EC (RO), SC (RO)	East-South-Carpathian endemic	Morariu & Beldie 1976; Strid & Tan 1991; Oprea 2005; Ciocârlan 2009
<i>Gentiana laciniata</i> Kit. ex Kanitz ¹⁰²	EC (UA)	East-Carpathian endemic	Rybczyński et al. 2014
<i>Gentianella amarella</i> subsp. <i>reussii</i> (Tocl) Holub (Syn.: <i>Gentiana reussii</i> Tocl)	WC (SK)	West-Carpathian endemic	Kliment 1999; Šípošová et al. 2004b
<i>Gentianella fatrae</i> (Borbás) Holub (Syn.: <i>G. austriaca</i> subsp. <i>fatrae</i> (Borbás) Á. Löve et D. Löve)	WC (SK)	West-Carpathian endemic	Kliment 1999; Šípošová et al. 2004a
<i>Gentianella lutescens</i> subsp. <i>tatrae</i> (Ronniger) Holub	WC (SK)	West-Carpathian endemic	Kliment 1999; Mirek & Piękoś-Mirkowa 2010
<i>Gypsophila petraea</i> (Baumg.) Rchb. ¹⁰³ (Syn.: <i>G. transsylvanica</i> Spreng.)	EC (RO), SC (RO)	East-South-Carpathian endemic	Morariu & Beldie 1976; Jalas & Suominen 1986; Oprea 2005; Hurdu et al. 2012b

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Helictotrichon decorum</i> (Janka) Henrard	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Morariu & Beldie 1976; Oprea 2005; CBIS 2008 (http://www.carpates.org/cbisec/bot.php?id=1222)
<i>Hepatica transsilvanica</i> Fuss	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Morariu & Beldie 1976; Oprea 2005
<i>Heracleum carpathicum</i> Porcius	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Čopyk 1976; Oprea 2005; Kobiv et al. 2007a; Ciocârlan 2009; Dihoru & Negrean 2009
<i>Heracleum sphondylium</i> subsp. <i>transsilvanicum</i> (Schur) Brummitt (Syn.: <i>H. transsilvanicum</i> Schur; <i>H. palmatum</i> Baumg.)	EC (UA, RO), SC (RO), AC	East-South-Apuseni-Carpathian endemic	Čopyk 1976; Oprea 2005; Čornej 2011; Sârbu et al. 2013
<i>Hesperis matronalis</i> subsp. <i>schurii</i> Soó ¹⁰⁴ (Syn.: <i>H. oblongifolia</i> Schur; <i>H. matronalis</i> subsp. <i>oblongifolia</i> (Schur) F. Dvořák)	SC (RO)	South-Carpathian endemic	Morariu & Beldie 1976; Beldie 1977; Ball 1993; Ciocârlan 2009
<i>Hesperis matronalis</i> subsp. <i>vrabelyiana</i> (Schur) Soó (Syn.: <i>H. vrabelyiana</i> (Schur) Borbás)	WC (HU)	West-Carpathian endemic (Bükk Mts)	Dvořák 1968; Soó 1968; Ball 1993; Šeffler et al. 2010
<i>Hesperis slovacica</i> (F. Dvořák) F. Dvořák ¹⁰⁵ (Syn.: <i>H. dinarica</i> subsp. <i>slovaca</i> F. Dvořák)	WC (SK)	West-Carpathian endemic (Nízke Tatry Mts)	Dvořák 1963, 1968; Zahradníková et al. 2002; Šípošová et al. 2004b
<i>Hieracium abietogenum</i> Nyár. ex Szeląg ¹⁰⁶	SC (RO)	South-Carpathian endemic (Retezat Mts)	Szeląg 2003a, 2006b
<i>Hieracium austrotatricum</i> Szeląg	WC (SK)	West-Carpathian endemic (Nízke Tatry Mts)	Ronikier & Szeląg 2008; Ilnicki & Szeląg 2011
<i>Hieracium bohatschianum</i> Zahn	SC (RO)	South-Carpathian endemic (Almajului Mts: Mt. Treskovat)	Ilnicki & Szeląg 2011; Szeląg 2011
<i>Hieracium borbasii</i> R. Uechtr. ¹⁰⁷ (Syn.: <i>H. sparsum</i> subsp. <i>borbasii</i> (R. Uechtr.) Zahn)	SC (RO)	South-Carpathian endemic (Retezat Mts)	Szeląg 2006b; Ilnicki & Szeląg 2011
<i>Hieracium borsanum</i> Mráz	EC (RO)	East-Carpathian endemic (Rodna Mts)	Mráz 2001b, 2003a
<i>Hieracium carpathicum</i> Besser subsp. <i>carpathicum</i>	WC (SK, PL)	West-Carpathian endemic	Zahn 1937; Chrtek jr. 2014 in litt.; cf. Piękoś-Mirkowa & Mirek 2003; Šeffler et al. 2010
<i>Hieracium cernaeglavae</i> (Hruby et Zahn) Mráz (Syn.: <i>H. rohacsense</i> subsp. <i>cernaeglavae</i> (Hruby et Zahn) Zahn)	EC (UA, RO)	East-Carpathian endemic	Mráz 2002
<i>Hieracium coldei</i> Szeląg (Syn.: <i>H. sparsum</i> subsp. <i>coldei</i> (Szeląg) Greuter)	EC (RO), SC (RO)	East-South-Carpathian endemic	Szeląg 2006a, b, 2012
<i>Hieracium crassipedipilum</i> (Pawl. et Zahn) Chrtek f.	WC (SK, PL)	West-Carpathian endemic	Chrtek jr. & Marhold 1998; Štorchová et al. 2002; Chrtek jr. et al. 2004a
<i>Hieracium czeremosense</i> Woł. et Zahn	EC (UA)	East-Carpathian endemic	Zahn 1936; Šljakov 1989; Szeląg 2007
<i>Hieracium dacicum</i> R. Uechtr.	SC (RO)	South-Carpathian endemic (Retezat Mts)	Zahn 1938; Szeląg 2014 in litt.
<i>Hieracium decipienteforme</i> (Woł. et Zahn) Schljakov	EC (UA)	East-Carpathian endemic	Chrtek jr. 1997; Chrtek jr. 2014 in litt.
<i>Hieracium deylii</i> Mráz (Syn.: <i>H. pietroszense</i> subsp. <i>deylii</i> (Mráz) Greuter)	EC (UA)	East-Carpathian endemic	Mráz 2003a
<i>Hieracium fagarasense</i> (Nyár. et Zahn) Nyár. ¹⁰⁸ (Syn.: <i>H. sparsum</i> subsp. <i>fagarasense</i> Nyár. et Zahn)	SC (RO)	South-Carpathian endemic (Făgăraș Mts)	Szeląg 2006b; Szeląg 2014 in litt.; cf. Beldie 1979, Oprea 2005, Ciocârlan 2009, Sârbu et al. 2013
<i>Hieracium filarszkyi</i> Jáv. et Zahn	SC (RO)	South-Carpathian endemic (Retezat Mts)	Zahn 1938; Szeląg 2014 in litt.
<i>Hieracium fritzeiforme</i> Zahn	SC (RO)	South-Carpathian endemic (Retezat Mts)	Zahn 1938; Szeląg 2014 in litt.
<i>Hieracium hryniawiense</i> Woł. (Syn.: <i>H. raddeanum</i> subsp. <i>hryniawiense</i> (Woł.) Greuter)	EC (UA)	East-Carpathian endemic	Šljakov 1989; Mráz 2003b; Szeląg 2007
<i>Hieracium jankae</i> R. Uechtr.	SC (RO)	South-Carpathian endemic (Almajului Mts)	Mráz & Szeląg 2004; Szeląg 2014 in litt.

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Hieracium jarzabczynum</i> (Pawl. et Zahn) Mráz et Chrtek f.	WC (SK, PL)	West-Carpathian endemic	Chrtek jr. & Mráz 2007; Chrtek jr. et al. 2007
<i>Hieracium kotschyanum</i> Heuff. (Syn.: <i>H. kotschyanum</i> Heuff. subsp. <i>kotschyanum</i> ; <i>H. sparsum</i> subsp. <i>kotschyanum</i> (Heuff.) Zahn)	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Mráz & Szeląg 2004; Szeląg 2006b; Ilnicki & Szeląg 2011
<i>Hieracium krivanense</i> (Wol. et Zahn) Schljakov	WC (SK, PL)	West-Carpathian endemic (Tatry Mts)	Chrtek jr. & Marhold 1998; Štorchová et al. 2002; Chrtek jr. et al. 2004a
<i>Hieracium lingelsheimii</i> Pax	WC (SK)	West-Carpathian endemic (Nízke Tatry Mts)	Schuhwerk & Lippert 1999; Chrtek jr. et al. 2004a
<i>Hieracium lomnicense</i> Wol.	EC (UA)	East-Carpathian endemic	Šljakov 1989; Szeląg 2007
<i>Hieracium lubricicaule</i> (Zahn) Borza (Syn.: <i>H. sparsum</i> subsp. <i>lubricicaule</i> Zahn)	SC (RO)	South-Carpathian endemic (Retezat Mts, Țarcu Mts)	Szeląg 2003b, 2006b; Mráz & Szeląg 2004
<i>Hieracium magocsyanum</i> Jáv. (Syn.: <i>H. sparsum</i> subsp. <i>magocsyanum</i> (Jáv.) Zahn)	SC (RO)	South-Carpathian endemic (Retezat Mts, Țarcu Mts)	Mráz & Szeląg 2004; Szeląg 2006b; Ilnicki & Szeląg 2011
<i>Hieracium mirekii</i> Szeląg	SC (RO)	South-Carpathian endemic (Retezat Mts)	Szeląg 2006b; Ilnicki & Szeląg 2011
<i>Hieracium mitkai</i> Szeląg (Syn.: <i>H. kotschyanum</i> subsp. <i>longidentatum</i> Nyár. ex Szeląg; <i>H. sparsum</i> subsp. <i>longidentatum</i> (Nyár. ex Szeląg) Greuter)	SC (RO)	South-Carpathian endemic (Retezat Mts)	Szeląg 2003a, 2006b
<i>Hieracium mlinicae</i> (Hruby et Zahn) Chrtek f. et Mráz (Syn.: <i>H. nigrescens</i> subsp. <i>mlinicae</i> Hruby et Zahn)	WC (SK, PL)	West-Carpathian endemic (Tatry Mts)	Zahn 1927; Chrtek jr. et al. 2004b; Chrtek jr. & Mráz 2007
<i>Hieracium napaeum</i> Zahn	EC (RO), SC (RO), AC	East-South-Apuseni-Carpathian endemic	Zahn 1936; Szeląg 2014 in litt.
<i>Hieracium negoiense</i> (Răvărut et Nyár.) Soó ¹⁰⁹ (Syn.: <i>Crepis negoiensis</i> Răvărut et Nyár.)	SC (RO)	South-Carpathian endemic (Retezat Mts)	Morariu & Beldie 1976; Oprea 2005; Sârbu et al. 2013
<i>Hieracium nigrescens</i> subsp. <i>koprovanum</i> Rech. f. et Zahn	WC (SK, PL)	West-Carpathian endemic (Tatry Mts, Nízke Tatry Mts)	Zahn 1927; Mráz 2001b, 2002; Chrtek jr. et al. 2004b, 2007
<i>Hieracium nigrilacus</i> Nyár.	SC (RO)	South-Carpathian endemic (Retezat Mts)	Szeląg 2006b
<i>Hieracium ostii-bucurae</i> Nyár. ex Szeląg ^{110, 111} (Syn.: <i>H. longifoliosum</i> Nyár. ex Szeląg; <i>H. sparsum</i> subsp. <i>sparsiforme</i> (Degen et Zahn) Greuter)	SC (RO)	South-Carpathian endemic	Szeląg 2003a, 2006b; Mráz & Szeląg 2004
<i>Hieracium palenicae</i> Rech. f. et Zahn	WC (SK)	West-Carpathian endemic (Západné Tatry Mts)	Zahn 1937; Procházka & Chrtek jr. 1999; Chrtek jr. 2014 in litt.
<i>Hieracium paltinae</i> Jáv. et Zahn	SC (RO)	South-Carpathian endemic (Retezat Mts)	Szeląg 2014 in litt.; cf. Zahn 1938
<i>Hieracium pawlowskianum</i> Nyár. (Syn.: <i>H. riumarense</i> Nyár. ex Szeląg; <i>H. tomasiforme</i> (Nyár.) Nyár.; <i>H. sparsum</i> subsp. <i>nigrovirenticeps</i> (Nyár. et Zahn) Greuter)	SC (RO)	South-Carpathian endemic (Retezat Mts, Țarcu Mts)	Szeląg 2003a, 2004a, 2006b
<i>Hieracium paxianum</i> Nyár. et Zahn	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Mráz et al. 2005
<i>Hieracium perfoliosum</i> Nyár. ex Szeląg	SC (RO)	South-Carpathian endemic (Godeanu Mts)	Szeląg 2003a, 2006b
<i>Hieracium pietroszense</i> Degen et Zahn ¹¹²	EC (RO)	East-Carpathian endemic (Rodna Mts)	Mráz 2003a
<i>Hieracium pinetophilum</i> (Degen et Zahn) Chrtek f.	WC (SK, PL)	West-Carpathian endemic	Chrtek jr. & Marhold 1998; Štorchová et al. 2002; Chrtek jr. et al. 2004a
<i>Hieracium pisatureense</i> Nyár.	SC (RO)	South-Carpathian endemic (Retezat Mts)	Szeląg 2003b, 2006b; Szeląg 2014 in litt.; cf. Zahn 1938
<i>Hieracium pocuticum</i> Wol.	EC (UA, RO)	East-Carpathian endemic	Šljakov 1989; Szeląg 2007; cf. Tasenkevyc 2003b

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Hieracium pojoritense</i> Woł.	EC (RO)	East-Carpathian endemic	Ștefureac & Tăcină 1979; Mráz 2003b; Mráz & Szeląg 2004; Mráz & Paule 2006; Szeląg 2006b, 2007
<i>Hieracium polyphyllobasis</i> (Nyár. et Zahn) Szeląg (Syn.: <i>H. sparsum</i> subsp. <i>polyphyllobasis</i> (Nyár. et Zahn) Greuter)	SC (RO)	South-Carpathian endemic (Retezat Mts)	Szeląg 2006b; Ilnicki & Szeląg 2011
<i>Hieracium porphyriticum</i> A. Kern. (Syn.: <i>H. sparsum</i> subsp. <i>porphyriticum</i> (A. Kern.) Zahn)	SC (RO), AC (RO)	South-Apuseni-Carpathian endemic	Szeląg 2006b; Ilnicki & Szeląg 2011; Szeląg 2014 in litt.
<i>Hieracium prassivae</i> Zahn	WC (SK)	West-Carpathian endemic (Vysoké Tatry Mts, Nízke Tatry Mts)	Šípošová et al. 2004b
<i>Hieracium pseudostygium</i> Woł. (Syn.: <i>H. nigritum</i> subsp. <i>pseudostygium</i> (Woł.) Zahn)	EC (UA)	East-Carpathian endemic	Šljakov 1989; Szeląg 2007; cf. Čornej 2011
<i>Hieracium rapunculoidiforme</i> Woł. et Zahn ¹¹³	EC (UA, ?RO)	East-Carpathian endemic	Zahn 1911; Šljakov 1989
<i>Hieracium ratezaticum</i> (Nyár. et Zahn) Mráz	SC (RO)	South-Carpathian endemic (Retezat Mts)	Mráz 2001b; Mráz & Szeląg 2004
<i>Hieracium rohacsense</i> Kit.	WC (SK, PL)	West-Carpathian endemic	Mráz 2001a, 2002, 2005a; Šípošová et al. 2004a
<i>Hieracium scitulum</i> Woł. ¹¹⁴	EC (UA)	East-Carpathian endemic	Šljakov 1989; Chrtek jr. 2004; Szeląg 2007
<i>Hieracium silesiacum</i> E. Krause ¹¹⁵	WC (SK, PL)	West-Carpathian subendemic	Chrtek jr. 1996; Chrtek jr. et al. 2002; Szeląg 2004b, 2006b; Mráz 2005b
<i>Hieracium slovacum</i> Chrtek f.	WC (SK)	West-Carpathian endemic (Belianske Tatry Mts)	Chrtek jr. & Marhold 1998; Štorchová et al. 2002; Chrtek jr. et al. 2004a
<i>Hieracium subsinuatum</i> Borbás (Syn.: <i>H. subserratosinuatum</i> Zahn, nom. illeg.)	WC (SK)	West-Carpathian endemic (Tatry Mts)	Zahn 1937; Kliment 1999; Procházka & Chrtek jr. 1999; Chrtek jr. 2014 in litt.
<i>Hieracium telekianum</i> Boros et Lengyel (Syn.: <i>H. sparsum</i> subsp. <i>telekianum</i> (Boros et Lengyel) Greuter)	EC (RO)	East-Carpathian endemic (Harghita Mts)	Mráz & Szeląg 2004; Szeląg 2006b; cf. Beldie 1977, Ciocârlan 2009, Sârbu et al. 2013
<i>Hieracium tomasae</i> (Nyár. et Zahn) Nyár. (Syn.: <i>H. sparsum</i> subsp. <i>tomasae</i> Nyár. et Zahn; <i>H. sparsum</i> var. <i>tomasae</i> (Nyár. et Zahn) Ciocârlan)	SC (RO)	South-Carpathian endemic (Țarcu Mts: Mt. Tomeasa)	Mráz & Szeląg 2004; Szeląg 2006b; Ilnicki & Szeląg 2011; cf. Dihoru & Pârvu 1987
<i>Hieracium tomosense</i> Simonk.	EC (RO), SC (RO)	East-South-Carpathian endemic	Ilnicki & Szeląg 2011; Szeląg 2013
<i>Hieracium tubulare</i> Nyár. ¹¹⁶ (Syn.: <i>H. sparsum</i> var. <i>tubulare</i> (Nyár.) Ciocârlan; <i>H. sparsum</i> subsp. <i>tubulatum</i> (Zahn) Greuter)	SC (RO)	South-Carpathian endemic (Retezat Mts)	Mráz & Szeląg 2004; Szeląg 2006a, b; Ilnicki & Szeląg 2011
<i>Hieracium ukierniae</i> Woł. et Zahn	EC (UA)	East-Carpathian endemic	Šljakov 1989; Szeląg 2014 in litt.
<i>Hieracium vapenicanum</i> (Lengyel et Zahn) Chrtek f. et Mráz	WC (SK, PL)	West-Carpathian endemic	Zahn 1927; Chrtek jr. et al. 2004b, 2007; Chrtek jr. & Mráz 2007
<i>Hieracium virgicale</i> Nägeli et Peter ¹¹⁷	WC (SK, PL, HU)	West-Carpathian endemic	Zahn 1930; Chrtek jr. et al. 2004a
<i>Hieracium worochtae</i> Woł.	EC (UA)	East-Carpathian endemic	Zahn 1938; Šljakov 1989; Szeląg 2007
<i>Hieracium zajacii</i> Szeląg	WC (SK)	West-Carpathian endemic (Veľká Fatra Mts)	Szeląg 2010
<i>Hieracium zanogae</i> Pax	SC (RO)	South-Carpathian endemic (Retezat Mts)	Szeląg 2006b
<i>Hylotelephium argutum</i> (Haw.) Holub (Syn.: <i>Sedum carpaticum</i> G. Reuss; <i>S. telephium</i> subsp. <i>fabaria</i> (W. D. J. Koch) Kirschl.)	WC (†CZ, SK, PL), EC (SK, UA, RO), SC (RO), AC (RO)	West-East-South-Apuseni-Carpathian (pan-Carpathian) endemic	Gulich 1984; Kliment 1999
<i>Jovibarba globifera</i> subsp. <i>preissiana</i> (Domin) Holub ¹¹⁸ (Syn.: <i>J. preissiana</i> (Domin) Omelczuk et Czopik)	WC (SK, PL), EC (UA, RO)	West-East-Carpathian subendemic	Letz 1998; Kliment 1999

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Jurinea transylvanica</i> (Spreng.) Simonk. ¹¹⁹ (Syn.: <i>J. mollis</i> subsp. <i>transylvanica</i> (Spreng.) Hayek)	Tr (RO)/EC (RO), SC (RO), AC (RO)	subendemic to Transylvanian Basin	Dihoru & Pârvu 1987; Oprea 2005
<i>Knautia kitaibelii</i> (Schult.) Borbás subsp. <i>kitaibelii</i> ^{120, 121}	WC (A, CZ, SK, PL)	West-Carpathian subendemic	Soják 1983b; Štěpánek 1985, 1997; Kliment 1999; Böhm & Facsar 2000; Kolář et al. 2009
<i>Knautia slovacica</i> Štěpánek	WC (SK)	West-Carpathian endemic	Štěpánek 1983, 1985
<i>Koeleria macrantha</i> subsp. <i>transsilvanica</i> (Schur) A. Nyár. ¹²² (Syn.: <i>K. transsilvanica</i> Schur; <i>K. macrantha</i> subsp. <i>transsilvanica</i> (Schur) Beldie, nom. illeg.)	EC (†UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Deyl 1934; Ghișa 1972; Soják 1983b; Popescu et al. 2003; Oprea 2005; Kricsfalusy & Budnikov 2007
<i>Koeleria tristis</i> Domin	WC (SK)	West-Carpathian endemic	Kliment 1999; Pečinka et al. 2006
<i>Lathyrus transsilvanicus</i> (Spreng.) Fritsch ^{123, 124, 125}	WC (SK, HU), EC (UA, RO), Tr (RO)/AC (RO)	subendemic to Western & Eastern Carpathians and Transylvanian Basin	Kliment 1999; Oprea 2005; Petrova & Vladimirov 2009; Proc' & Kiš 2009; Kobiv 2010; Tosheva et al. 2011; Marinov et al. 2014
<i>Leontodon kulczynskii</i> Popov (Syn.: <i>L. repens</i> Schur, nom. illeg.)	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Čopyk 1976; Hel'tman 1989; Oprea 2005; Čornej 2011
<i>Leucanthemopsis alpina</i> subsp. <i>tatrae</i> (Vierh.) Holub ¹²⁶ (Syn.: <i>L. tatrae</i> (Vierh.) Holub; <i>Chrysanthemum alpinum</i> f. <i>tatrae</i> Vierh.)	WC (SK, PL)	West-Carpathian endemic (Tatry Mts, Nízke Tatry Mts)	Holub 1977b; Kliment 1999; Mirek & Piękoś-Mirkowa 2009, 2010
<i>Leucanthemum rotundifolium</i> (Waldst. et Kit. ex Willd.) DC. ¹²⁷ (Syn.: <i>L. waldsteinii</i> (Sch. Bip.) Pouzar; <i>Chrysanthemum rotundifolium</i> Waldst. et Kit. ex Willd.)	WC (SK, PL), EC (SK, PL, UA), SC (RO), AC (RO)	West-East-South-Apuseni-Carpathian (pan-Carpathian subendemic	Zelený 1970; Čopyk 1976; Kliment 1999; Oprea 2005
<i>Linum extraaxilare</i> Kit. ¹²⁸ (Syn.: <i>L. perenne</i> subsp. <i>extraaxillare</i> (Kit.) Nyman)	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian subendemic	Kliment 1999; Oprea 2005; Petrova & Vladimirov 2009; Vladimirov et al. 2011
<i>Linum uninerve</i> (Rochel) Jáv. ^{129, 130}	SC (RO)/EC (RO)	East-South-Carpathian subendemic	Dihoru & Pârvu 1987; Oprea 2005; Petrova 2011
<i>Luzula alpinopilosa</i> subsp. <i>obscura</i> S. Fröhner	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999; Oprea 2005; Mirek & Piękoś-Mirkowa 2010; Čornej 2011
<i>Melampyrum saxosum</i> Baumg. ^{131, 132} (Syn.: <i>M. sylvaticum</i> subsp. <i>saxosum</i> (Baumg.) P. Beauv.; <i>M. herbichii</i> Woł.)	EC (SK, PL, UA, RO), SC (RO)	East-South-Carpathian endemic	Štech & Drábková 2005; Těšitel & Štech 2007; Michalik & Mitka 2008; Těšitel et al. 2009
<i>Minuartia oxypetala</i> (Woł.) Kulcz. (Syn.: <i>M. verna</i> subsp. <i>oxypetala</i> (Woł.) G. Halliday)	EC (UA, RO)	East-Carpathian endemic	Pawłowski 1939; Beldie 1977; Fedorončuk & Diduch 2002c; Oprea 2005; Kobiv et al. 2007a; Kobiv 2010; Čornej 2011
<i>Minuartia pauciflora</i> (Kit. ex Kanitz) Dvořáková ¹³³ (Syn.: <i>M. zarenczyi</i> (Zapał.) Klokov)	WC (SK, PL), EC (UA)	West-East-Carpathian endemic	Dvořáková 2003
<i>Nigritella carpatica</i> (Zapał.) Teppner, E. Klein et M. Zagulskij (Syn.: <i>Gymnadenia carpatica</i> (Zapał.) Teppner et E. Klein)	EC (UA, RO)	East-Carpathian endemic	Teppner et al. 1994; Teppner & Klein 1998; Teppner 2004; Čornej 2009
<i>Noccaea banatica</i> (R. Uechtr.) F. K. Mey. ¹³⁴ (Syn.: <i>Thlaspi dacicum</i> subsp. <i>banaticum</i> (R. Uechtr.) Nyár.; <i>T. dacicum</i> subsp. <i>banaticum</i> (R. Uechtr.) Dvořáková, nom. illeg.)	SC (RO)	South-Carpathian endemic	Beldie 1977; Dihoru & Pârvu 1987; Popescu et al. 2003; Oprea 2005; Ciocârlan 2009; Dihoru & Negrean 2009; Sârbu et al. 2013
<i>Noccaea caeruleascens</i> subsp. <i>tatrensis</i> (Zapał.) Holub (Syn.: <i>Thlaspi caeruleascens</i> subsp. <i>tatrense</i> (Zapał.) Dvořáková)	WC (SK, PL)	West-Carpathian endemic	Kliment 1999; Hodálová & Mártonfi 2002
<i>Noccaea dacica</i> (Heuff.) F. K. Mey. subsp. <i>dacica</i> (Syn.: <i>Thlaspi dacicum</i> Heuff. subsp. <i>dacicum</i>)	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Morariu & Beldie 1976; Jalas et al. 1996; Tasenkevych 2003b; Oprea 2005; Ciocârlan 2009

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Noccaea jankae</i> (A. Kern.) F. K. Mey. ¹³⁵ (Syn.: <i>Thlaspi jankae</i> A. Kern.)	WC (SK, HU)	West-Carpathian endemic	Clapham & Akeyrod 1993; Kliment 1999; Hodálová & Mártonfi 2002
<i>Onosma pseudarenaria</i> Schur subsp. <i>pseudarenaria</i> ¹³⁶	Tr (RO)/EC (RO), AC (RO)	subendemic to Transylvanian Basin	Oprea 2005; Dihoru & Negrean 2009; Șuteu 2012
<i>Onosma viridis</i> (Borbás) Jáv. ^{137, 138} (Syn.: <i>Onosma tornensis</i> Jáv.)	WC (SK, HU), SC (RO), AC (RO), Tr (RO)	endemic to Western, Southern & Apuseni Carpathians and Transylvanian Basin	Grințescu & Nyárády 1960a; Kolarčík et al. 2010; Mártonfi et al. 2014
<i>Ophrys holubyana</i> Andras. ¹³⁹	WC (CZ, SK)	West-Carpathian endemic	Kliment 1999; Dítě 2014 in litt.
<i>Ornithogalum orthophyllum</i> subsp. <i>acuminatum</i> (Schur) Zahar.	SC (RO)	South-Carpathian endemic (Bârsei Mts: Postăvarul massif)	Dihoru & Pârvu 1987; Oprea 2005
<i>Oxytropis campestris</i> subsp. <i>tatrae</i> (Borbás) Dostál	WC (SK, PL), EC (RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999; Tasenkevich 2011
<i>Oxytropis carpatica</i> Uechtr. (Syn.: <i>O. jacquinii</i> subsp. <i>carpatica</i> (R. Uechtr.) Hausskn.)	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999; Oprea 2005
<i>Papaver corona-sancti-stephani</i> Zapał. (Syn.: <i>P. alpinum</i> subsp. <i>corona-sancti-stephani</i> (Zapał.) Borza; <i>P. pyrenaicum</i> subsp. <i>corona-sancti-stephani</i> (Zapał.) Borza)	EC (RO), SC (RO)	East-South-Carpathian endemic	Beldie 1977; Dihoru & Pârvu 1987; Oprea 2005; Ciocârlan 2009; Sârbu et al. 2013
<i>Papaver tatricum</i> (A. Nyár.) Ehrend. (Syn.: <i>P. alpinum</i> subsp. <i>tatricum</i> A. Nyár.)	WC (SK, PL)	West-Carpathian endemic	Kliment 1999
<i>Papaver tatricum</i> (A. Nyár.) Ehrend. subsp. <i>tatricum</i>	WC (SK, PL)	West-Carpathian endemic (Tatry Mts)	Bernátová 2002; Mirek & Piękoś-Mirkowa 2010
<i>Papaver tatricum</i> subsp. <i>fatraemagnae</i> Bernátová	WC (SK)	West-Carpathian endemic (Veľká Fatra Mts)	Bernátová 2002; Šipošová et al. 2004a; Kliment et al. 2008
<i>Pedicularis baumgartenii</i> Simonk. ¹⁴⁰	SC (RO)/EC (RO)	East-South-Carpathian endemic	Beldie 1977; Dihoru & Pârvu 1987; Oprea 2005; Ciocârlan 2009; Sârbu et al. 2013
<i>Peucedanum rochelium</i> Heuff. ^{141, 142}	Tr (RO)	subendemic to Transylvanian Basin	Boșcaiu 1965; Boșcaiu & Rațiu 1965; Dihoru & Pârvu 1987; Paucă-Comănescu & Negrean 1994; Jakab et al. 2008; Kovács 2011
<i>Phyteuma tetramerum</i> Schur	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Čopyk 1976; Morariu & Beldie 1976; Negrean & Oltean 1989; Oprea 2005; Čornej 2011
<i>Phyteuma vagneri</i> A. Kern. (Syn.: <i>P. spiciforme</i> Rochel)	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Dihoru & Pârvu 1987; Negrean & Oltean 1989; Oprea 2005
<i>Pilosella plaicensis</i> (Wol.) Soják (Syn.: <i>Hieracium plaicense</i> Wol.)	EC (UA)	East-Carpathian endemic	Szeląg 2007
<i>Pilosella ullepitschii</i> (Błocki) Szeląg ¹⁴³ (Syn.: <i>P. alpicola</i> subsp. <i>ullepitschii</i> (Błocki) Soják; <i>Hieracium alpicola</i> subsp. <i>ullepitschii</i> (Błocki) Zahn)	WC (SK, PL)/EC (RO), SC (RO)	West-East-South-Carpathian endemic	Szeląg 2008; Šingliarová et al. 2008, 2011a, b, 2013; Šingliarová & Mráz 2009; Ilnicki & Szeląg 2011
<i>Plantago atrata</i> subsp. <i>carpatica</i> (Soó) Soó	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999; Oprea 2005; Sârbu et al. 2013
<i>Poa babiogorensis</i> Bernátová, Májovský et Obuch	WC (PL)	West-Carpathian endemic (Beskid Żywiecki Mts: Mt. Babia Góra)	Bernátová et al. 1999, 2003a
<i>Poa carpatica</i> (V. Jirásek) Chopik ¹⁴⁴ (Syn.: <i>P. nemoralis</i> subsp. <i>carpatica</i> V. Jirásek)	WC (SK, PL), EC (UA)	West-East-Carpathian endemic	Čopyk 1976; Bernátová et al. 2006

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Poa granitica</i> Braun-Blanq.	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999
<i>Poa granitica</i> Braun-Blanq. subsp. <i>granitica</i>	WC (SK, PL)	West-Carpathian endemic (Tatry Mts)	Kliment 1999; Mirek & Piękoś-Mirkowa 2008c
<i>Poa granitica</i> subsp. <i>disparilis</i> (Nyár.) Nyár. ¹⁴⁵ (Syn.: <i>P. deylii</i> Chrtek et V. Jirásek)	EC (UA, RO), SC (RO)	East-South-Carpathian endemic	Ghişa & Beldie 1972; Beldie 1972; Oprea 2005; Ciocârlan 2009; Kobiv 2010; Sârbu et al. 2013
<i>Poa margilicola</i> Bernátová et Májovský	WC (SK)	West-Carpathian endemic (Veľká Fatra Mts: Mt. Borišov)	Bernátová & Májovský 1997; Bernátová et al. 2003b; Šipošová et al. 2004a
<i>Poa nobilis</i> Skalińska ¹⁴⁶	WC (SK, PL)	West-Carpathian endemic (Tatry Wysokie Mts)	Piękoś-Mirkowa et al. 1996; Kliment 1999; Piękoś-Mirkowa 2008
<i>Poa pannonica</i> subsp. <i>scabra</i> (Asch. et Graebn.) Soó ¹⁴⁷	WC (SK, HU), †EC (UA)/SC (RO)	West-South-Carpathian endemic	Kliment 1999; Oprea 2005; Dúbravková et al. 2010; Borhidi et al. 2012
<i>Poa rehmannii</i> (K. Richt.) Woł. ¹⁴⁸ (Syn.: <i>P. rehmannii</i> (K. Richt.) Szafer, nom. illeg.; <i>P. nemoralis</i> subsp. <i>rehmannii</i> (K. Richt.) Asch. et Graebn.)	EC (UA, RO)	East-Carpathian endemic	Čopyk 1976; Beldie 1979; Oprea 2005; Dihoru & Negrean 2009; Čornej 2011; Sârbu et al. 2013
<i>Poa sejuncta</i> Bernátová, Májovský et Obuch	WC (SK)	West-Carpathian endemic (Západné Tatry Mts: Mt. Osobitá)	Bernátová et al. 1999, 2003a
<i>Prangos carinata</i> Griseb. ex Degen ^{149, 150} (Syn.: <i>P. carinata</i> Griseb. ex Grecescu, nom. illeg.; <i>P. ferulacea</i> subsp. <i>carinata</i> (Griseb. ex Degen) Dihoru)	SC (RO)	South-Carpathian endemic (Almajului Mts: Porțile de Fier Gorge)	Morariu & Beldie 1976; Beldie 1977; Dihoru & Pârvu 1987; Oprea 2005; Dihoru & Negrean 2009
<i>Primula auricula</i> subsp. <i>hungarica</i> (Borbás) Soó ¹⁵¹	WC (SK, PL)	West-Carpathian subendemic	Soó 1964, 1980a; Simon 1992; Kliment 1999
<i>Primula auricula</i> subsp. <i>serratifolia</i> (Rochel) Jáv.	SC (RO, SRB)	South-Carpathian endemic	Beldie 1979; Popescu et al. 2003; Oprea 2005; Ciocârlan 2009; Dihoru & Negrean 2009
<i>Primula leucophylla</i> Pax ¹⁵² (Syn.: <i>P. elatior</i> subsp. <i>leucophylla</i> (Pax) Hesl.-Harr.f. ex W. W. Sm. et H. R. Fletcher)	EC (RO)	East-Carpathian endemic	Șuteu 2012; Șuteu et al. 2011, 2013; Hurdu et al. 2014 in litt.
<i>Primula wulfeniana</i> subsp. <i>baumgarteniana</i> (Degen et Moesz) Lüdi (Syn.: <i>P. baumgarteniana</i> Degen et Moesz)	SC (RO)	South-Carpathian endemic	Dihoru & Pârvu 1987; Oprea 2005; Dihoru & Negrean 2009; Negrean 2011; Hurdu et al. 2012a; Sârbu et al. 2013
<i>Pulmonaria rubra</i> subsp. <i>filarszkyana</i> (Jáv.) Domin ¹⁵³ (Syn.: <i>P. filarszkyana</i> Jáv.)	EC (UA, RO)	East-Carpathian endemic	Čopyk 1976; Beldie 1979; Oprea 2005; Čornej 2011; Ciocârlan 2009; Sârbu et al. 2013
<i>Pulsatilla slavica</i> G. Reuss ¹⁵⁴ (Syn.: <i>P. halleri</i> subsp. <i>slavica</i> (G. Reuss) Zamels)	WC (SK, PL)	West-Carpathian endemic	Goliašová 1985; Kliment 1999; Šipošová et al. 2004b; Ciocârlan 2009
<i>Pulsatilla subslavica</i> Futák ex Goliašová	WC (SK)	West-Carpathian endemic	Goliašová 1985; Kliment 1999
<i>Pyrola carpaticea</i> Holub et Křísa (Syn.: <i>P. rotundifolia</i> subsp. <i>carpaticea</i> (Holub et Křísa) Beldie et Váczy)	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999; Oprea 2005; Ciocârlan 2009
<i>Ranunculus altitatrencensis</i> Paclová et Murin	WC (SK)	West-Carpathian endemic (Vysoké Tatry Mts)	Paclová 1999
<i>Ranunculus carpaticeus</i> Herlich	EC (SK, UA, RO), SC (RO)	East-South-Carpathian endemic	Kliment 1999; Oprea 2005
<i>Ranunculus flabellifolius</i> Heuff. ex Rchb. ¹⁵⁵	SC (RO, SRB)	South-Carpathian endemic	Jalas & Suominen 1989; Stevanović et al. 1991; Dunkel 2011
<i>Ranunculus malinovskii</i> Elenevsky et Derv.-Sok. (Syn.: <i>R. kladnii</i> auct. non Schur)	EC (UA)	East-Carpathian endemic	Cvelev 2001; Diduch et al. 2004a; Čornej 2011; Turlaj 2011

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Ranunculus pseudomontanus</i> Schur ^{156, 157} (Syn.: <i>R. montanus</i> subsp. <i>pseudomontanus</i> (Schur) Ciocârlan)	WC (SK, PL), EC (UA, RO), SC (RO), AC	West-East-South-Apuseni-Carpathian (pan-Carpathian) endemic	Kožuharov & Petrova 1988; Kliment 1999; Oprea 2005
<i>Rosa coziae</i> Nyár. ^{158, 159} (Syn.: <i>R. villosa</i> subsp. <i>coziae</i> (Nyár.) Ciocârlan)	SC (RO)	South-Carpathian endemic	Oprea 2005; Ciocârlan 2009; Kerényi-Nagy 2011; Hurdu et al. 2012a
<i>Salix kitaibeliana</i> Willd. (Syn.: <i>S. retusa</i> subsp. <i>kitaibeliana</i> (Willd.) Jáv.)	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999
<i>Salvia transsylvanica</i> (Schur ex Griseb. et Schenk) Schur ¹⁶⁰	Tr, EC (RO)/SC (RO)	endemic to Transylvanian Basin, Eastern and Southern Carpathians	Dihoru & Pârvu 1987; Oprea 2005
<i>Saussurea porcii</i> Degen ¹⁶¹	EC (UA, RO)	East-Carpathian endemic	Kobiv et al. 2007b; Dihoru & Negrean 2009; Bahlej 2010; Kobiv 2010; Počynok & Prokopiv 2010; Derevenko 2011; Mátyás et al. 2014
<i>Saxifraga moschata</i> subsp. <i>dominii</i> Soó	WC (SK, PL)	West-Carpathian endemic	Kliment 1999
<i>Saxifraga moschata</i> subsp. <i>kotulae</i> S. Pawł.	WC (SK, PL)	West-Carpathian endemic (Tatry Mts, Nizke Tatry Mts)	Kliment 1999; Mirek & Piękoś-Mirkowa 2010
<i>Saxifraga mutata</i> subsp. <i>demissa</i> (Schott et Kotschy) D. A. Webb (Syn.: <i>S. demissa</i> Schott et Kotschy; <i>S. transsylvanica</i> Fuss)	SC (RO)	South-Carpathian endemic	Beldie 1977; Dihoru & Pârvu 1987; Webb 1993a; Oprea 2005; Ciocârlan 2009; Sârbu et al. 2013
<i>Saxifraga wahlenbergii</i> Ball	WC (SK, PL)	West-Carpathian endemic	Kliment 1999; Cieślak et al. 2013
<i>Scabiosa columbaria</i> subsp. <i>banatica</i> (Waldst. et Kit.) Diklić ¹⁶² (Syn.: <i>S. banatica</i> Waldst. et Kit.; <i>S. columbaria</i> subsp. <i>banatica</i> (Waldst. et Kit.) Soó, nom. illeg.)	SC (RO, SRB), AC (RO)/EC (RO)	East-South-Apuseni-Carpathian endemic	Prodan 1961; Diklić 1973; Mardari 2009; Hurdu et al. 2012b
<i>Scabiosa lucida</i> subsp. <i>barbata</i> Nyár. (Syn.: <i>S. pseudobanatica</i> subsp. <i>barbata</i> (Nyár.) Chrték; <i>S. opaca</i> Klokov)	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Kobiv et al. 2007a; Oprea 2005
<i>Scilla kladnii</i> Schur ¹⁶³ (Syn.: <i>S. bifolia</i> subsp. <i>subtriphylla</i> (Schur) Domin; <i>S. alpina</i> Schur)	WC (CZ, SK, HU, PL), EC (SK, PL, UA), SC (RO), AC (RO), Tr (RO)	subendemic to Western, Eastern, Southern & Apuseni Carpathians and Transylvanian Basin (pan-Carpathian subendemic)	Kereszty 1993; Kricsfalusy & Vajnagi 1994; Kliment 1999; Kochjarová et al. 2004, 2005; Trávníček et al. 2010
<i>Scorzoneroideis pseudotaraxaci</i> (Schur) Holub (Syn.: <i>Leontodon pseudotaraxaci</i> Schur; <i>L. montanus</i> subsp. <i>pseudotaraxaci</i> (Schur) Finch et P. D. Sell)	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999; Oprea 2005; Čornej 2011
<i>Sempervivum carpathicum</i> Wettst. ex Prodan ¹⁶⁴	WC (SK, PL), EC (PL, UA, RO), SC (RO)	West-East-South-Carpathian endemic	Letz 1998; Kliment 1999
<i>Sempervivum carpathicum</i> Wettst. ex Prodan subsp. <i>carpathicum</i> ¹⁶⁵ (Syn.: <i>S. montanum</i> subsp. <i>carpathicum</i> (Wettst. ex Prodan) A. Berger)	WC (SK, PL), EC (PL, UA, RO), SC (RO)	West-East-South-Carpathian endemic	Letz 1998, 2002; Letz & Marhold 1998; Kliment 1999
<i>Sempervivum carpathicum</i> subsp. <i>heterophyllum</i> (Hazsl.) Letz ¹⁶⁶ (Syn.: <i>S. montanum</i> subsp. <i>heterophyllum</i> (Hazsl.) Jáv. ex Soó)	WC (SK)	West-Carpathian endemic	Letz 1998, 2002; Letz & Marhold 1998; Kliment 1999
<i>Sempervivum matricum</i> Letz ¹⁶⁷	WC (SK, HU)	West-Carpathian subendemic	Blanár & Letz 2005; Letz 2009
<i>Senecio dacicus</i> Hodálová et Marhold (Syn.: <i>S. hercynicus</i> subsp. <i>dacicus</i> (Hodálová et Marhold) Greuter)	SC (RO), AC (RO)	South-Apuseni-Carpathian endemic	Hodálová & Marhold 1998; Hodálová 1999a, b; Negrean 2011
<i>Senecio ucranicus</i> Hodálová ¹⁶⁸ (Syn.: <i>S. hercynicus</i> subsp. <i>ucranicus</i> (Hodálová) Greuter)	EC (SK, PL, UA, RO), SC (RO)	East-South-Carpathian subendemic	Hodálová 1999b; Rola 2014
<i>Sesleria heuflerana</i> Schur	WC (SK, HU), EC (UA, RO), SC (RO), AC (RO), Tr (RO)	subendemic to Western, Eastern, Southern & Apuseni Carpathians and Transylvanian Basin (pan-Carpathian subendemic)	Kliment 1999

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Sesleria heuflerana</i> Schur subsp. <i>heuflerana</i> ¹⁶⁹	WC (SK, HU), EC (UA, RO), SC (RO), AC (RO), Tr (RO)	subendemic to Western, Eastern, Southern & Apuseni Carpathians and Transylvanian Basin (pan-Carpathian subendemic)	Čopyk 1976; Gejdeman 1986; Kliment 1999; Oprea 2005
<i>Sesleria heuflerana</i> subsp. <i>hungarica</i> (Ujhelyi) Deyl ¹⁷⁰ (Syn.: <i>S. hungarica</i> Ujhelyi)	WC (HU)	West-Carpathian endemic (Bükk Mts)	Lysák 1996; Kliment 1999; Borhidi et al. 2012
<i>Sesleria rigida</i> Heuff. ex Rchb. ¹⁷¹ (Syn.: <i>S. haynaldiana</i> Schur; <i>S. rigida</i> subsp. <i>haynaldiana</i> (Schur) Beldie)	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Morariu & Beldie 1976; Beldie 1979; Oprea 2005; Kuzmanović et al. 2013
<i>Sesleria tatrae</i> (Degen) Deyl ¹⁷² (Syn.: <i>S. sadlerana</i> subsp. <i>tatrae</i> (Degen) Deyl)	WC (SK, PL)	West-Carpathian subendemic	Fabiszewski 1970; Hendrych 1987; Kliment 1999; Budzáková et al. 2014
<i>Silene dinarica</i> Spreng.	SC (RO)	South-Carpathian endemic	Beldie 1977; Chater et al. 1993; Ciocârlan 2009
<i>Silene nivalis</i> (Kit.) Rohrb. (Syn.: <i>Lychnis nivalis</i> Kit.; <i>Polyschemone nivalis</i> (Kit.) Schott, Nyman et Kotschy)	EC (RO)	East-Carpathian endemic (Rodna Mts)	Coldea 1997; Oprea 2005; Ciocârlan 2009; Dihoru & Negrean 2009; Sârbu et al. 2013
<i>Silene nutans</i> subsp. <i>dubia</i> (Herbich) Zapał. (Syn.: <i>S. dubia</i> Herbich)	EC (SK, PL, UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Kliment 1999; Oprea 2005; Mered'a et al. 2012
<i>Silene zawadzkii</i> Herbich ^{173, 174} (Syn.: <i>Elisanthe zawadzkii</i> (Herbich) Klokov; <i>Melandrium zawadzkii</i> (Herbich) A. Braun; <i>Silenanthe zawadzkii</i> (Herbich) Griseb. et Schenk)	EC (UA, RO)	East-Carpathian endemic	Klokov 1952; Morariu & Beldie 1976; Beldie 1977; Chater et al. 1993; Fedorončuk & Diduch 2002d; Cvelev 2004a; Ciocârlan 2009; Kobiv 2010; Hurdu et al. 2012a; Sârbu et al. 2013
<i>Soldanella carpatica</i> Vierh.	WC (SK, PL)	West-Carpathian endemic	Kliment 1999; Zhang & Kadereit 2002; Kochjarová & Hroudá 2006
<i>Sorbus amici-petri</i> Mikoláš ¹⁷⁵	WC (SK)	West-Carpathian endemic (Čierna hora Mts)	Mikoláš 2003
<i>Sorbus atrimontis</i> Bernátová et Májovský	WC (SK)	West-Carpathian endemic (Veľká Fatra Mts)	Bernátová & Májovský 2003; Uhlířová & Bernátová 2004; Kliment et al. 2008
<i>Sorbus borbasii</i> Jáv.	SC (RO)	South-Carpathian endemic	Kárpáti 1960; Oprea 2005; Ciocârlan 2009; Dihoru & Negrean 2009; Hurdu et al. 2012a
<i>Sorbus caeruleomontana</i> Bernátová et Májovský	WC (SK)	West-Carpathian endemic (Nízke Tatry Mts: Mt. Siná)	Bernátová & Májovský 2003; Šípošová et al. 2004b
<i>Sorbus dacica</i> Borbás	AC (RO)	Apuseni-Carpathian endemic (Trascău Mts)	Kárpáti 1960; Dihoru & Negrean 2009; Hurdu et al. 2012a
<i>Sorbus diversicolor</i> Bernátová et Májovský	WC (SK)	West-Carpathian endemic (Veľká Fatra Mts)	Bernátová & Májovský 2003; Šípošová et al. 2004a; Uhlířová & Bernátová 2004
<i>Sorbus dolomiticola</i> Mikoláš ¹⁷⁶	WC (SK)	West-Carpathian endemic (Čierna hora Mts)	Mikoláš 1996
<i>Sorbus haljamovae</i> Bernátová et Májovský ¹⁷⁷	WC (SK)	West-Carpathian endemic	Bernátová & Májovský 2003; Uhlířová & Bernátová 2004
<i>Sorbus hazslinszkyana</i> (Soó) Májovský (Syn.: <i>S. austriaca</i> subsp. <i>hazslinszkyana</i> (Soó) Kárpáti)	WC (SK, HU)	West-Carpathian endemic	Kárpáti 1960; Kliment 1999
<i>Sorbus margittaiana</i> (Jáv.) Kárpáti (Syn.: <i>S. hostii</i> subsp. <i>margittaiana</i> Jáv.)	WC (SK)	West-Carpathian endemic (Krivánska Fatra Mts)	Bernátová et al. 1998; Májovský et al. 1998
<i>Sorbus montisalpae</i> Bernátová et Májovský	WC (SK)	West-Carpathian endemic (Veľká Fatra Mts)	Bernátová & Májovský 2003; Šípošová et al. 2004a; Uhlířová & Bernátová 2004

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Sorbus pekarovae</i> Májovský et Bernátová	WC (SK)	West-Carpathian endemic (Veľká Fatra Mts: Mt. Pekárová)	Májovský & Bernátová 1996; Bernátová & Májovský 1999; Šipošová et al. 2004a
<i>Sorbus salatinii</i> Bernátová et Májovský	WC (SK)	West-Carpathian endemic (Nízke Tatry Mts: Mt. Salatin)	Bernátová & Májovský 2003; Šipošová et al. 2004b
<i>Sorbus scepusiensis</i> Kovanda	WC (SK)	West-Carpathian endemic (Voloenské vrchy Mts)	Kovanda 1985, 1986; Kliment 1999
<i>Sorbus umbellata</i> subsp. <i>banatica</i> (Jáv.) Kárpáti	SC (RO)	South-Carpathian endemic	Kárpáti 1960; Ciocârlan 2009
<i>Sorbus zuzanae</i> Májovský et Bernátová ¹⁷⁸	WC (SK)	West-Carpathian endemic	Bernátová & Májovský 2003; Šipošová et al. 2004a; Uhlířová & Bernátová 2004
<i>Stipa crassiculmis</i> subsp. <i>heterotricha</i> Dihoru et Roman ¹⁷⁹	SC (RO)	South-Carpathian endemic (Cozia Mts)	Dihoru & Pârnu 1987; Negrean & Oltean 1989; Dihoru & Negrean 2009; Negrean 2011; Vázquez & Gutiérrez 2011
<i>Stipa danubialis</i> Dihoru et Roman	SC (RO)	South-Carpathian endemic (Almajului Mts: Porțile de Fier Gorge)	Beldie 1979; Dihoru & Pârnu 1987; Oprea 2005; Ciocârlan 2009; Dihoru & Negrean 2009
<i>Swertia punctata</i> Baumg. ¹⁸⁰ (Syn.: <i>S. perennis</i> subsp. <i>punctata</i> (Baumg.) Ciocârlan)	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian subendemic	Beldie 1979; Tan & Vladimirov 2001; Oprea 2005; Kricsfalussy & Budnikov 2007; Kobiv 2010, 2012a
<i>Symphytum cordatum</i> Waldst. et Kit. ex Willd. ¹⁸¹ (Syn.: <i>S. cordatum</i> Waldst. et Kit., nom. illeg.; <i>S. cordifolium</i> Baumg.)	WC (SK, PL), EC (SK, PL, UA, RO), SC (RO), AC (RO), Tr (RO)	subendemic to Western, Eastern, Southern & Apuseni Carpathians and Transylvanian Basin (pan-Carpathian subendemic)	Čopyk 1976; Roman et al. 1996; Kliment 1999; Oprea 2005; Čornej 2011
<i>Syringa josikaea</i> J. Jacq. ex Rchb. f.	EC (UA), AC (RO)	East-Apuseni-Carpathian endemic	Dihoru & Negrean 2009; Bilz 2013; Lendvay et al. 2013
<i>Taraxacum carpaticum</i> Štěpánek et Kirschner ¹⁸²	SC (RO)	South-Carpathian endemic	Štěpánek et al. 2011
<i>Taraxacum erythrocarpum</i> Kirschner et Štěpánek	WC (SK)	West-Carpathian endemic	Kirschner & Štěpánek 1985; Kliment 1999
<i>Taraxacum nigricans</i> (Kit.) Rchb. ¹⁸³	WC (SK)	West-Carpathian endemic (Nízke Tatry Mts)	Štěpánek et al. 2011
<i>Taraxacum pawlowskii</i> Soest ¹⁸⁴	WC (PL, ?SK)	West-Carpathian endemic (Tatry Wysokie Mts)	Tacik 1980; Mirek & Piękoś-Mirkowa 2009, 2010
<i>Taraxacum pieninicum</i> Pawł. ¹⁸⁵ (Syn.: <i>T. hoppeanum</i> subsp. <i>pieninicum</i> (Pawł.) Pawł.)	WC (PL)	West-Carpathian endemic (Pieniny Mts: Mt. Trzy Korony)	Zarzycki 1986; Mirek et al. 1995; Wróbel & Zarzycki 2008
<i>Tephrosieris longifolia</i> subsp. <i>moravica</i> Holub	WC (CZ, SK)	West-Carpathian endemic	Kochjarová 1998; Kochjarová & Hrouda 2004; Janišová et al. 2012; Hegedúšová et al. 2013; Olšovská et al. 2015
<i>Thalictrum minus</i> subsp. <i>carpaticum</i> (B. Kotula) Osvač.	WC (SK, PL)	West-Carpathian endemic (Tatry Mts)	Kliment 1999; Mirek & Piękoś-Mirkowa 2010
<i>Thesium kernerianum</i> Simonk.	EC (RO), SC (RO)	East-South-Carpathian endemic	Negrean & Oltean 1989; Oprea 2005; Dihoru & Negrean 2009
<i>Thymus alternans</i> Klokov ¹⁸⁶	EC (SK, UA, RO)/SC (RO), AC (RO)	East-South-Apuseni-Carpathian subendemic	Klokov 1960; Mártonfi 1996; Kliment 1999; Oprea 2005; Mártonfi 2014 in litt.
<i>Thymus bihoriensis</i> Jalas (Syn.: <i>T. marginatus</i> A. Kern. non Sm. ex Dickson, nom. illeg.)	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Beldie 1979; Dihoru & Pârnu 1987; Oprea 2005
<i>Thymus comosus</i> Heuff. ex Griseb. et Schenk (Syn.: <i>T. comosus</i> Schur, nom. illeg.)	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Morariu & Beldie 1976; Dihoru & Pârnu 1987; Oprea 2005
<i>Thymus dacicus</i> Borbás ¹⁸⁷ (Syn.: <i>T. porcii</i> Borbás)	EC (RO), SC (RO, SRB), AC (RO)	East-South-Apuseni-Carpathian subendemic	Gușuleac 1961; Diklić & Vasić 2000
<i>Thymus pulcherrimus</i> Schur	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian subendemic	Kliment 1999; Oprea 2005

Taxon	Occurrence in the Carpathians	Category of endemism	References
<i>Thymus pulcherrimus</i> Schur subsp. <i>pulcherrimus</i> (Syn.: <i>T. circumcinctus</i> Klokov)	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Mártonfi 1997; Kliment 1999; Čornej 2011
<i>Thymus pulcherrimus</i> subsp. <i>sudeticus</i> (Lyka) P. A. Schmidt ^{188, 189} (Syn.: <i>T. carpathicus</i> Čelak.)	WC (SK, PL)	West-Carpathian subendemic	Kliment 1999
<i>Trifolium orbelicum</i> subsp. <i>monticolum</i> (Domin) Májovský	WC (SK, PL)	West-Carpathian endemic	Kliment 1999; Mirek & Piękoś-Mirkowa 2010
<i>Trifolium pratense</i> subsp. <i>kotulae</i> (Pawl.) Soják ¹⁹⁰ (Syn.: <i>T. pratense</i> var. <i>kotulae</i> Pawl.; <i>T. frigidum</i> Schur)	WC (SK, PL), EC (UA, RO), SC (RO)	West-East-South-Carpathian endemic	Kliment 1999; Oprea 2005; Kricsfalusy & Budnikov 2007
<i>Trifolium sarosienae</i> Hazsl. ¹⁹¹ (Syn.: <i>T. medium</i> subsp. <i>sarosienae</i> (Hazsl.) Simonk.; <i>T. sarosienae</i> subsp. <i>banaticum</i> (Heuff.) Holub)	WC (SK, HU), EC (?UA, RO), SC (RO), AC (RO), Tr (RO)	subendemic to Western, Eastern, Southern & Apuseni Carpathians and Transylvanian Basin (pan-Carpathian subendemic)	Hendrych 1993, 1995; Kliment 1999
<i>Trisetum flavescens</i> subsp. <i>tatricum</i> Chrtek ¹⁹²	WC (SK), EC (SK, PL, †UA)	West-East-Carpathian endemic	Kliment 1999; Čornej 2011
<i>Trisetum fuscum</i> (Kit. ex Schult.) Schult. (Syn.: <i>T. ciliare</i> (Kit.) Domin)	WC (SK, PL), EC (UA, RO), SC (RO), AC (RO)	West-East-South-Apuseni-Carpathian (pan-Carpathian) endemic	Kliment 1999; Oprea 2005
<i>Trisetum macrotrichum</i> Hack.	EC (RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Buia & Morariu 1972; Morariu & Beldie 1976; Oprea 2005
<i>Tulipa hungarica</i> Borbás ^{193, 194} (Syn.: <i>T. hungarica</i> subsp. <i>undulatifolia</i> (Roman) Roman et Beldie)	SC (RO, †SRB)	South-Carpathian endemic	Ciocârlan 2009; Dihoru & Negrean 2009; Negrean 2011; Čalić et al. 2012; Stevanović 2013
<i>Valeriana tripteris</i> subsp. <i>heterophylla</i> (Baumg.) Rostański ^{195, 196, 197} (Syn.: <i>V. tripteris</i> var. <i>heterophylla</i> Baumg.; <i>V. transilvanica</i> Schur; <i>V. sisymbriifolia</i> Schur non Vahl, nom. illeg.)	EC (SK, PL, UA, RO), SC (RO), AC (RO)/WC (PL)	East-South-Apuseni-Carpathian endemic	Schur 1866; Simonkai 1887; Katina 1961; Morariu 1961; Rostański 1967, 1970; Prokudin 1987; Dmytrach 2010; Tasenkevych 2014
<i>Viola declinata</i> Waldst. et Kit. ¹⁹⁸	EC (UA, RO), SC (RO), AC (RO)	East-South-Apuseni-Carpathian endemic	Oprea 2005; Velez & Apostolova 2009
<i>Viola jooi</i> Janka ¹⁹⁹	EC (UA, RO), SC (RO), AC (RO), Tr (RO)	subendemic to Eastern, Southern & Apuseni Carpathians and Transylvanian Basin	Grințescu et al. 1955; Oprea 2005; Čornej 2011; Cristea 2014

Notes

¹ The occurrence of *Aconitum degenii* subsp. *degenii* was reported from the Eastern, Southern and Apuseni Carpathians (e.g. Mucher 1993, Starmühler 1998a, 2000). In Slovak herbarium collections, Novikov (2013a, 2014 in litt.) found also several specimens from the Slovak part of the Western Carpathians (cf. Eliáš jr. et al. 2015).

² Outside the Carpathian arch *Aconitum firmum* subsp. *fissurae* is known from Velebit Mts in Croatia (Starmühler & Mitka 2001). In the record from the central Dneper Valley (Starmühler 1997) the taxon might have been changed with the local endemic *Aconitum odontandrum* Wissjul. (Novikov 2014 in litt.).

³ In the included literature *Aconitum firmum* subsp. *fissurae* is reported as *Aconitum napellus* subsp. *fissurae* (Nyár.) Seitz; *A. callibotryon* subsp. *hunyadense* (Degen) Soó; *A. firmum* subsp. *hunyadense* (Degen) Morariu et Beldie; *A. tatarae* subsp. *hunyadense* (Degen) Soó; *A. tauricum* subsp. *hunyadense* (Degen) Ciocârlan or *A. firmum* subsp. *romanicum* (Wol.) Beldie.

⁴ Eligibility of a separate subspecies *Aconitum firmum* subsp. *skerisorae* was justified by Starmühler (2000); it was accepted also by Mitka (2003), Tasenkevich (2011) and Novikov (2014 in litt.).

⁵ *Aconitum lasianthum* has centre of its distribution in the Romanian Southern Carpathians, it occurs sparsely also in the Eastern (Penteleu Mts) and Apuseni Carpathians (Trascău Mts) (Grințescu 1953, Oprea 2005).

- ⁶ The record of *Aconitum lasiocarpum* subsp. *lasiocarpum* from the Beskid Niski Mts (Mitka & Starmühler 2000) concerns subsp. *kotulae* (Mitka 2003).
- ⁷ Outside the Carpathians *Aconitum lasiocarpum* subsp. *kotulae* was reported from Podillja and Polissja in Ukraine (Mitka 2001, 2003, Novikoff & Mitka 2011a, b; cf. Diduch 2009).
- ⁸ In the Southern Carpathians *Aconitum lasiocarpum* subsp. *kotulae* is documented by a single older record (Walz 1884 CL) from Mt. Cristianul Mare in the Postavărul Massif (Mitka & Starmühler 2000, Mitka 2003).
- ⁹ In Poland and Ukraine *Aconitum moldavicum* subsp. *moldavicum* occurs also outside the Carpathians at lower altitudes in Kotlina Sandomierska, Wyżyna Małopolska, Wyżyna Lubelska; Opillja, Podillja (Mitka 2003, 2008). Naturalized plants from a cultivation were found near Graz in Austria (Starmühler 2004).
- ¹⁰ In Poland and Ukraine *Aconitum moldavicum* subsp. *hosteanum* has been recorded also outside the Carpathians: Kotlina Sandomierska, Pogórze Przemyskie, Góry Świętokrzyskie, Wyżyna Małopolska; Podillja, Roztoččja, Mal'e Polissje, Voronjaky, Holohory (Mitka 2008).
- ¹¹ Outside the Carpathians *Aconitum toxicum* subsp. *toxicum* is known from two localities in the surrounding of Sarajevo (Bosnia) and from Mt. Ranisava (Durmitor Mts) and Mt. Balj in Montenegro (Mucher 1993, Ilnicki & Mitka 2011).
- ¹² Based on the Soják's herbarium material from Făgăraș Mts, Plocek (1983) reported *Alchemilla boleslai* also from the Southern Carpathians; later he reported its distribution area as restricted to the Western Carpathians (Plocek 1992). Negrean (2011) wrongly localized the record of Plocek to Parâng Mts. As Boruz (2014 in litt.) failed in finding the species in the Romanian herbaria and in the field, she considered its occurrence in Romania as doubtful (cf. Kurtto et al. 2007). The reference material was not found in the Soják's collections (PR) (Šída 2014 in litt.), nor in the collections of PRC (Hadinec 2014 in litt.).
- ¹³ Syčák (1992) described *Alchemilla bucovinensis* from the Ukrainian Carpathians and identified it also with populations from the Romanian Eastern Carpathians (Maramureș Mts), provisionally reported as *A. romanica* Plocek ined. (cf. Kurtto et al. 2007).
- ¹⁴ *Alchemilla czywczyńska* was described from Čyvcyny Mts (Pawłowski 1952). Outside the Eastern Carpathians it was found in Pamporovo (Rodopi Mts, Bulgaria) (Fröhner 1986, Kurtto et al. 2007, 2009).
- ¹⁵ In the original diagnosis of *Alchemilla eugenii*, Pawłowski (1952) reported its occurrence in Mt. Kresanica (2 110 m a.s.l.) at the Slovak-Polish border. Plocek (1992) considered this record, documented by a herbarium specimen, as dubious.
- ¹⁶ Apart from frequent occurrence in the Moravsko-sliezske Beskydy Mts and Javorníky Mts *Alchemilla gruneica* is known also from the Hrubý Jeseník Mts in the Sudeten Mts (Plocek 1992).
- ¹⁷ Syčák (2002) reported *Alchemilla ladislai* also from the Ukrainian Carpathians; however, these data are erroneous (Syčák sec. Kurtto et al. 2007, Syčák 2011).
- ¹⁸ According to Plocek (Plocek in Kliment 1999), the populations from the Veľká Fatra Mts and Javorníky Mts reported as *Alchemilla laxa* (Plocek 1990, 1992) belong to a different taxon.
- ¹⁹ According to Plocek (1992), similar populations occur also in the Southern Carpathians (Bucegi Mts). Kurtto et al. (2007) report *Alchemilla marginata* only from Slovakia. Boruz (2014 in litt.) has not found this species in the Southern Carpathians; according to her knowledge it is absent from Romania.
- ²⁰ Plocek (1992) considered *Alchemilla sokolowskii* to be a synonym of *A. pseudincisa*. In our study we follow the concept of Kurtto et al. (2007), who consider *A. sokolowskii* to be a separate species.
- ²¹ With respect to the relative alpine flora strongly affected by the Pleistocene glaciation we recognize endemics to Tatry Mts s. str. occurring in the Západné Tatry Mts, Vysoké Tatry Mts and Belianske Tatry Mts, and endemics to Tatry Mts s. lat. occurring also at higher altitudes of the Nízke Tatry Mts (for details see Kliment 2003); to distinguish between them we report the precise information on their occurrence.

- ²² *Alchemilla suavis* was described from the Moravian part of the Bílé Karpaty Mts and found also on Mt. Krompašský vrch in the Slovenské rudohorie Mts (Plocek 1973, 1992); such distribution was accepted also by Kurtto et al. (2007). According to Plocek (1992), the species area involves also the Ukrainian and Romanian Eastern Carpathians. However, Syčák (2002, 2011) has not reported this species from the Ukrainian Carpathians. Neither it was found in the Romanian herbarium collections nor in the Romanian Carpathians (Boruz 2014 in litt.). Based on these facts we reduced its distribution area to the Western Carpathians.
- ²³ Plocek (1992) reported *Alchemilla versipiloides* as synonym of *A. versipila* Buser; however, Kurtto et al. (2007) accepted it as a separate species (cf. Piękoś-Mirkowa & Mirek 2009).
- ²⁴ There is only a single, recently not confirmed, evidence of *Alchemilla zapalowiczii* occurrence from the Western Carpathians, specifically from the Belianske Tatry Mts (Dostál 1949 PRC). Therefore, Kurtto et al. (2007) consider this species as probably extinct in Slovakia. Based on the second volume of Flora Europaea (Walters & Pawłowski 1968; cf. Negrean 2011) the occurrence of *A. zapalowiczii* is reported also from the Romanian Carpathians although it was confirmed neither in herbaria nor during the field research (Boruz 2014 in litt.). Kurtto et al. (2009) reported sporadic occurrence of this species from the Jakupica Mts in Macedonia.
- ²⁵ Plocek (1992) reported *Alchemilla zmudae* as synonym of *A. stanislaae*; however, Kurtto et al. (2007) accepted it as a separate species (cf. Mirek & Piękoś-Mirkowa 2009, 2010).
- ²⁶ According to Brullo et al. (1996) *Allium fussii* differs from *A. fuscum* in several traits of leaves and inflorescences, therefore it was regarded as a separate species within the *Allium paniculatum* group (cf. Mráz 2005a, Šafářová et al. 2011).
- ²⁷ Based on common morphological seed traits Sennikov (1999a) ordered *Andryala laevimentosa* together with another narrow endemic (SE Spain, Morocco) relic alpine species *A. agardhii* Haens. ex DC. to oligotypic genus *Pietrosia* Nyár. ex Sennikov (cf. Negrean 2004). The results of the most recent molecular-phylogenetic study (Ferreira et al. 2015) support the opinion of Greuter (2003) and justify its backward classification within the monophyletic genus *Andryala* L. Up-to-date note on taxonomy, nomenclature, biology, phylogeny and conservation status of *Andryala laevitomentosa*, particularly important microchoric species endemic to the Romanian Carpathians, were published by Negrea & Pricop (2009a, b ut *Pietrosia laevitomentosa*). Records from the Ukrainian Carpathians demand of a confirmation (cf. Kricsfalussy & Budnikov 2007).
- ²⁸ In the included literature it is reported also using invalid names *Andryala levitomentosa* (Nyár.) P. D. Sell., *Hieracium levimentosum* (Nyár.) Soó and *Pietrosia levitomentosa* Nyár.
- ²⁹ The occurrence of *Antennaria carpatica* subsp. *carpatica* in Romania is reported only in older recently not confirmed records from the Gutâi, Ceahlău, Bucegi and Făgăraș Mts (Oprea 2005; cf. Ciocârlan 2009, Sârbu et al. 2013), therefore the subspecies has been considered as extinct in Romania already by Oltean et al. (1994). Chrtek & Pouzar (1985) report its occurrence only from the Western (Vysoké Tatry Mts, Nízke Tatry Mts) and Eastern Carpathians (Mt. Blyznychja, Mt. Drahobrat).
- ³⁰ In the Eastern Carpathians *Anthemis cretica* subsp. *pyrethroides* grows only in the Ceahlău Mts (Beldie 1979, Oprea 1985, Sârbu et al. 2013).
- ³¹ In some Romanian literature (Dihoru & Pârvu 1987, Negrean & Oltean 1989, Oltean et al. 1994, Popescu & Sanda 1998, Sârbu et al. 2013) it is invalidly reported as *Anthemis carpatica* subsp. *pyrethroides* (Schur) Beldie (cf. Fernandes 1976).
- ³² The occurrence of *Aquilegia transsilvanica* is reported also from the Ukrainian Carpathians (e.g. Čopyk 1976, Stojko & Tasenkewitsch 1991, Stoyko & Tasenkevich 1993, Vasil'eva 2001, 2012, Kricsfalussy & Budnikov 2002, 2007, Tasenkevych 2003b, Ziman et al. 2009). According to Kobiv (2012b), it certainly grows in the Southern Carpathians (Paríng, Făgăraș, Iezer and Bucegi Mts) and outside them only in the southeastern part of the Eastern Carpathians (Buzău Mts), while all records from the Ukrainian Carpathians are erroneous.
- ³³ Sporadic records of *Arabidopsis neglecta* in the Austrian Alps (Janichen 1957, Čopyk 1976, Piękoś-Mirkowa et al. 1996, etc.) refer alpine forms of *Arabidopsis arenosa* (cf. Kliment 1999).
- ³⁴ *Armeria pocutica* is a very rare stenochoric endemic described in the Ukrainian Carpathians and known only from the Čornyj Čeremoš Valley in the Čyvčyny Mts (Pawłowski 1962) in Ukraine and a single locality (Borșa, under Mt. Luhi in the Cizla Valley) in the Romanian Maramureș Mts (Beldie & Váczy 1976, Morariu & Beldie 1976, Beldie 1979, Ciocârlan 1988, 2009, Popescu & Sanda 1998, Negrean 2011). Its occurrence in its *locus classicus* has not been confirmed since the last 30 years (Kahalo & Syčák 2009).
- ³⁵ Apart its frequent occurrence in Hășmaș Mts, where there is a centre of its distribution, *Asperula carpatica* occurs also in the Rarău Mts (here only within its *locus classicus* in Pietrele Doamnei); in the Southern Carpathians its occurrence was reported from a single locality in the Piatra Craiului Mts (Negrean 2011).

³⁶ *Astragalus australis* subsp. *krajinae* is an alpine taxon known from a sub-alpine belt of the Ukrainian mountains Svydovec Mts and Marmaroš Mts (Ziman 2009). It was described by Domin (1931a) from Svydovec Mts as a species distinct from *A. australis* (L.) Lam., but later it was re-classified to subspecies (Domin 1935). Since its description it is unequivocally accepted by the Ukrainian authors (in both taxonomic and syntaxonomic literature); and it is also accepted in the Carpathian Red List (Witkowski et al. 2003). In some comprehensive taxonomic studies (Chater 1968; Podlech 2008, 2011) it is evaluated as synonym of *A. australis*.

³⁷ Rare and endangered stenoendemic of the Romanian flora, *Astragalus peterfii*, is recently known from two adjacent localities situated only 5 km from each other (Suatu, Căianu) located east of Cluj (Șuteu et al. 2003, Bartha 2012).

³⁸ *Astragalus roemeri* has centre of its distribution in the Romanian Carpathians (Hășmaș Mts). For a longer time not confirmed sporadic occurrence from the Apuseni Carpathians (cf. Váczy & Beldie 1976) was confirmed by Bartha & Bartók (2013), who found two flowering and one sterile individuals on Mt. Scărița. According to these authors, the population is at the border of extinction; it represents a rare exclave occurrence outside the Eastern Carpathians.

³⁹ The occurrence of *Athamanta turbith* subsp. *hungarica* in Serbia is restricted to the Serbian part of the Southern Carpathians (Gornjačka klisura; cf. Stevanović et al. 1991, Hurdu et al. 2012b).

⁴⁰ The subspecies *Aubrieta columnae* subsp. *platycarpa*, known only from the Piatra Craiului Mts (Marele Grohotiș) in the Southern Carpathians, is reported using distinct names in the Romanian literature: *Aubrieta deltoidea* (L.) DC., *A. intermedia* Heldr. et Orph. ex Boiss., *A. intermedia* subsp. *falcata* Ciocârlan and also using invalidly published name *A. deltoidea* subsp. *falcata* (Ciocârlan) Adr. Oprea. To avoid confusions, Ciocârlan (2006) published a new name *A. columnae* subsp. *platycarpa*, but probably due to its publication in less accessible conference proceedings this name is not reported in any supranational database. In the Romanian Red Book (Dihoru & Negrean 2009) the subspecies is mentioned in the note to another subspecies, *A. columnae* subsp. *croatica* (Schott, Nyman et Kotschy) Mattf. (Parâng Mts), which Anastasiu & Negrean (2005) classified as alien taxon of the Romanian flora.

⁴¹ In the Southern Carpathians *Barbarea lepuznica* is recently occurring in the Retezat Mts (including Mt. Piule in the Retezat Mic Mts) and in the Godeanu Mts (Borăscu massif). Outside the Southern Carpathians it grows in Vršacki breg (Vršac Mts) within the Serbian Vojvodina, in close vicinity to the western margin of the Southern Carpathians (Dihoru & Negrean 2009, Strajeru & Stevanović 2013).

⁴² Soó (1973, 1980a) reported *Bromus monocladius* (ut *B. pannonicus* var. *monocladius* (Domin) Soó) also from Hungary (Budai Mts). According to recent information (Somlyay 2014 in litt.), this species does not occur in Hungary, it is also not included in the Hungarian vegetation database (Dúbravková 2014 in litt.).

⁴³ In the past, the name *Campanula kladniana* was used also for populations of other alpine species of *C. rotundifolia* agg., so that the information on its distribution area was distorted.

⁴⁴ Distinct conceptions of the species *Campanula tatrae* (for details see Kliment 1999, Goliašová et al. 2008) is reflected in distinct size of its distribution area and distinct endemic status reported in the literature – from the West-Carpathian endemic to the pan-Carpathian endemic.

⁴⁵ In Moravia, Poland and Ukraine *Cardamine glanduligera* sporadically occurs also in the adjacent geographic units (Czech Republic: Opavská pahorkatina Mts, Jesenícké podhůří Mts; Poland: Kotlina Śląska Basin, Wyżyna Krakowsko-Wieluńska Mts, Góry Świętokrzyskie Mts, Puszcza Sandomierska, Kotlina Sandomierska Basin, Wyżyna Lubelska Mts; Ukraine: Zakarpattia, Roztoččja-Opillja). Isolated localities are in the primeval forest Puszcza Knyszyńska in Poland and in the vicinity of Dnester estuary in Ukraine; it occurs scarcely also in shady forest of central Moldavia (Kliment 1999, Diduch 2007).

⁴⁶ In the literature *Carduus kernerii* subsp. *kernerii* is evaluated as a Carpathian-Balkan taxon. According to Franco (1976), data from the north Balkan mountains are related to subsp. *austro-orientalis* Franco (i.e. subsp. *scardicus* (Griseb.) Kazmi).

⁴⁷ In the Romanian literature *Carduus kernerii* subsp. *lobulatiformis* is also invalidly reported as *C. viridis* subsp. *lobulatiformis* (Csűrös et Nyár.) Beldie.

⁴⁸ *Centaurea reichenbachii* belongs to the *Centaurea stoebe* group. It was accepted by Dostál (1976), Ochsmann (2000) and Greuter (in Euro+Med). *Centaurea calvescens* Pančić was also included as synonym by Ochsmann (l. c.). According to Dostál (l. c.), Beldie (1979), Popescu & Sanda (1998), Dihoru & Negrean (2009), Ciocârlan (2009), Sârbu et al. (2013), Hurdu (2014 in litt.) and Vonica (2014 in litt.), these two species are close but different. Occurrence of *C. reichenbachii* in Romania is restricted to the Trascău Mts and Scărița-Belioara

Massif in the Apuseni Carpathians (Hurdu et al. 2012a); the data from the Iron Gates Gorge refer probably *C. calvescens* (cf. Beldie 1979, Dihoru & Negrean 2009). According to Mráz et al. (2011a), this species can hardly be morphologically distinguished from *C. stoebe*. Mráz et al. (2012) include it to *Centaurea stoebe*.

⁴⁹ *Centaurea rodnensis* Simonk. was invalidly reported from several Ukrainian and Romanian mountains of the Eastern Carpathians as *Centaurea carpatica* (Porcius) Porcius and it was considered as the local endemic. The revision of herbarium material indicated that *C. rodnensis* occurs rarely only in the Rodna Mts (5 localities) and is a narrow endemic of this mountain range. Older data from other mountains are erroneous and refer mainly *Centaurea phrygia* s. str. (Koutecký 2013).

⁵⁰ Following Beldie & Alexandrescu (1976), Morariu & Beldie (1976) and Beldie (1979), we preliminarily included *Centaurea rarauensis* to synonyms of *Centaurea rodnensis*. Koutecký (2014 in litt.) considers *C. rarauensis* to be a problematic taxon. The type does not exist and no plants matching the description were found (neither in herbaria nor in the field).

⁵¹ In the included literature *Centaurea rodnensis* was referred also as *Centaurea carpatica* (Porcius) J. Wagner, *C. carpatica* (Porcius) J. Wagner subsp. *carpatica*, *C. carpatica* (Porcius) Porcius subsp. *carpatica*, *C. phrygia* subsp. *carpatica* (Porcius) Dostál, *C. carpatica* subsp. *rarauensis* (Prodan) Ciocârlan and *C. phrygia* subsp. *rarauensis* (Prodan) Dostál.

⁵² *Centaurea triniifolia* belongs to the *Centaurea stoebe* group, which was accepted also by Dostál (1976), Ochsmann (2000) and Greuter (in Euro+Med). It grows in the southwestern part of the Southern Carpathians in broader vicinity of the Iron Gates Gorge; older records from the Trascău Mts need revision (Hurdu 2014 in litt.). According to Mráz et al. (2011a), it can hardly be morphologically distinguished from *C. stoebe*. Mráz et al. (2012) include it to *Centaurea stoebe*.

⁵³ *Cephalaria radiata* occurs predominantly in the Transylvanian Basin and adjacent Carpathian promontories. Scarcely it occurs also on the adverse side of the Southern Carpathians, in the vicinity of the city Ploiești (Prodan 1961).

⁵⁴ *Cephalaria uralensis* subsp. *multifida* is known only from several localities in the vicinity of the Iron Gates Gorge in Romania (Beldie et Váczy 1976, Dihoru & Pârvu 1987, Ciocârlan 2011, Sârbu et al. 2013 and others).

⁵⁵ Based on an older record (Hayek 1924) *Cerastium arvense* subsp. *lerchenfeldianum* was reported from Serbia or the former Yugoslavia (Morariu & Beldie 1976, Jalas & Suominen 1983, Jalas et al. 1993, Oprea 2005, Ciocârlan 2009); consequently this subspecies was evaluated as a Carpathian-Balkan taxon (e.g. Boșcaiu & Ehrendorfer 1996) or subendemic to the Eastern and Southern Carpathians (Heltmann 1985). According to Hurdu et al. (2012b), the record from Serbia is erroneous; they consider *Cerastium *lerchenfeldianum* to be endemic to the South-Eastern Carpathians.

⁵⁶ Letz (in Letz & Michalková 2012) found, that plants from the type locality of *Cerastium lerchenfeldianum* Schur are octoploid, similarly to populations of *Cerastium arvense* from almost the whole Carpathian region except the Tatry Mts (the distribution area of *Cerastium tatrae*). Regarding the leaf indument, the plants fall within the variability of *C. arvense*. For that reason, the authors include *Cerastium lerchenfeldianum* to synonyms of *Cerastium arvense* L. So far these preliminary results were not supported by a published comparative study. Until this problem is definitively resolved, we respect a separate position *Cerastium *lerchenfeldianum* at the subspecies level as well as its endemic status.

⁵⁷ Outside the Romanian and Serbian part of the Southern Carpathians *Clinopodium pulegium* scarcely grows in the Tara planina Plateau (Zaovine Village) in Serbia (Diklić & Nikolić 1986a, Hurdu et al. 2012b); a very infrequent occurrence was reported from Bosnia and Herzegovina nearby the cities Zvornik and Višegrad (Đug et al. 2013).

⁵⁸ Outside the Carpathians *Crocus banaticus* sporadically occurs in the vicinity of the city Šabac in Serbia (Hurdu et al. 2012b). Myhal' (2009) reports that the single population site in Serbia (without specifying the locality) was destroyed due to local economic activities.

⁵⁹ A single occurrence of *Cyanus maramarosiensis* in Slovakia („Mt. Stinka; determined by prof. Dostál“) was reported without further details by Májovský (1970 ut *Centaurea montana* subsp. *marmarossica*) within a note to the record of *Ranunculus carpaticus*. Hadač, Terray et al. (1991) reported from Stinská Mt. only the occurrence of *Centaurea mollis*. Also Greuter (in Euro+Med) and Olšovská (2014 in litt.) regard the occurrence of *Cyanus maramarosiensis* in Slovakia as questionable.

⁶⁰ Outside the Carpathians *Cyanus mollis* was found on Mt. Kobilica (Šar planina Mts) at the border of Macedonia and Kosovo and in two Croatian localities (Plitvička jezera and Mrzin) (Kliment 1999).

⁶¹ *Cyanus pinnatifidus* subsp. *sooanus* was described by Borhidi (1957 ut *Centaurea achtarovii* subsp. *sooana*) from the Ceahlău Mts (Eastern Carpathians); outside this mountain range it has not been found so far.

⁶² In the last decades, the narrow endemic subspecies *Cyclamen purpurascens* subsp. *immaculatum* was reported usually at the species level (as *Cyclamen fatrense*) in the literature focussing on the Western Carpathians. The current molecular-taxonomic study by Kučera et al. (2013) confirmed its separate position, however only at the level of subspecies.

⁶³ Soó (1980b) reported occurrence of *Dactylorhiza cordigera* subsp. *siculorum* also from the western Ukraine (i.e. from the Ukrainian Carpathians). However, the Ukrainian authors (Čopyk 1976, Malynovs'kyj et al. 2002, Kricsfalussy & Budnikov 2007, Kobiv 2010, etc.) report from the Ukrainian Carpathians only the Balkan-Carpathian species *D. cordigera*.

⁶⁴ *Dactylorhiza maculata* subsp. *schurii* was reported from the Eastern, Southern and Apuseni Carpathians (Soó 1967, Paucă & Beldie 1972); according to Hurdu (2014 in litt.) it is known also from several localities in the Transylvanian Basin.

⁶⁵ In the majority of literature, *Delphinium elatum* subsp. *nacladense* is evaluated as endemic of the Eastern Carpathians; however, Starmühler (1996b) reported also herbarium materials from the Piatra Craiului Mts in the Southern Carpathians.

⁶⁶ In several overviews of endemic plants (Stojko & Tasenkewitsch 1991, Stoyko & Tasenkevich 1993, Kricsfalussy & Budnikov 2002) *Dianthus carthusianorum* subsp. *tenuifolius* is reported also from the Ukrainian Carpathians. According to Fedorončuk & Čornej (2005), the occurrence of this subspecies does not extend up to Ukraine and the data from the Ukrainian Carpathians are erroneous (cf. Čornej 2011). On the other hand, Kuz'mina (2004) reports that its occurrence in the Ukrainian Carpathians is documented by a single herbarium specimen (Marmaroš Mts, Mt. Pop Ivan, Popov 1946).

⁶⁷ In older literature *Dianthus praecox* subsp. *praecox* was reported using nomenclatorically incorrect names *D. praecox* Kit. subsp. *praecox*, *D. plumarius* subsp. *praecox* (Kit. ex Schultes) Domin and *D. plumarius* subsp. *praecox* (Kit.) Pawł.

⁶⁸ To *Dianthus *praecox* also *D. hungaricus* subsp. *pseudopraecox* was included by Jalas & Suominen (1986), although its taxonomic position they consider to be controversial; in accord with this concept they report *Dianthus *praecox* also from Hungary.

⁶⁹ Baksay (1972) reported the occurrence of *Dianthus praecox* subsp. *lumnitzeri* also from the Pilis Mts in the northern Hungary (cf. Kmeťová 1985). According to current molecular studies records of this subspecies from Hungary should be referred to *Dianthus plumarius* subsp. *regis-stephani* (Rapaics) Baksay (Somogyi et al. 2012, Barina 2014 in litt.).

⁷⁰ *Dianthus praecox* subsp. *pseudopraecox* is a rare subspecies, occurring only in the Slovenský kras Karst including its Hungarian part (Aggteleki karszt Karst), and sporadically also in the Bükk Mts (Kmeťová 1985, 2012).

⁷¹ Baksay (1972) reported *Dianthus plumarius* subsp. *praecox* from the northwestern Hungary. According to Kmeťová (1985), these data are erroneous and concern subsp. *pseudopraecox* (cf. Kmeťová 2012).

⁷² *Dianthus spiculifolius* was reported (Klokov 1952, Prokudin 1987, Tasenkevych 2003b) also from the Ukrainian Carpathians (the southern part of Bukovina County), although with certain ambiguity in some studies (Fedorončuk & Diduch 2002b, Kuz'mina 2004). According to Čornej (2011), the data from Ukraine are erroneous and the species grows only in the Romanian part of Bukovina County.

⁷³ Occurrence of *Doronicum carpaticum* is reported also from the Apuseni Carpathians (Oprea 2005), Bulgaria (Pawłowski 1970) and Serbia (Greuter in Euro+Med). According to Álvarez Fernández (2003), its distribution area is restricted only to the Carpathians; according to Pachswöll (2013), it grows only in the Eastern and Southern Carpathians in the region spreading from mountain ranges Čornohora and Svydovec in Ukraine to the Țarcu Mts and Retezat Mts in Romania. Records from the Apuseni Carpathians in Romania, and from Bulgaria and Serbia refer a closely relative species *Doronicum columnae* Ten.

⁷⁴ Several authors (Oprea 2005, Dihoru & Negrean 2009, Sârbu et al. 2013, and others) report the occurrence of *Draba dorneri* also from the Făgăraș Mts. This record was not confirmed and it probably concerns *Draba kotschyi* (Ion 2014 in litt.).

⁷⁵ Older records of *Draba kotschyi* from the Alps concern *Draba norvegica* Gunnerus (Hurdu et al. 2012b; cf. Walters & Akeyrod 1993, Jalas et al. 1996).

- ⁷⁶ In the included literature *Draba simonkaiana* was reported also as *D. stellata* subsp. *simonkaiana* (Jáv.) Beldie.
- ⁷⁷ In the included literature *Erigeron hungaricus* is wrongly reported also as *Erigeron neglectus* A. Kern.
- ⁷⁸ The results of molecular analyses (Şuteu 2012) proved that the Carpathian (*Eritrichium jankae*) and Alpine populations (*Eritrichium nanum*) are hardly genetically differentiated, however, they are well isolated biogeographically (cf. Tribsch & Schönswetter 2003).
- ⁷⁹ *Erysimum hungaricum* is a very rare species, since its description (Zapałowicz 1913) known only from the limestone rocks on the southern slope of the borderline Mt. Malyj Lostun (Čyvčyny Mts) / (Maramureş Mts, Mt. Lostun Mic), already in the Romanian part of the mountain range (Kobiv 2010). According to Sârbu et al. (2013), its occurrence in Romania needs to be confirmed (cf. Oprea 2005, Ciocârlan 2009).
- ⁸⁰ Tasenkevich (2002) considers *Erysimum hungaricum* to be subendemic to the Alps and Carpathians. This evaluation is based on a broader species concept (*E. hungaricum* incl. *E. wahlenbergii*) and enlargement of its area to the Austrian Alps (cf. Melzer & Polatschek 1971).
- ⁸¹ Based probably on the same chromosome number (2n = 14), invalidly described *Erysimum vagicum* Holub et Tomšovic (syn. *E. witmannii* subsp. *vagicum* (Holub et Tomšovic) Dostál, nom. inval.) was also often included to *E. witmannii* although it was evaluated by Michalková (2002) as a diploid cytotype of *Erysimum odoratum* Ehrh.
- ⁸² Reports on occurrence of *Erysimum witmannii* subsp. *witmannii* in the Carpathians are very conflicting. Some authors, e.g. Konětopský (1963) and Čopyk (1976) reduce the distribution of *E. witmannii* s. str. only to the Western Carpathians, others (e.g. Beldie 1977, Ciocârlan 2009, Sârbu et al. 2013) only to the Eastern Carpathians. Nyárády (1955) and Tomšovic (1988) report it also from the Southern Carpathians (Făgăraş Mts).
- ⁸³ Borza (1964) published the occurrence of *Erysimum baumgartenianum* from Cheile Gâlzii (Trascău Mts). It is the single known record of *Erysimum witmannii* in the Apuseni Carpathians.
- ⁸⁴ The older records of *Erysimum baumgartenianum* Schur from Bulgaria concern *Erysimum cuspidatum* (M. Bieb.) DC. (cf. Ančev & Polatschek 2006).
- ⁸⁵ In the newer Romanian handbooks (Beldie 1977, Ciocârlan 2009, Sârbu et al. 2013) the distribution area of *Erysimum witmannii* subsp. *transsilvanicum* is restricted to the Southern Carpathians. However, Nyárády (1955) reported its occurrence also from the Eastern Carpathians; currently it was reported e.g. by Oprea & Sîrbu (2012, 2013) and Vojtkó et al. (2012). Tomšovic (1988) delimited its distribution in Romania to the Eastern and Southern Carpathians, from the Maramureş Mts to the Retezat Mts.
- ⁸⁶ From the Ukrainian Carpathians a single occurrence of *Erysimum witmannii* subsp. *transsilvanicum* is known from Mt. Velykyj Kamiň in the Čyvčyny Mts (Čopyk 1976, Il'jins'ka et al. 2007, Kobiv 2010).
- ⁸⁷ *Euphorbia carpatica* has centre of its distribution in the Ukrainian Carpathians. In the Romanian Eastern Carpathians it is known only from Oaş and Gutâi Mts (Dihoru & Negrean 2009). From Poland (Bieszczady Mts) it was reported by an error (cf. Mirek et al. 2002); the data concern *Euphorbia sojakii*.
- ⁸⁸ Malynovs'kyj et al. (2002) reported occurrence of *Euphorbia carpatica* using an invalid name *Euphorbia jasiewiczii* (Chrtek et Křísa) Dubovik.
- ⁸⁹ For a long time, narrow endemic *Euphrasia exaristata* has been known only from the Slovak part of the Červené vrchy Massif (Západné Tatry Mts), however, recently it was discovered also in its Polish part (Staszkievicz 2009).
- ⁹⁰ Our evaluation of endemic status of *Euphrasia tatarae* is based on Smejkal (1963) and Smejkal & Čerovský (1999), where the species distribution area is restricted to the Western Carpathians and the western part of the Eastern Carpathians. In Romanian literature (Răvăruţ 1960, Beldie 1967b, Oprea 2005), the species is reported from the Romanian Eastern (Maramureş, Rodna, Ceahlău & Rarău Mts) and Southern Carpathians (Bucegi & Retezat Mts); however, Mihoková & Mikoláš (1994) question the records from the Southern Carpathians (possibly confusion with *Euphrasia minima* Jacq. ex DC.). Records from the Krkonoše Mts concern local narrow endemic *Euphrasia corcontica* (Smejkal) Smejkal et Dvořáková (Dvořáková 1999).

- ⁹¹ According to the current data (Lendvay & Kalapos 2014), *Ferula sadlerana* occurs recently in seven localities, four of them located in the Western Carpathians, one in the Apuseni Carpathians and two outside the Carpathians in the northern part of the Transdanubian Mountains in Hungary.
- ⁹² In older (not included) studies, *Ferula sadlerana* is usually evaluated as subendemic to the Pannonian Basin.
- ⁹³ In several sources (e.g. Markgraf-Dannenberg 1980, Oprea 2005, Ciocârlan 2009, Sârbu et al. 2013), the occurrence of *Festuca amethystina* subsp. *orientalis* is reported also from the western part of the Balkan Peninsula. However, this taxon was not included in recent Balkan floras neither as species nor as subspecies (Krahulec 2014 in litt.).
- ⁹⁴ In the included literature *Festuca saxatilis* is reported also using invalid name *Festuca rupicola* subsp. *saxatilis* (Schur) Jáv.
- ⁹⁵ Records of *Festuca tatrae* from the Ukrainian Carpathians are erroneous (Čornej 2011, Kobiv 2014 in litt., Tasenkevich 2014 in litt., Kobiv 2014 in litt.). Beldie (1967b) reported scarce occurrence of *Festuca amethystina* subsp. *tatrae* from the Bucegi Mts in the Southern Carpathians (cf. Beldie 1972). Ciocârlan (2009) considered this record to be erroneous; his opinion is shared also by other Romanian botanists (Hurdu 2014 in litt.).
- ⁹⁶ Outside the Carpathians *Festuca versicolor* subsp. *versicolor* occurs scarcely in the Bohemian and Polish part of the Krkonoše Mts (see Kliment 1999 for details).
- ⁹⁷ The eligibility of delimitation of the subspecies *Galium album* subsp. *suberectum* was questioned in some comprehensive studies (cf. Ehrendorfer & Krendl 1976); but according to the current results (Gynda 2004), it differs from subsp. *album* morphologically and ecologically, and these two subspecies are reproductively isolated.
- ⁹⁸ In the present overviews (Čornej 2011, Tasenkevich 2011) *Galium album* subsp. *suberectum* was evaluated as endemic to the Eastern Carpathians. Michalková (1993) reported this subspecies from several mountain ranges and basins in the Slovak part of the Western Carpathians.
- ⁹⁹ Outside the Carpathians *Galium kitaibelianum* scarcely occurs also in Serbia: Velika Remeta, Krušedol (Hurdu et al. 2012b). In several studies (e.g. Popescu & Sanda 1998, Oprea 2005) is it considered to be a Carpathian-Balkan species.
- ¹⁰⁰ In the included literature *Genista tinctoria* subsp. *oligosperma* (Andrae) Jáv. is reported also using nom. illeg. (younger homonyma) *G. tinctoria* subsp. *oligosperma* (Andrae) Prodan, *G. tinctoria* subsp. *oligosperma* (Andrae) Borza and *G. tinctoria* subsp. *oligosperma* (Andrae) Soó.
- ¹⁰¹ Records of *Gentiana cruciata* subsp. *phlogifolia* from Greece (Falakron Mts) are erroneous; according to Strid & Tan (1991), the populations of the place belong to subsp. *cruciata*.
- ¹⁰² Tutin (1972) identified *Gentiana laciniata* with *Gentiana pyrenaica* L., which is distributed from the Pyrenees through the Carpathians to the mountains of the southwestern Bulgaria. Current study by Rybczyński et al. (2014) confirmed the eligibility of delimitation of *G. laciniata* as a separate species, narrow endemic to the Ukrainian Carpathians (cf. Čopyk 1976, Kricsfalussy 1999, Malynovs'kyj et al. 2002, Antosyak & Kozurak 2011) with a scarce occurrence in the Čornohora, Svydovec and Boržava Mts. Based on explicit differences in the length and width of sepal teeth, both species were distinguished already by Cvelev (1978).
- ¹⁰³ Records of *Gypsophila petraea* from Bulgaria (Rodopi Mts) are erroneous (cf. Jalas & Suominen 1986, Hurdu et al. 2012b).
- ¹⁰⁴ Taxonomic concept of the genus *Hesperis* is based on a monography by Dvořák (1968), with regards to the current nomenclatoric information.
- ¹⁰⁵ In the description of *Hesperis dinarica* subsp. *slovaca*, Dvořák (1963) along with the Nízke Tatry Mts wrongly reported also the locality „Transsilvania, Carpati Orientales, comit. Máramaros, mons Nagy Pietrosz“. In his monography (Dvořák 1968), *Hesperis slovaca* is already reported only from the Nízke Tatry Mts as a local endemite.
- ¹⁰⁶ The nomenclature of the species of the genus *Hieracium* is in accordance with current taxonomic studies, in the section *Cernua* following mainly Szeląg (2003b).
- ¹⁰⁷ Further synonyms of *Hieracium borbasii* are as follows: *Hieracium evolutum* (Nyár. et Zahn) Zahn; *H. pseudokotschyanum* (Nyár. et Zahn) Nyár.; *H. pseudotubulare* (Nyár. et Zahn) Nyár.; *H. sublubricicaule* (Nyár. et Zahn) Nyár. (cf. Szeląg 2006b).
- ¹⁰⁸ In the included literature *Hieracium fagarasense* is also reported as follows: *Hieracium kotschyanum* subsp. *fagarasense* (Nyár. et Zahn) Beldie; *H. sparsum* subsp. *borbasii* var. *fagarasense* (Nyár. et Zahn) Ciocârlan.

- ¹⁰⁹ In the included literature *Hieracium negoiense* is also reported using invalid name *Hieracium krasanii* var. *negoiense* (Răvărut et Nyár.) Beldie et L. Alex.
- ¹¹⁰ *Hieracium ostii-bucurae* is a narrow endemic known from only three adjacent mountain ranges Retezat, Țarcu and Godeanu Mts (cf. Szeląg 2006b).
- ¹¹¹ The tetraploid chromosome number ($2n = 36$) reported for *Hieracium longifoliosum* in Mráz (2006) concern *Hieracium polyphyllobasis* (Szeląg 2006b).
- ¹¹² Zahn (1936) reported occurrence of *Hieracium pietroszense* besides of *locus classicus* (Rodna Mts: Mt. Pietrosu) also from several sites at high altitudes in the Southern Carpathians (Retezat Mts) and the Alps. According to the current knowledge (Mráz 2003a), it occurs only in the Eastern Carpathians (Rodna Mts). Records from the Southern Carpathians were confirmed as erroneous based on the study of herbarium material; the occurrence in the Alps is very improbable.
- ¹¹³ *Hieracium rapunculoidiforme* was described from Mt. Pietrosz (Ukraine: Čornohora Mts, Mt. Petros) on the border of former Kingdom of Hungary and Galicia region (Zahn 1911). It is considered to be endemic to the Eastern Carpathians in numerous Ukrainian and Romanian overviews. Nyárády (1965) located *locus classicus* wrongly to Mt. Pietrosu in the Maramureș Mts. Szeląg (2014 in litt.) expects species occurrence also in the Romanian Eastern Carpathians.
- ¹¹⁴ Record of *Hieracium scitulum* from the other Carpathian subunits concern *H. scitulum* s. lat. (Szeląg 2014 in litt.); records from the Krkonoše Mts are erroneous (Chrtěk jr. 2004).
- ¹¹⁵ Outside the Western Carpathians *Hieracium silesiacum* recently scarcely occurs only in the Hrubý Jeseník Mts (Velká kotlina Cirque) (Szeląg 2004b).
- ¹¹⁶ Record of *Hieracium tubulare* in the Harghita Mts (Mráz & Szeląg 2004) concerns *Hieracium coldei* (Szeląg 2006a).
- ¹¹⁷ The occurrence of *Hieracium virgicale* outside the Western Carpathians is reported by a single historical evidence from the Maramureș Mts in the Eastern Carpathians (Zahn 1930 ut *H. virgicale* subsp. *nudatum*). Šljakov (1989) does not report this species from Ukraine; it was also not confirmed in the Romanian Eastern Carpathians (Nyárády 1965, Oprea 2005). Chrtěk et al. (2004a) considered the above-mentioned record as dubious.
- ¹¹⁸ In Poland *Jovibarba globifera* subsp. *preissiana* scarcely occurs also outside the Carpathians (Letz 1998).
- ¹¹⁹ *Jurinea transylvanica* has a centre of its distribution in the Transylvanian Basin. Sporadically, it was found also in the Romanian Eastern (Hășmaș Mts), Southern (Retezat Mts) and Apuseni Carpathians (Cheile Turzii Gorge); several localities are also in jud. Galați, already outside the Carpathians (Oprea 2005). Records from Bulgaria are erroneous (Greuter in Euro+Med).
- ¹²⁰ Outside the Western Carpathians *Knautia kitaibelii* subsp. *kitaibelii* occurs rarely in the Záhorská nížina Basin (Štěpánek 1985). Böhm & Facsar (2000) reported its sporadic occurrence from the Pilis Mts in Hungary, together with subsp. *tomentella* (Szabó) Baksay.
- ¹²¹ Up to this day, *Knautia kitaibelii* has been reported also from Ukraine and Romania (currently e.g. Kricsfalussy & Budnikov 2002, 2007, Ciocârlan 2009). The distribution area of *K. kitaibelii* towards the east reaches the Pieniny Mts, records from the Ukrainian and Romanian Carpathians are erroneous (cf. Soják 1983b, Štěpánek 1985, 1997).
- ¹²² Deyl (1934), and based on his collections later also Soják (1983b), reported *Koeleria macrantha* subsp. *transsilvanica* also from the Ukrainian Carpathians, from where its occurrence was later not confirmed (cf. Kricsfalussy & Budnikov 2007).
- ¹²³ From the Transylvanian Basin *Lathyrus transsilvanicus* sporadically spreads beyond to the Cheile Turzii Gorge on the eastern margin of the Apuseni Carpathians (Oprea 2005).
- ¹²⁴ In 1903, *Lathyrus transsilvanicus* was collected in a beech forest in the Armankaja River Valley (Sredna Stara planina Mts), ca 1 600 m a.s.l., where it has not been confirmed since 1905 and was considered to be extinct in Bulgaria (Petrova & Vladimirov 2009, Tosheva et al. 2011). During 2007–2011 it was recorded in three near microlocalities (Triglav Massif, seven gorges locality) in the Sredna Stara Planina Mts, 1 450–1 500 m a.s.l. (Marinov et al. 2014).
- ¹²⁵ In the included literature it is also reported as *Lathyrus transsilvanicus* (Spreng.) Rchb. f.
- ¹²⁶ Holub (1977b) constrained the occurrence of diploid endemic subspecies *Leucanthemopsis alpina* subsp. *tatrae* only to the Tatry Mts with a note that the relations of the Tatry populations with populations of *Leucanthemopsis alpina* from the Eastern and Southern Carpathians require further study. *Leucanthemopsis *tatrae* was evaluated as endemic to the

Tatry Mts e.g. by Čopyk (1976), Kliment (1999), Piękoś-Mirkowa & Mirek (2003) and Mirek & Piękoś-Mirkowa (2009, 2010). Oprea (2005) reported it from the Romanian Eastern and Southern Carpathians remarking that it is a single subspecies growing in the Romanian Carpathians. Without a detailed information it was reported also by Mosyakin & Fedoronchuk (1999) from the Ukrainian Carpathians and by Sârbu et al. (2013) from the Romanian Carpathians. Further authors, e.g. Popescu & Sanda (1998) and Ciocârlan (1990, 2009), report from Romania only the species (using diverse names) without information on its distribution. According to the current information, reliable records of *Leucanthemopsis alpina* at the subspecies level are missing as from the Ukrainian Carpathians (Kozurak 2014 in litt.), so from the Romanian Carpathians (Puşcaş 2014 in litt.). Until the taxonomic affiliation of *L. alpina* populations from the Ukrainian and Romanian Carpathians will be resolved, we retain the present endemic status of *Leucanthemopsis *tatrae*.

¹²⁷ Outside the Carpathians *Leucanthemum rotundifolium* sporadically occurs in the Vranica Planina Mts in Bosnia; an older record from the southern Croatia proved to be erroneous (Zelený 1970, Kliment 1999, and others).

¹²⁸ Outside the Carpathians *Linum extraaxillare* very scarcely occurs only in Bulgaria. In the localities Mt. Karlovska (Centralna Stara planina Mts) and Mt. Vitoša (Vitoša Mts) its occurrence have not been confirmed for more than 80 years, therefore the species was considered to be extinct in Bulgaria (Petrova & Vladimirov 2009). In 1995, it was collected on Mt. Kucheto in the Centralna Stara planina Mts (Vladimirov et al. 2011).

¹²⁹ Outside the Carpathians *Linum uninerve* occurs very scarcely (including only three localities) in the Centralna Stara planina Mts (Kuru Dere) and the Rodopi Mts (Dobrostan Massif), in small populations with restricted reproductive ability (Petrova 2011). In Romania it was found in the vicinity of the city Lugoj at the border of the Southern Carpathians (Poiana Ruscă Mts) and the Pannonian Basin (Dihoru & Pârvu 1987).

¹³⁰ In the included literature it is also reported as *Linum uninerve* (Rochel) Borbás.

¹³¹ *Melampyrum saxosum* (with white-coloured petals) and *M. herbichii* (with yellow-coloured petals) were so-far evaluated as two separate taxa (microspecies). Těšitel et al. (2009) in their thorough morphometric and molecular study did not find traits, which would differentiate between the white-coloured and the yellow-coloured populations; these also do not differ in their geographical distribution. Therefore they suggested to consider both microspecies as a single species with an older valid name *Melampyrum saxosum* Baumg.

¹³² Towards the west, an introgressive hybridization between yellow-flowering plants and *Melampyrum sylvaticum* populations probably occurs, which hindered identification of the western border of *M. herbichii* distribution (cf. Štech & Drábková 2005, Těšitel & Štech 2007). As a consequence, the species was evaluated variously in the present overviews of endemics: from an endemic to the Eastern and Southern Carpathians to a pan-Carpathian subendemic or a Sudeten-Carpathian species. Similarly to other east-Carpathian species (s. l.), also in this case the biogeographic border between the Western and Eastern (South-Eastern) Carpathians is of special importance, and the typical populations of *Melampyrum saxosum* (incl. *M. herbichii*) do not extend over this border (for details see Těšitel et al. 2009).

¹³³ In the included literature *Minuartia pauciflora* is wrongly reported as *Minuartia verna* subsp. *gerardii* (Willd.) Graebn.; this name concerns the alpine plants.

¹³⁴ In the included literature *Noccaea banatica* is reported using an invalid name *Thlaspi dacicum* subsp. *banaticum* (R. Uechtr.) Jáv. (cf. Holub 1984).

¹³⁵ Records of *Noccaea jankae* in Romania (Beldie & Váczy 1976, Beldie 1977, Dihoru & Negrean 2009, Sârbu et al. 2013, and others) concern probably a Carpathian-Balkan species *Noccaea kovatsii* (Heuff.) F. K. Mey. (*T. jankae* Borbás non A. Kern. and *T. jankae* Velen. non A. Kern. also belong to synonyms of *N. kovatsii*; cf. Čopyk 1976, Ančev 2007).

¹³⁶ Outside the Transylvanian Basin *Onosma pseudarenaria* subsp. *pseudarenaria* sporadically occurs also in the Eastern (Hășmaș Mts) and Apuseni Carpathians (Trascău Mts: Cheile Turzii Gorge) (Oprea 2005).

¹³⁷ According to molecular analyses (Kolarčík et al. 2010), *Onosma viridis* occurs only in the western, south-western and southern parts of the Transylvanian Basin (mainly close to its border with the Apuseni and the Southern Carpathians), in the Apuseni Carpathians (Trascău Mts), in the southern part of the Banat Mts (cf. Grințescu & Nyárády 1960a), as well as in the Slovenský kras and Aggtelek Karsts (populations formerly reported as *O. tornensis*; see next note). In this respect it belongs to the taxa endemic to the Western, Southern and Apuseni Carpathians and the Transylvanian Basin. According to a broader concept (cf. Valdés in Euro+Med), it represents a Carpathian-Balkan species distributed from Serbia, Bulgaria and Greece to Turkey. The whole taxonomic complex of closely related species within *Onosma heterophylla* s. lat. (see Kolarčík et al. 2010, 2014 for details), requires deeper taxonomic (and subsequently also chorological) revision.

¹³⁸ Since its description (Jávorka 1906), *Onosma tornensis* has been considered as one of the rarest stenoendemic of the Western Carpathians with a restricted occurrence in a small area in the eastern part of the Slovenský kras Karst including the Slovak-Hungarian plateau Dolný vrch. With regard to its rarity it was included in the World Red Book of endangered species as well as to the international directives (Bern1, Annex II of Habitat Directive 92/43/EEC). The recent study of molecular variation (Kolarčík et al. 2010) confirmed significant DNA similarity of *Onosma tornensis* populations from Slovakia with *Onosma viridis* populations from the Banat part of the Southern Carpathians and taxonomic identity of both species. Regarding this fact, Mártonfi et al. (2014), in accordance with regulations of the International Code of Nomenclature for algae, fungi, and plants (McNeill et al. 2012, Art. 11.5), reports *O. tornensis* as synonym of *O. viridis*.

¹³⁹ So-far *Ophrys holubyana* have been reported as subendemic to the Western Carpathians. Records from the northern Hungary (cf. Kliment 1999) concern recent hybrids of *Ophrys holoserica* (Burm. f.) Greuter and *Ophrys scolopax* subsp. *cornuta* (Steven) E. G. Camus (Dítě 2014 in litt.).

¹⁴⁰ In the Eastern Carpathians *Pedicularis baumgartenii* is reported only from the Ceahlău Mts (Dihoru & Pârvu 1987, Ciocârlan 2009, Sârbu et al. 2013).

¹⁴¹ *Peucedanum rochelianum* is a species of wet aluvial meadows (*Peucedano rocheliani-Molinietum caeruleae*) and forests (*Quercus-Betuletum molinietosum*). The centre of its distribution is in the sub-Carpathian depressions in the south-western part of the Transylvanian Basin wedged in the adjacent mountain ranges of the Southern Carpathians (Caransebeș Depression, Hațeg Depression, Sibiu Depression), at altitudes between 230 and 400 m (Boșcaiu 1965, Boșcaiu & Rațiu 1965, Kovács 2011). Records from the central and eastern Transylvania (Dihoru & Pârvu 1987, Jakab et al. 2008) can be dubious (Kovács l. c.). Sporadic occurrence was reported also in the forests near the villages Mavrodolu and Plopeni at the southern foothill of the Southern Carpathians (Paucă-Comanescu & Negrean 1994) and in the surroundings of Galați city in the south-eastern Romania (Dihoru & Pârvu 1987).

¹⁴² Kovács (2011), with the reference to Boșcaiu (1965), reports occurrence of *Peucedanum rochelianum* also from Croatia. However, Boșcaiu (l. c.) reported from the Croatian Dalmatia occurrence of a vicariant taxon *Peucedanum coriaceum* subsp. *pospichalii* (Thell.) Horvatić and association *Peucedano pospichalii-Molinietum litoralis*. *Peucedanum rochelianum* was not reported in the relevant Croatian literature (Krahulec 2014 in litt.).

¹⁴³ *Pilosella ullepitschii* occurs predominantly in the Western Carpathians; only scarcely it occurs in the Eastern (three localities) and Southern (one locality) Carpathians (Šingliarová & Mráz 2009).

¹⁴⁴ Bernátová et al. (2006) distinguished two subspecies within *Poa carpatica*: subsp. *carpatica* (endemic to the Western and Eastern Carpathians) and subsp. *supramontana* Bernátová et al. (endemic to the Veľká Fatra Mts, Krivánska Fatra Mts and Nízke Tatry Mts). However, the later mentioned subspecies was invalidly described because the original diagnosis lacked specification of herbarium collection or institution in which holotype specimen was conserved (Art. 40.7 of Melbourne Code, McNeill et al. 2012).

¹⁴⁵ In older Romanian sources (Ghișa & Beldie 1972) also *Poa deylii* var. *retezatensis* (A. Nyár.) Ghișa et Beldie (only Retezat Mts) is recorded, evaluated as a separate taxon (*Poa granitica* subsp. *retezatensis* A. Nyár) also by Greuter (in Euro+Med). However, in the current Romanian identification keys (Ciocârlan 2009, Sârbu et al. 2013) only occurrence of *Poa granitica* subsp. *disparilis* is reported.

¹⁴⁶ *Poa nobilis* is an alpine hybridogenous species (probably *Poa alpina* f. *vivipara* × *Poa granitica*) described from the Polish side of the Tatra Mts (Skalińska 1955), which was stabilised thanks to its vivipary (Pogan 1977). During the field survey in the nineties of the last century, its occurrence has not been confirmed (Piękoś-Mirkowa et al. 1996), which is reflected also in the current evaluation within the category DD (Piękoś-Mirkowa 2008).

¹⁴⁷ *Poa pannonica* subsp. *scabra* has centre of its distribution on neutral and alkaline substrates (volcanic rocks and limestones) in the Slovak part of the Western Carpathians with outstrokes towards the south in the Északi-középhegység Mts and in the northern part of the Dunántúli-középhegység Mts (Visegrádi Mts). In the Ukrainian Carpathians only older records exist from slopes of Chorna Hora near the town Sevluš (Vinohradiv) (Jirásek 1934, 1935); in more recent floras (Čopyk 1976, etc.) it was not reported any more. In Romanian Carpathians the subspecies was reported near the Iron Gates Gorge; older record is from the vicinity of Arad City (Oprea 2005).

¹⁴⁸ In the included literature it is usually reported as *Poa rehmannii* (Asch. et Graebn.) Woł. As a synonym of *Poa nemoralis* B. [subsp.] *rehmannii* Asch. et Graebn. Ascherson & Graebner (1900: 413) reported the name *Poa caesia* d) *Rehmanni* Richter Pl. Eur. I. 83 (1890), which is a reference to the first published name and should be regarded as a bazionym (Marhold 2014 pers. comm.).

- ¹⁴⁹ *Prangos carinata* is a rare species known only from the Iron Gates Gorge (between the locality of Vîrciorova and the village of Gura Văii) in the Southern Carpathians (Morariu & Beldie 1976, Beldie 1977, Dihoru & Negrean 2009).
- ¹⁵⁰ In the included literature it is reported also under invalid name *Prangos carinata* Griseb.
- ¹⁵¹ Outside the Western Carpathians *Primula auricula* subsp. *hungarica* sporadically occurs on dolomites in some mountain ranges (Vértes, Bakony, Balaton-vidék) of the Transdanubian Mountains in Hungary (Soó 1964, 1980a, Simon 1992, and others).
- ¹⁵² *Primula leucophylla* is an alpine taxon of the *Primula elatior* group adapted to a cool climate, ecologically, phenologically and genetically isolated from *P. elatior* s. str. Its occurrence is restricted to the limestone mountain ranges in the central part of the Eastern Carpathians, from the Rarău Mts to the Vrancei Mts (Şuteu 2012, Şuteu et al. 2011, 2013, Hurdu 2014 in litt.; cf. Ciocârlan 2009, Sârbu et al. 2013).
- ¹⁵³ According to Şuteu (2012), „Studies based on sequences analysis indicate that the taxon *P. filarszkyana* should be considered a subspecies of the species *Pulmonaria rubra*“. Therefore, in our study we included the taxon at the level of subspecies.
- ¹⁵⁴ The West-Carpathian endemic *Pulsatilla slavica* was reported also from the north-eastern Romania (Mîrzesti) by some Romanian authors (Beldie & Váczy 1976, Beldie 1977, Sanda et al. 1983, Ciocârlan 1988, Popescu & Sanda 1998). Goliašová (1985) considered its occurrence in Romania as unlikely. Ciocârlan (2009) evaluated the published records as erroneous.
- ¹⁵⁵ *Ranunculus flabellifolius* is a narrow species of the *Ranunculus auricomus* group with rather ambiguous taxonomic evaluation. While e.g. Jalas & Suominen (1989), Stevanović et al. (1991) and Dunkel (2011) considered it as a separate species of this group (cf. Beldie 1977, Popescu & Sanda 1998, Ciocârlan 2009, Sârbu et al. 2013, Hörandl & Raab-Straube in Euro+Med, and others), Tutin & Akeroyd (1993) included it within *Ranunculus cassubicus* L. (cf. Oprea 2005). According to Dunkel (2011), *Ranunculus flabellifolius* represents the corner stone of the whole complex. In our study, we therefore accept *R. flabellifolius* as a separate species.
- ¹⁵⁶ Outside the Carpathians, *Ranunculus pseudomontanus* sporadically occurs in the Bulgarian mountain ranges Vitoša and Zapadni Rodopi Mts (Kožuharov & Petrova 1988).
- ¹⁵⁷ In the included literature it is also reported under invalid name *Ranunculus montanus* subsp. *pseudomontanus* (Schur) Beldie.
- ¹⁵⁸ *Rosa coziae* represents one of four separate species of the *Rosa villosa* agg. (Kerényi-Nagy 2011), restricted by its occurrence to the Cozia and Căpătâinii Mts in the Southern Carpathians (Oprea 2005, Ciocârlan 2009, Hurdu et al. 2012a).
- ¹⁵⁹ In the included literature it is also reported using invalid names *Rosa villosa* subsp. *coziae* (Nyár.) Beldie and *R. villosa* subsp. *coziae* (Nyár.) Beldie et L. Alex.
- ¹⁶⁰ In the Southern Carpathians the occurrence of *Salvia transsylvanica* is known only from the vicinity of the village Boteni (Dihoru & Pârnu 1987, Oprea 2005).
- ¹⁶¹ Current distribution of *Saussurea porcii*, a narrow endemic species of the Eastern Carpathians, was studied by Kobiv et al. (2007b), later also by Počynok & Prokopiv (2010), and its biology was studied by Bahlej (2010). Počynok & Prokopiv (l. c.) reported its occurrence on seven confirmed localities in the Ukrainian mountain ranges: Čyvčyny Mts (five localities), Čornohora Mts (one locality) and Svydovec Mts (one locality). In the Rodna Mts (Romania) the species was considered extinct (cf. Dihoru & Negrean 2009). The occurrence not confirmed since 1902 has been verified by Mátiš, Szabó & Bartha in August 2014 (Mátiš et al. 2014).
- ¹⁶² Diklić (1973) reported *Scabiosa columbaria* subsp. *banatica* from the eastern and south-eastern part of Serbia and from Vojvodina; Hurdu et al. (2012b) delimited its occurrence in Serbia to the Serbian part of the Southern Carpathians and ranked it as endemic to the South-Eastern Carpathians.
- ¹⁶³ In Ukraine and Hungary *Scilla kladnii* sporadically occurs also outside the Carpathians (cf. Speta 1977, Kereszty et al. 1986, Kereszty 1993, Kricsfalusy & Vajnagi 1994).
- ¹⁶⁴ In the included literature *Sempervivum carpathicum* is reported also as *Sempervivum wettsteinii* Letz ined.; wrongly also as *Sempervivum montanum* L.
- ¹⁶⁵ In the included literature *Sempervivum carpathicum* subsp. *carpathicum* is reported also as *Sempervivum wettsteinii* Letz ined. subsp. *wettsteinii*.
- ¹⁶⁶ In the included literature *Sempervivum carpathicum* subsp. *heterophyllum* is reported also as *Sempervivum wettsteinii* subsp. *heterophyllum* (Hazsl.) Letz ined.

- ¹⁶⁷ Outside the Carpathians *Sempervivum matricum* grows also in the northern part of the Transdanubian Mountains in Hungary (Letz 2009).
- ¹⁶⁸ *Senecio ucranicus* is distributed from the Polish and Slovak part of the Eastern Carpathians to the Făgăraș Mts in the Southern Carpathians; in Ukraine it rarely outreaches to Volyn and Podillja, in Poland to Pogórze Dynowskie Mts at the border of the Eastern and Western Carpathians (Hodálová 1998b, Rola 2014).
- ¹⁶⁹ *Sesleria heuflerana* subsp. *heuflerana* extends rarely to Podillja from the Ukrainian Carpathians (Čopyk 1976) and to rocky limestone hillsides of Moldavian Priprutia from the Romanian Carpathians (Gejdeman 1986).
- ¹⁷⁰ In several surveys (e.g. Deyl 1980, Májovský, Murín et al. 1987, Dostál 1989, 1992) *Sesleria heuflerana* subsp. *hungarica* is reported also from Slovakia. The only occurrence of octoploid plants from Slovakia (Slovenský kras Karst) was evaluated as an octoploid cytotype of *S. heuflerana* by Lysák (1996).
- ¹⁷¹ Deyl (1980) reported the occurrence of *Sesleria rigida* also from the northern part of the Balkan Peninsula. According to the results of the molecular-taxonomic study by Kuzmanović et al. (2013), *S. rigida* s. str. occurs only in the Romanian Carpathians; records from the Balkan mountains concern closely related species of *Sesleria rigida* s. lat.: *S. filifolia* Hoppe, *S. serbica* (Adamović) Ujhelyi and *S. achtarovii* Deyl.
- ¹⁷² Outside the Western Carpathians *Sesleria tatrae* occurs very scarcely in a single locality in the Śnieżnik Kłodzki Massif in Poland (Fabiszewski 1970, Hendrych 1987, Kliment 1999, Budzáková et al. 2014, and others).
- ¹⁷³ In the Ukrainian Carpathians *Silene zawadzki* occurs rarely only in two localities (Velykyj Kamin' and Mokryniv Kamin') in the Čyvčyny Mts (Kobiv 2010).
- ¹⁷⁴ Based on Drăgulescu (2003), Oprea (2005) reported the occurrence of *Silene zawadzki* also from the Făgăraș Mts (Mt. Arpașu) in the Southern Carpathians although he himself considered this species to be endemic to the Eastern Carpathians. In later studies (Ciocârlan 2009, Hurdu et al. 2012a, Sârbu et al. 2013), this report was not considered.
- ¹⁷⁵ *Sorbus amici-petri* is a narrow endemic hybridogenous taxon described from sunny hills in the vicinity of Obišovce, Kysak and Trebejov Villages (Mikoláš 2003) in the phytogeographical district Stredné Pohornádíe (eastern Slovakia), geomorphologically ordered to the Čierna hora Mts (Mazúr & Lukniš 1980).
- ¹⁷⁶ *Sorbus dolomiticola* is a stenoendemic of the area between Kysak and Trebejov in the eastern Slovakia (Mikoláš 1996).
- ¹⁷⁷ *Sorbus haljamovae* is an alpine hybridogenous species in its occurrence restricted to relic calcareous dwarf mountain pine (*Pinus mugo*) communities in the Veľká Fatra, Krivánska Fatra and Nízke Tatry Mts (Bernátová & Májovský 2003).
- ¹⁷⁸ *Sorbus zuzanae* is an alpine hybridogenous species, in its occurrence restricted to relic calcareous dwarf mountain pine (*Pinus mugo*) communities in the Veľká Fatra, Krivánska Fatra and Nízke Tatry Mts (Bernátová & Májovský 2003).
- ¹⁷⁹ *Stipa crassiculmis* subsp. *heterotricha* is a subspecies with ambiguous taxonomic evaluation (cf. Ciocârlan 2009, Sârbu et al. 2013), however, considered as a separate taxon by Vázquez & Gutiérrez (2011) (cf. Popescu & Sanda 1998, Dihoru & Negrean 2009, Negrean 2011).
- ¹⁸⁰ Outside the Carpathians two small populations of *Swertia punctata* are recently known only from two adjacent localities (1 300 and 2 000 m a.s.l.) on Mt. Midžur (Zapadna Stara planina Mts) on the Bulgarian-Serbian border; it is also reported from Kosovo (Tan & Vladimirov 2001).
- ¹⁸¹ In Ukraine, *Symphytum cordatum* scarcely occurs also outside the Carpathians in the vicinity of Lviv as an isolated population in Volyn near the Sluč River (Čopyk 1976).
- ¹⁸² A narrow endemic *Taraxacum carpaticum* was up to now recorded only in two neighbouring mountain ranges, Bucegi Mts and Piatra Craiului Mts within the Southern Carpathians (Štěpánek et al. 2011).
- ¹⁸³ *Taraxacum nigricans* is an inconsistently evaluated species described from the Nízke Tatry Mts (Ďumbier Mt.), in the past referred also from the Polish, Ukrainian and Romanian parts of Carpathians, or from even larger area. The checking of the type material confirmed, that it is restricted to the region of original diagnosis (Štěpánek et al. 2011).

- ¹⁸⁴ Besides the classical locality in the Polish part of the Vysoké Tatry Mts, Tacik (1980) reports the occurrence of *Taraxacum pawlowskii* (he evaluates this occurrence as unsure) also from the Velická dolina Valley in the Slovak part of the mountain range.
- ¹⁸⁵ *Taraxacum pieninicum* is probably the best known stenoendemic species of the Pieniny Mts. Its occurrence is known only from the rocky walls of Mt. Okraǵlica in the Trzy Korony Massif in the Polish part of the Pieniny Mts. After extinction of the original population in its *locus classicus* (crash of the rocky wall), the species has been considered as missing for longer time (Zarzycki 1986 to Mirek et al. 1995). During a detailed field survey of the rocky walls in 1999–2000 it has been re-discovered under the top of Mt. Okraǵlica (Wróbel & Zarzycki 2008).
- ¹⁸⁶ Along with recently known occurrences in the Slovak, Ukrainian and Romanian Carpathians and their surroundings (see Kliment 1999 for details), a sporadic occurrence of *Thymus alternans* documented by herbarium specimens is known also from the Southern (Făgăraş Mts: Nucşoara, Mártonfi 2014 in litt.) and Apuseni Carpathians (Bihor Mts; cf. Kliment 1999). Our recent evaluation is most compatible with the evaluation by Čornej (2011), who denoted *Thymus alternans* as subendemic to the Carpathians.
- ¹⁸⁷ Outside the Carpathians *Thymus dacicus* sporadically occurs in Moldova (Iaşi City) and in the Walachian (Câmpia Română) Basin (Guşuleac 1961, Diklić & Vasić 2000).
- ¹⁸⁸ Outside the Western Carpathians *Thymus pulcherrimus* subsp. *sudeticus* occurs in two isolated localities in the Hrubý Jeseník Mts.
- ¹⁸⁹ In the included literature *Thymus pulcherrimus* subsp. *sudeticus* was incorrectly reported as *Thymus macrophyllus* Rchb.
- ¹⁹⁰ In the Romanian literature *Trifolium pratense* subsp. *kotulae* is reported as *T. pratense* subsp. *nivale* (Sieber) Arcang. and *T. pratense* var. *frigidum* Gaudin.
- ¹⁹¹ From the Carpathians, *Trifolium sarosiense* sporadically reaches also the northern part of the Transdanubian Mountains, eastern part of the Pannonian Basin and south-eastern part of Serbia: Vršac, Bor, Pirot (Hendrych 1993, 1995); the author assumes also its occurrence in the Ukrainian Carpathians.
- ¹⁹² From the Ukrainian Carpathians *Trisetum flavescens* subsp. *tatricum* is documented only by older herbarium specimens (K. Igošina 1948 LE) from Mt. Petros in the Čornohora Mts (Cvelev 1974, 1976, Prokudin et al. 1977, Čornej 2011).
- ¹⁹³ In Serbia a single locality of *Tulipa hungarica* was known from the southern side of the Iron Gate Gorge (Đerdap National Park; cf. Čalić et al. 2012); recently the species is evaluated as extinct in Serbia (Stevanović 2013, Turis et al. 2014).
- ¹⁹⁴ Oprea (2005) incorrectly reported *Tulipa hungarica* var. *undulatifolia* Roman as synonym of *Tulipa undulatifolia* Boiss. (cf. Negrean 2011).
- ¹⁹⁵ Rostański (1967, 1970) ordered to *Valeriana tripteris* subsp. *heterophylla* plants with at least some stem leaves composed of 5 leaflets, with a large oblong-ovate terminal divisions and about 0.5 mm long hairs on the leaf margin. From the nominate subspecies it differs also ecologically; it occurs in springs, aluvial alder forests, moist habitats along the mountain brooks and in *Alnus viridis* stands (Schur 1866, Dmytrach 2010). It is accepted as a separate subspecies e.g. by Mirek et al. (2002), and as a separate species (ut *Valeriana transsilvanica* Schur) by Katina (1961) and Prokudin (1987).
- ¹⁹⁶ Rostański (1967) reported the occurrence of *Valeriana tripteris* subsp. *heterophylla* from the Polish and Ukrainian part of the Eastern Carpathians. From the Polish Bieszczady Mts the subspecies sporadically protrudes to the neighbouring Beskid Niski Mts (Rostański 1967, 1970) ordered already to the Western Carpathians. Probably based on these facts, it was evaluated as endemic to the Western and Eastern Carpathians by Tasenkevych (2014). However, Morariu (1961 ut *V. tripteris* var. *heterophylla*) reported several localities of this taxon not only from the Eastern but also from the Southern and Apuseni Carpathians (cf. Schur 1866, Simonkai 1887 ut *Valeriana tripteris* var. *bijuga* Simk.).
- ¹⁹⁷ Kirschner (2007) ordered similar populations from the alluvial alder forests and springs in the Bukovské vrchy Mts to *Valeriana montana* var. *ternata* Schur. The occurrence of *Valeriana montana* L. (without closer specification) in the Bukovské vrchy Mts (springs and alluvia in the Zbojský potok Valley) was first reported by Hadač (1989); species distribution in the mountain range was later substantially completed by Hadač, Terray et al. (1991).
- ¹⁹⁸ In some of the included literature (e.g. Beldie 1977, Oprea 2005, Ciocârlan 2009) *Viola declinata* is evaluated as a Carpathian-Balkan species. According to Velez & Apostolova (2009), it does not occur in Serbia nor in Bulgaria (cf. Diklić 1972, Delipavlov 1979).

¹⁹⁹ In Ukraine, *Viola jooi* grows also in the Pokuttja region (Čornej 2011).

References

- Álvarez Fernández I. (2003): Systematics of Eurasian and Nord African *Doronicum* (Asteraceae: Senecioneae). – Ann. Missouri Bot. Gard. 90: 319–389.
- Anastasiu P. & Negrean G. (2005): Alien plants in Romania. – Analele Ști. Univ. Al. I. Cuza Iași, Biol. Veget. 51: 87–96.
- Ančev M. (2007): Catalogue of the family Brassicaceae (Cruciferae) in the flora of Bulgaria. – Phytol. Balc. 13: 153–178.
- Ančev M. & Polatschek A. (2006): The genus *Erysimum* (Brassicaceae) in Bulgaria. – Ann. Naturhist. Mus. Wien 107 B: 227–273.
- Antosyak T. & Kozurak A. (2011): Distribution of endemic and rare vascular plant species in the upper Tysa Basin within the Carpathian biosphere reserve (Ukraine). – Transylv. Rev. Syst. Ecol. Res. 11: 85–94.
- Ascherson P. & Graebner P. (1898–1901): Synopsis der mitteleuropäischen Flora 2/1. – Gebrüder Borntraeger Verlag, Leipzig.
- Bahlej O. V. (2010): Kompleksna ocinka stanu populjacji *Saussurea porcii* Degen (Asteraceae) v Ukraïns'kykh Karpatach [Integrated assessment of state of populations of *Saussurea porcii* Degen (Asteraceae) in the Ukrainian Carpathians]. – In: Roslynnij svit u Červonyj knyzi Ukraïny: vprovadžennja Hlobal'noï stratehii zberežennja roslyn. Materialy mižnarodnoï konferencii (11–15 žovtnja 2010 r., Kyïv), p. 55–57. Alterpress, Kyïv.
- Baksay L. (1972): Biosystematik der *Dianthus plumarius* L. (sensu lato) in Ungarn. – Symp. Biol. Hung. 12: 149–161.
- Ball P. W. (1993): *Hesperis* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 336–337. Cambridge University Press, Cambridge.
- Bartha L. (2012): Molecular phylogenetic studies on the European species of the genus *Astragalus* L. section *Dissitiflori*. – Summary of the PhD thesis, Babeș-Bolyai University, Cluj-Napoca.
- Bartha L. & Bartók A. (2013): Rediscovery of *Astragalus roemeri* Simonk. in the Scărița-Belioara Natural Reserve (Apuseni Mountains, Romania). – Contr. Bot. Cluj 48: 23–26.
- Beldie A. (1967b): Flora și vegetația Munților Bucegi [Flora and vegetation of the Bucegi Mts]. – Edit. Acad. Rep. Soc. România, București.
- Beldie A. (1972): *Festuca* L. – In: Nyárády E. I. et al. (eds), Flora Republicii Socialiste România 12, p. 459–559. Edit. Acad. Rep. Soc. România, București.
- Beldie A. (1977): Flora României. Determinator ilustrat al plantelor vasculare I [Flora of Romania. Illustrated identification key of vascular plants I]. – Edit. Acad. Rep. Soc. România, București.
- Beldie A. (1979): Flora României. Determinator ilustrat al plantelor vasculare II [Flora of Romania. Illustrated identification key of vascular plants II]. – Edit. Acad. Rep. Soc. România, București.
- Beldie A. & Alexandrescu L. (1976): Modificări de taxonomie sau nomenclatură și sinonimizări la principalii taxonii din vol. I–XII Flora R. S. R. [Changes in taxonomy or nomenclature and synonymy of the main taxa in volumes I–XII of the Flora R. S. R.]. – In: Beldie A. & Morariu I. (eds), Flora Republicii Socialiste România 13, p. 72–96. Edit. Acad. Rep. Soc. România, București.
- Beldie A. & Váczy C. (1976): Taxoni noi pentru Flora României, publicați după editarea volumelor respective din „Flora R. S. R.” [New taxa of the flora of Romania published after „Flora R. S. R.” volumes editing]. – In: Beldie A. & Morariu I. (eds), Flora Republicii Socialiste România 13, p. 35–53. Edit. Acad. Rep. Soc. România, București.
- Bernátová D. (2002): *Papaver tatricum* (A. Nyár.) Ehrend. – In: Goliašová K. & Šípošová H. (eds), Flóra Slovenska V/4, p. 47–54. Veda, Bratislava.
- Bernátová D. & Májovský J. (1997): The *Poa glauca* group in the Western Carpathians I.: *Poa margilicola* sp. nova. – Biologia 52: 27–31.
- Bernátová D. & Májovský J. (1999): *Sorbus pekarovae* Májovský et Bernátová. – In: Čerovský J. et al. (eds), Červená kniha ohrozených a vzácných druhov rastlín a živočíchov SR a ČR 5, p. 350. Vyššie rastliny. Príroda, Bratislava.
- Bernátová D. & Májovský J. (2003): New endemic hybridogenous species of the genus *Sorbus* in the Western Carpathians. – Biologia 58: 781–790.
- Bernátová D., Uhlířová J. & Topercer J. (1998): Aktuálne poznatky o subalpínskej vegetácii Krivánskej Fatry a návrhy na jej manažment [Some new information on subalpine vegetation of the Krivánska Fatra Mts. and management recommendations]. – In: Korňan M. (ed.), Výskum a ochrana Krivánskej Fatry, p. 49–51. Správa Národného parku Malá Fatra, Varín.
- Bernátová D., Májovský J. & Obuch J. (1999): The *Poa glauca* group in the Western Carpathians II.: *Poa sejuncta* sp. nova and *Poa babiogorensis* sp. nova. – Biologia 54: 37–41.
- Bernátová D., Kliment J. & Uhlířová J. (2003a): K cenológii lipnice osobitej (*Poa sejuncta*) a lipnice babiohorskej (*Poa babiogorensis*) [On the coenology of *Poa sejuncta* and *Poa babiogorensis*]. – Bull. Slov. Bot. Spoločen. 25: 231–237.
- Bernátová D., Jarolímek I., Kliment J. & Uhlířová J. (2003b): The association *Poa margilicolae-Primuletum hungaricae* ass. nova in the Veľká Fatra Mts (Central Slovakia). – Biologia 58: 817–822.

- Bernátová D., Májovský J., Kliment J. & Topercer J. (2006): Taxonomy and distribution of *Poa carpatica* in the Western Carpathians. – *Biologia* 61: 387–392.
- Bilz M. (2013): *Syringa josikaea*. – The IUCN Red List of Threatened Species. Version 2014.3. URL: <http://www.iucnredlist.org/details/162267/0> (accessed 16 December 2014).
- Blanár D. & Letz D. R. (2005): *Sempervivum marmoreum* agg. na Muránskej planine [*Sempervivum marmoreum* agg. in the Muránska planina Mts]. – *Reussia* 2: 129–151.
- Bogosavljević S., Zlatković B. & Randelović V. (2007): Flora klisure Svrliškog Timoka [Flora of the Svrlijski Timok Gorge]. – In: Proceeding of the 9th Symposium on flora of Southeastern Serbia and neighbouring regions. Niš September 01–03, 2007, p. 41–54. Niš.
- Böhm É. I. & Facsar G. (2000): A Kitaibel-varfű [*Knautia kitaibelii* (Schultes) Borbás] hazai elő fordulásáról [*Knautia kitaibelii* (Schultes) Borbás in Hungary]. – *Kitaibelia* 5: 347–349.
- Borhidi A. (1957): *Centaurea achtarovii* Urumoff dans les Carpathes orientaux. Annotations sur le groupe *Centaurea triumfettii* All. – *Ann. Hist.-Nat. Mus. Natl. Hung.*, S. N. 8: 219–225.
- Borhidi A., Kevey B. & Lendvai G. (2012): Plant communities of Hungary. – Akadémiai kiadó, Budapest.
- Boroń P., Zalewska-Gałosz J., Sutkowska A., Zemanek B. & Mitka J. (2011): ISSR analysis point to relict character of *Aconitum bucovinense* Zapal. (Ranunculaceae) at the range margin. – *Acta Soc. Bot. Polon.* 80: 315–326.
- Borza A. (1964): Rezervația cu *Leontopodium alpinum* de la Întregalde [The natural reserve with *Leontopodium alpinum* from Întregalde]. – *Ocrot. Nat.* 8: 99–103.
- Boșcaiu M. & Ehrendorfer F. (1996): *Cerastium villosum* Baumg. in der Flora der Karpaten und Ostalpen. – *Stapfia* 45: 31–35.
- Boșcaiu N. (1965): Cercetări fitocenologice asupra asociației *Peucedano (rocheliani)-Molinietum coeruleae* din Banat și Țara Hațegului [Phytocoenological researches on the association of *Peucedano (rocheliani)-Molinietum coeruleae* from Banat and the Hațeg Region]. – *Contr. Bot. Cluj* 1965: 251–264.
- Boșcaiu N. & Rațiu F. (1965): Observații sistematice și anatomice asupra unor specii de *Peucedanum* din secția *Peucedanum* [Observations upon the systematics and anatomy of certain species of *Peucedanum* section *Peucedanum*]. – *Contr. Bot. Cluj* 1965: 299–312.
- Burillo S., Guglielmo A., Pavone P., Scelsi F. & Terassi M. C. (1996): Cytotaxonomic consideration of *Allium fuscum* Waldst. et Kit. (Liliaceae), a critical species the European flora. – *Folia Geobot. Phytotax.* 31: 465–472.
- Budžáková M., Hodálová I., Mered'a P. jr., Somlyay L., Bisbing S. M. & Šibík J. (2014): Karyological, morphological and ecological differentiation of *Sesleria caerulea* and *S. tatrae* in the Western Carpathians and adjacent regions. – *Preslia* 86: 245–277.
- Buia A. & Morariu I. (1972): *Trisetum* Pers. – In: Nyárády E. I. et al. (eds), *Flora Republicii Socialiste România* 12, p. 287–295. Edit. Acad. Rep. Soc. România, București.
- Čalić D., Devrnja N., Milojević J., Zdravković-Korać S., Tubić L., Djuričković M. S. & Vinterhalter B. (2012): Pollen morphology and variability of *Tulipa hungarica* Borb. – *African J. Biotechnol.* 11: 616–620.
- Catană R., Mitoi M. & Ion R. (2013): The RAPD techniques used to assess the genetic diversity in *Draba dorneri*, a critically endangered plant species. – *Advances Biosci. & Biotechnol.* 2013, 4: 164–169.
- Chater A. O. (1968): *Astragalus* L. – In: Tutin T. G. et al. (eds), *Flora Europaea* 2, p. 108–124. Cambridge University Press, Cambridge.
- Chater A. O., Walters S. M., Akeyrod J. K. & Wrigley F. (1993): *Silene* L. – In: Tutin T. G. et al. (eds), *Flora Europaea* 1. Ed. 2, p. 191–218. Cambridge University Press, Cambridge.
- Chorney I., Budzhak V. & Tokaryuk A. (2008): Rare, endangered and endemic species of plants of the Chyvchyn / Civcin Mountains (Carpathians). – *Transylv. Rev. Syst. Ecol. Res.* 5: 37–44.
- Chrtek J. & Pouzar Z. (1985): Further comments on the problem of *Antennaria carpatica*. – *Preslia* 57: 193–198.
- Chrtek J. jr. (1996): Chromosome numbers in selected species of *Hieracium* (Compositae) in the Sudeten Mts. and the Western and Ukrainian Eastern Carpathians. – *Fragm. Florist. Geobot.* 41: 783–790.
- Chrtek J. jr. (1997): *Hieracium decipiensiforme* (the *H. nigrescens* group) – an interesting species of the Ukrainian Carpathians. – *Preslia* 69: 121–128.
- Chrtek J. jr. (2004): *Hieracium* L. – In: Štěpánek J. & Štěpánková J. (eds), *Květena České republiky* 7, p. 540–701. Academia, Praha.
- Chrtek J. jr. & Marhold K. (1998): Taxonomy of the *Hieracium fritzei* group (Asteraceae) in the Sudeten Mts and the West Carpathians (Studies in *Hieracium* sect. *Alpina* II). – *Phyton (Horn)* 37: 181–217.
- Chrtek J. jr. & Mráz P. (2007): Taxonomic revision of *Hieracium nigrescens* agg. in the Western Carpathians. – *Preslia* 79: 45–62.
- Chrtek J. jr., Szeląg Z., Mráz P. & Severa M. (2002): *Hieracium silesiacum* Krause [*Hieracium sparsum* subsp. *silesiacum* (Krause) Zahn] v Západních Karpatech [*Hieracium silesiacum* Krause [*Hieracium sparsum* subsp. *silesiacum* (Krause) Zahn] in the Western Carpathians]. – *Bull. Slov. Bot. Spoločn.* 24: 81–90.

- Chrtěk J. jr., Mráz P. & Severa M. (2004a): Chromosome numbers in selected species of *Hieracium* s. str. (*Hieracium* subgen. *Hieracium*) in the Western Carpathians. – *Preslia* 76: 119–139.
- Chrtěk J. jr., Tonková M., Plačková I., Mráz P. & Marhold K. (2004b): *Hieracium nigrescens* group in the Sudeten Mountains and the Western Carpathians. – *Thaiszia-J. Bot.* 14, Suppl. 1: 3–4.
- Chrtěk J. jr., Tonková M., Mráz P., Marhold K., Plačková I., Krahulcová A. & Kirschner J. (2007): Morphological and allozyme diversity in the *Hieracium nigrescens* group (Compositae) in the Sudety Mountains and Western Carpathians. – *Bot. J. Linn. Soc.* 153: 287–300.
- Cieślak E., Schönswetter P. & Ronikier M. (2013): Phylogenetic analysis confirms the status of *Saxifraga wahlenbergii* Ball (Saxifragaceae) as a distinct endemic of the Western Carpathians. – *Acta Biol. Cracov., Bot.* 55, Suppl. 1: 40.
- Ciocîrlan V. (1988): Flora ilustrată a României 1 [Illustrated flora of Romania 1]. – Editura Ceres, Bucureşti.
- Ciocîrlan V. (1990): Flora ilustrată a României 2 [Illustrated flora of Romania 2]. – Editura Ceres, Bucureşti.
- Ciocîrlan V. (2006): Genul *Aubrieta* Adanson în flora României [Genus *Aubrieta* Adanson in the flora of Romania]. – In: Sesiunea ştiinţifică Conservarea diversităţii plantelor “in situ” şi “ex situ”, Iaşi 25–28 mai 2006, p. 36–37.
- Ciocîrlan V. (2009): Flora ilustrată a României. Pteridophyta et Spermatophyta [Illustrated flora of Romania. Pteridophyta et Spermatophyta]. – Editura Ceres, Bucureşti.
- Ciocîrlan V. (2011): The variability of *Cephalaria uralensis* (Murray) Roem. et Schult. – *J. Pl. Developm.* 18: 105–108.
- Clapham A. R. & Akeyrod J. R. (1993): *Thlaspi* L. – In: Tutin T. G. et al. (eds), *Flora Europaea* 1. Ed. 2, p. 384–388. Cambridge University Press, Cambridge.
- Coldea Gh. (ed.) (1997): Les associations végétales de Roumanie 1. Les associations herbacées naturelles. – Presses Universitaires de Cluj, Cluj.
- Cristea V. (2014): Phytodiversity hotspots for the city of Cluj-Napoca. – *Contr. Bot. Cluj* 49: 191–200.
- Cvelev N. N. (1972): Rod ovsjanica (*Festuca* L.) v SSSR [Genus *Festuca* L. in USSR]. – *Novosti Sist. Vyssh. Rast.* 9: 15–47.
- Cvelev N. N. (1974): Poaceae Barnh. (Graminae Juss. nom. altern.). – In: Fedorov A. A. (ed.), *Flora evropejskoj časti SSSR* 1, p. 117–368. Nauka, Leningrad.
- Cvelev N. N. (1976): Zlaki SSSR [Gramineae of USSR]. – Nauka, Leningrad.
- Cvelev N. N. (1978): Gentianaceae Juss. – In: Fedorov A. A. (ed.), *Flora evropejskoj časti SSSR* 3, p. 57–86. Nauka, Leningrad.
- Cvelev N. N. (2001): *Ranunculus* L. – In: Cvelev N. N. (ed.), *Flora Vostočnoj Evropy* 10, p. 100–158. Tovariščestvo naučnyh izdanij KMK, Moskva & Sankt-Peterburg.
- Cvelev N. N. (2004a): *Silenanthe* (Fenzl) Griseb. et Schenk. – In: Cvelev N. N. (ed.), *Flora Vostočnoj Evropy* 11, p. 225–226. Tovariščestvo naučnyh izdanij KMK, Moskva & Sankt-Peterburg.
- Čopýk V. I. (1976): Vysokohirna flora Ukraïns'kyh Karpat [High-mountain flora of the Ukrainian Carpathians]. – *Naukova dumka*, Kyïv.
- Čornej I. I. (2009): *Nigritella carpatica* (Zapał.) Teppner, Klein et Zagulskij. – In: Diduch J. P. (ed.), Červona knyha Ukraïny. Roslynnyj svit, p. 198. Globalconsalting, Kyïv.
- Čornej I. I. (2011): Krytyčna revizija taksoniv, ščo navodylis' jak endemični dlja flory Ukraïns'kyh Karpat [Critical revision of taxa reported as endemic to flora of the Ukrainian Carpathians]. – *Nauk. Zap. Bukovins'k. Tovar. Pryr.* 1: 23–59.
- Delipavlov D. (1979): Violaceae Batsch. – In: Kuzmanov B. (ed.), *Flora na Narodna republika Bălgaria* 7, p. 338–395. Izdatelstvo na Bălgarskata akademija na naukite, Sofija.
- Derevenko T. (2011): Ex situ conservation of *Saussurea porcii* Degen in Y. Fedkovych National University Botanic Garden. – *J. Pl. Developm.* 18: 151–157.
- Deyl M. (1934): Tři transsilvanské subspecie trav na Podkarp. Rusi [Three transylvanian subspecies of grasses in Subcarpathian Ruthenia]. – *Věda Přir.* 15: 244–245.
- Deyl M. (1980): *Sesleria* Scop. – In: Tutin T. G. et al. (eds), *Flora Europea* 5, p. 173–177. Cambridge University Press, Cambridge.
- Diduch J. P. (ed.) (2007): Ekoflora Ukraïny 5 [Ecoflora of Ukraine 5]. – Fitosociocentr, Kyïv.
- Diduch J. P. (ed.) (2009): Červona knyha Ukraïny. Roslynnyj svit [Red data book of Ukraine. Vegetable kingdom]. – Globalconsalting, Kyïv.
- Diduch J. P., Ziman S. M., Burda P. I. & Čornej I. I. (2004a): *Ranunculus malinovskii* A. Jelen. et Derviz-Sokolova. – In: Diduch J. P. (ed.), *Ekoflora Ukraïny* 2, p. 167–168. Fitosociocentr, Kyïv.
- Dihoru G. & Negrean G. (2009): Cartea roşie a plantelor vasculare din România [Red data book of vascular plants of Romania]. – Edit. Acad. Române, Bucureşti.
- Dihoru G. & Pârnu C. (1987): Plante endemice în flora Românie [Endemic plants in the flora of Romania]. – Editura Ceres, Bucureşti.
- Diklić N. (1972): Violaceae De Cand. – In: Josifović M. (ed.), *Flora SR Srbije* 3, p. 128–164. Srpska Akademija Nauka i Umetnosti, Beograd.
- Diklić N. (1973): Dipsacaceae B. Juss. – In: Josifović M. (ed.), *Flora SR Srbije* 5, p. 536–584. Srpska Akademija Nauka i Umetnosti, Beograd.

- Diklić N. & Nikolić V. (1986a): *Micromeria pulegium* (Rochel) Benth. – In: Sarić R. M. & Diklić N. (eds), Flora SR Srbije 10. Dodatak (2), p. 186. Srpska Akademija Nauka i Umetnosti, Beograd.
- Diklić N. & Vasić O. (2000): A taxonomic assessment of *Thymus dacicus*, a neglected member of the flora of Yugoslavia. – Bot. Chron. 13: 179–186.
- Dmytrach R. I. (2010): Heneratyvne rozmnožennja vysokohirnych vydiv roslyn ta osoblyvosti nasinnevoho vidnovlennja ichnich populacij v Ukraïns'kykh Karpatach [Seed reproduction of the high-mountain plant species and characteristics of seed restoration of populations in the Ukrainian Carpathians]. – Nauk. Visn. NLTU Ukrainy 20.9: 45–51.
- Domin K. (1931a): Schedae ad floram Čechoslovenicam exsiccata. Centuria III. – Acta Bot. Bohem. 10: 3–79.
- Domin K. (1935): Plantarum Čechoslovakiae enumeratio species vasculares indigenas et introductas exhibens. – Preslia 13–15: 3–305.
- Dostál J. (1976): *Centaurea* L. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 254–301. Cambridge University Press, Cambridge.
- Dostál J. (1989): Nová květena ČSSR [New flora of the Czechoslovak Socialist Republic]. – Academia, Praha.
- Dostál J. (1991–1992): Velký klíč na určování vyšších rostlin 1, 2 [The big key for the determination of vascular plants 1, 2]. – SPN, Bratislava.
- Drăgulescu C. (2003): Cormoflora județului Sibiu [Cormophyte flora of Sibiu County]. – Pelecanus, Braşov.
- Dúbravková D., Chytrý M., Willner W., Illyés E., Janišová M. & Kállayné Szerényi J. (2010): Dry grasslands in the Western Carpathians and the northern Pannonian Basin: a numerical classification. – Preslia 82: 165–221.
- Dug S., Muratović E., Drešković N., Boškailo A. & Dudević S. (2013): Crvena lista flore Federacije Bosne i Hercegovine [Red list of the flora of the Federation of Bosnia and Hercegovina]. – NVO „Greenway“ Sarajevo, Sarajevo.
- Dunkel F. G. (2011): Typification of *Ranunculus binatus* Kit. ex Rchb. (Ranunculaceae). – Willdenowia 41: 239–243.
- Dvořák F. (1963): *Hesperis dinarica* G. Beck subsp. *slovaca* Dvořák subsp. nova. – Preslia 35: 243–244.
- Dvořák F. (1968): Rod *Hesperis* v Československu. Taxonomická studie [The genus *Hesperis* in Czechoslovakia. Taxonomic study]. – Biol. Práce 14/3: 1–56.
- Dvořáková M. (1999): *Euphrasia corcontica*, eine endemische Art aus dem Gebirge Krkonoše (Riesengebirge, Westsudeteten). – Preslia 71: 33–35.
- Dvořáková M. (2003): *Minuartia pauciflora*, das karpatische Endemit aus der *M. verna*-Gruppe. – Preslia 75: 349–356.
- Ehrendorfer F. & Krendl F. (1976): *Galium*. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 14–36. Cambridge University Press, Cambridge.
- Eliáš P. jr., Dítě D., Kliment J., Hrivnák R. & Feráková V. (2015): Red list of ferns and flowering plants of Slovakia, 5th edition (October 2014). – Biologia 70: 218–228 & Electronic Appendix.
- Erdelská O. & Turis P. (eds) (1996): Lykovec muránsky (*Daphne arbuscula* Čelak.). – Správa CHKO Muránska planina, Rožňava.
- Fabiszewski J. (1970): Wstępna charakterystyka geobotaniczna otoczenia Jaskini Niedźwiedziej w masywie Śnieżnika [Geobotanical characterization of Jaskinia Niedźwiedzia cave environment in Śnieżnik Mts]. – Acta Univ. Wratislaviensis 127, Stud. Geogr. 14: 85–117.
- Fedorončuk M. M. & Čornej I. I. (2005): Rid *Dianthus* L. (Caryophyllaceae Juss.) flory Ukraïny: taksonomičnyj i sozologičnyj analiz [Genus *Dianthus* L. (Caryophyllaceae Juss.) in the flora of Ukraine – taxonomic and zoological analysis]. – Zapovidna sprava v Ukraïni 11, 2: 9–18.
- Fedorončuk M. M. & Diduch J. P. (2002b): *Dianthus spiculifolius* Schur. – In: Diduch J. P. (ed.), Ekoflora Ukraïny 3, p. 446–447. Fitosociocentr, Kyïv.
- Fedorončuk M. M. & Diduch J. P. (2002c): *Minuartia oxypetala* (Wol.) Kulcz. – In: Diduch J. P. (ed.), Ekoflora Ukraïny 3, p. 154–155. Fitosociocentr, Kyïv.
- Fedorončuk M. M. & Diduch J. P. (2002d): *Elisanthe zawadzkyi* (Herbich) Klokov. – In: Diduch J. P. (ed.), Ekoflora Ukraïny 3, p. 324–325. Fitosociocentr, Kyïv.
- Fernandes R. (1976): *Anthemis* L. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 145–159. Cambridge University Press, Cambridge.
- Ferreira M. Z., Zahradníček J., Kadlecová J., de Sequeira M. M., Chrtek J. jr. & Fehrer J. (2015): Tracing the evolutionary history of the little-known Mediterranean-Macaronesian genus *Andryala* (Asteraceae) by multigene sequencing. – Taxon 64: 535–551 & Electronic Supplement.
- Franco A. (1976): *Carduus* L. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 220–232. Cambridge University Press, Cambridge.
- Fröhner S. (1968): Floristische Neuheiten aus der Hohen und Belauer Tatra (1965). – Preslia 40: 417–426.
- Fröhner S. (1975): Kritik an der europäischen *Alchemilla*-Taxonomie. – Feddes Repert. 86: 119–169.
- Fröhner S. (1986): Zur infragenerischen Gliederung der Gattung *Alchemilla* L. in Eurasien. – Gleditschia 14: 3–49.
- Gejdeman T. S. (1986): Opredelitel' vyššich rastenij Moldavskoj SSR [Identification key of vascular plants of Moldavian Soviet Socialist Republic]. Ed. 3. – Știința, Kișinev.
- Ghișa E. (1972): *Koeleria* Pers. – In: Nyárády E. I. et al. (eds), Flora Republicii Socialiste România 12, p. 227–238. Edit. Acad. Rep. Soc. România, București.
- Ghișa E. & Beldie A. (1972): *Poa* L. – In: Nyárády E. I. et al. (eds), Flora Republicii Socialiste România 12, p. 364–429. Edit. Acad. Rep. Soc. România, București.

- Goliašová K. (1985): Variabilita *Pulsatilla slavica*, *P. grandis*, *P. subslavica* a problém introgresívnej hybridizácie [Die Variabilität *Pulsatilla slavica*, *P. grandis*, *P. subslavica* und das Problem der introgressiven Hybridisation]. – Biol. Práce 31/5: 91–172.
- Goliašová K., Michalková E. & Mráz P. (2008): *Campanula* L. – In: Goliašová K. & Šípošová H. (eds), Flóra Slovenska VI/1, p. 239–349. Veda, Bratislava.
- Greuter W. (2003): The Euro+Med treatment of Cichorieae (Compositae) – generic concepts and required new names. – Willdenowia 33: 229–238.
- Griñtescu G. (1953): *Aconitum* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 3, p. 460–511. Edit. Acad. Rep. Pop. Romîne, București.
- Griñtescu G. & Nyárády E. I. (1960a): *Onosma* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 7, p. 214–226. Edit. Acad. Rep. Pop. Romîne, București.
- Griñtescu G., Gușuleac M. & Nyárády E. I. (1955): *Viola* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 3, p. 553–625. Edit. Acad. Rep. Pop. Romîne, București.
- Grulich V. (1984): Notes on the Czechoslovak species of the genus *Hylotelephium*. – Preslia 56: 47–53.
- Gușuleac M. (1961): Labiatae B. Juss. – In: Nyárády E. I. (ed.), Flora Republicii Populare Romîne 8, p. 82–394. Edit. Acad. Rep. Pop. Romîne, București.
- Gynda L. (2004): Variation of two endemic taxa of *Galium* L. (Rubiaceae) from the Eastern Carpathians. – Visnyk Lviv Univ., Ser. Biol. 36: 96–100.
- Hadač J., Terray J. et al. (1991): Kvetena Bukovských vrchov [Flora of the Bukovské vrchy Mts]. – Příroda, Bratislava.
- Hayek A. (1924): Prodromus Florae peninsulae Balcanicae 1(2). – Feddes Repert., Beih. 30: 161–352.
- Hegedúšová K., Škodová I., Janišová M. & Kochjarová J. (2013): Phytosociological affiliation of Annex II species *Tephoseris longifolia* subsp. *moravica* in comparison with two related species *T. integrifolia* and *T. crispa*. – Biologia 68: 861–871.
- Heltmann H. (1985): Zum Endemismus des Südost-Karpatenraumes. – In: Heltmann H. & Wendelberger G. (eds), Naturwissenschaftliche Forschungen über Siebenbürgen. III. Beiträge zur Pflanzengeographie des Südost-Karpatenraumes. Siebenbürg. Arch., III. Folge, 20, p. 157–216.
- Hel'tman D. V. (1989): *Leontodon* L. – In: Cvelev N. N. (ed.), Flora evropejskoj časti SSSR 8, p. 27–32. Nauka, Leningrad.
- Hendrych R. (1987): Karpatische Migrationen und Florenbeziehungen in den tschechischen Ländern der Tschechoslowakei. – Acta Univ. Carol., Biol. 1985: 105–250.
- Hendrych R. (1993): *Trifolium sarosiense* in der Slowakei. – Preslia 65: 33–52.
- Hendrych R. (1995): *Trifolium sarosiense* in Ungarn, Rumänien und Jugoslawien. – Preslia 66 (1994): 305–321.
- Hodálová I. (1999a): Multivariate analysis of the *Senecio nemorensis* group (Compositae) in the Carpathians with a new species from the East Carpathians. – Folia Geobot. 34: 321–335.
- Hodálová I. (1999b): Taxonomy of the *Senecio nemorensis* group (Compositae) in the Carpathians. – Biologia 54: 395–404.
- Hodálová I. & Marhold K. (1998): Morphometric comparison of *Senecio germanicus* and *Senecio nemorensis* (Compositae) with a new species from Romania. – Bot. J. Linn. Soc. 128: 277–290.
- Hodálová I. & Mártonfi P. (2002): *Thlaspi* L. – In: Goliašová K. & Šípošová H. (eds), Flóra Slovenska V/4, p. 600–618. Veda, Bratislava.
- Holobiuc M., Blindu R., Mitoi M., Helepciuc F. & Cristea V. (2009): The establishment of an in vitro gene bank in *Dianthus spiculifolius* Schur and *D. glacialis* ssp. *gelidus* (Schott, Nym. et Kotschy) Tutin. I. The initiation of a tissue collection and the characterization of the cultures in minimal growth conditions. – Ann. Forest. Res. 52: 117–128.
- Holub J. (1977b): New Names in Phanerogamae 5. – Folia Geobot. Phytotax. 12: 293–311.
- Holub J. (1984): Subspecies names in Jávorka, Magyar Flóra (1924–1925). – Preslia 56: 303–318.
- Hurdu B. I., Pușcaș M., Turtureanu P. D., Niketić M., Coldea G. & Zimmermann N. E. (2012a): Patterns of plant endemism in the Romanian Carpathians (South-Eastern Carpathians). – Contr. Bot. Cluj 47: 25–38.
- Hurdu B. I., Pușcaș M., Turtureanu P. D., Niketić M., Vonica G., Coldea G. (2012b): A critical evaluation of the Carpathian endemic plant taxa list from the Romanian Carpathians. – Contr. Bot. Cluj 47: 39–47.
- Il'jins'ka A. P., Diduch J. P. & Korotchenko I. A. (2007): *Erysimum transsilvanicum* Schur. – In: Diduch J. P. (ed.), Ekoflora Ukraïny 5, p. 228–229. Fitosociocentr, Kyïv.
- Ilnicki T. & Mitka J. (2009): Chromosome numbers in *Aconitum* sect. *Aconitum* (Ranunculaceae) from the Carpathians. – Caryologia 62: 198–203.
- Ilnicki T. & Mitka J. (2011): Chromosome numbers in *Aconitum* sect. *Cammarum* (Ranunculaceae) from the Carpathians. – Caryologia 64: 446–452.
- Ilnicki T. & Szeląg Z. (2011): Chromosome numbers in *Hieracium* and *Pilosella* (Asteraceae) from Central and Southeastern Europe. – Acta Biol. Cracov., Bot. 53: 102–110.
- Ion R. (2012): First record of *Draba dorneri* Heuff. (Brassicaceae) in Parâng Mountains (Romanian Carpathians). – The 52th Annual Scientific Conference „A New Approach of the Academic Research in Biology“. Institute of Biology Bucharest of the Romanian Academy (poster).
- Jakab G., Csörgő A.-M. & Ambrus L. (2008): Kiegészítések Székelyföld flórájának ismeretéhez [Contributions to the flora of Szeklerland (Romania)]. – Kitaibelia 13: 107.

- Jalas J. & Suominen J. (eds) (1983): Atlas Florae Europaeae 6. Caryophyllaceae (Alsinoideae and Paronychioideae). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J. & Suominen J. (eds) (1986): Atlas Florae Europaeae 7. Caryophyllaceae (Silenoideae). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J. & Suominen J. (eds) (1989): Atlas Florae Europaeae 8. Nymphaeaceae to Ranunculaceae. – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J. & Suominen J. (eds) (1994): Atlas Florae Europaeae 10. Cruciferae (*Sisymbrium* to *Aubrieta*). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J., Suominen J. & Lampinen R. (eds) (1996): Atlas Florae Europaeae 11. Cruciferae (*Ricotia* to *Raphanus*). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J., Jackson M. B. W., Sell P. D. & Whitehead F. H. (1993): *Cerastium* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 164–175. Cambridge University Press, Cambridge.
- Janchen E. (1956–1960): Catalogus Florae Austriae. 1. Pteridophyten und Anthophyten (Farne und Blütenpflanzen). – Springer Verlag, Wien.
- Janišová M., Hegedúšová K., Král P. & Škodová I. (2012): Ecology and distribution of *Tephrosia longifolia* subsp. *moravica* in relation to environmental variation at a micro-scale. – Biologia 67: 97–109.
- Jávorka S. (1906): Hazai *Onosma*-fajaink [Species Hungaricae generis *Onosma*]. – Ann. Hist.-Nat. Mus. Natl. Hung. 4: 406–449.
- Jirásek V. (1934): *Poa sterilis* M. Bieb. a *Poa capillifolia* Kalchbr. v Československu [*Poa sterilis* M. Bieb. and *Poa capillifolia* Kalchbr. in Czechoslovakia]. – Věda Přir. 15: 205–206.
- Jirásek V. (1935): The distribution of the species of the genus *Poa* in Czechoslovakia. – Bull. Int. Acad. Sci. Bohem. 1934: 1–9.
- Joachimiak A., Ilnicki T. & Mitka J. (1999): Karyological studies on *Aconitum lasiocarpum* (Rchb.) Gayer (Ranunculaceae). – Acta Biol. Cracov., Bot. 41: 205–211.
- Kahalo O. O. & Syčák A. N. (2009): *Armeria pocutica* Pawl. – In: Diduch J. P. (ed.), Červona knyha Ukraïny. Roslynnij svit, p. 517. Globalconsalting, Kyïv.
- Kanka R., Turis P. & Chilová V. (2008): Phytosociological characteristic of the plant communities with the occurrence of endemic species *Cyclamen fatrense*. – Hacquetia 7: 21–32.
- Kárpáti Z. (1960): Die *Sorbus*-Arten Ungarns und der angrenzender Gebiete. – Feddes Repert. 62: 71–331.
- Katina Z. F. (1961): Valerianaceae DC. – In: Kotov M. I. (ed.), Flora URSS 10, p. 291–339. Vydavnyctvo Akademii nauk Ukraïns'koï RSR, Kyïv.
- Kerényi-Nagy V. (2011): A szentendrei rózsza (*Rosa sancti-andreae*) nevezéktani és taxonómiai problémái [Nomenclatural and taxonomical problems related to the *Rosa sancti-andreae*]. – Kanitzia 18: 13–28.
- Kereszty Z. (1993): The distribution of the genus *Scilla* in Hungary. – Stud. Bot. Hung. 24: 51–75.
- Kereszty Z., Szilágyi L. & Borhidi A. (1986): Biosystematic studies of the *Scilla bifolia* complex in Hungary. – Uppsala Univ. Arsskr. 27: 107–112.
- Kirschner J. (2007): Valerianaceae Batsch. – Ms., Institute of Botany, Slovak Academy of Sciences, Bratislava.
- Kirschner J. & Štěpánek J. (1985): „*Taraxacum hoppeanum*“ and its allies (Studies in *Taraxacum* 4). – Preslia 37: 111–134.
- Kliment J. (1999): Komentovaný prehľad vyšších rastlín flóry Slovenska, uvádzaných v literatúre ako endemické taxóny [Annotated survey of the vascular plants of Slovak flora recorded in literature as endemic taxa]. – Bull. Slov. Bot. Spoločn. 21, Suppl. 4: 1–434.
- Kliment J. (2003): Zamyslenie sa nad súčasným fyto geografickým členením Slovenska (poznámky k vybraným fytochoriónom) [Thoughts on recent phytogeographical regionalisation of Slovakia (notes to selected phytochorions)]. – Bull. Slov. Bot. Spoločn. 25: 199–224.
- Kliment J., Bernátová D., Dítě D., Janišová M., Jarolímek I., Kochjarová J., Kučera P., Obuch J., Topercer J., Uhlířová J. & Zaliberová M. (2008): Paprad'orasty a semenné rastliny [Ferns and flowering plants]. – In: Kliment J. (ed.), Příroda Vel'kej Fatry. Lišajníky, machorasty, cievnaté rastliny, p. 109–367. Vydavateľstvo Univerzity Komenského, Bratislava.
- Klokov M. V. (1952): Caryophyllaceae Juss. – In: Kotov M. I. (ed.), Flora URSS 4, p. 421–646. Vydavnyctvo Akademii nauk Ukraïns'koï RSR, Kyïv.
- Klokov M. V. (1960): Labiatae Juss. – In: Kotov M. I. (ed.), Flora URSS 9, p. 5–364. Vydavnyctvo Akademii nauk Ukraïns'koï RSR, Kyïv.
- Klokov M. V. (1961): Rubiaceae Juss. – In: Kotov M. I. (ed.), Flora URSS 10, p. 90–249, 455–474. Vydavnyctvo Akademii nauk Ukraïns'koï RSR, Kyïv.
- Kmeťová E. (1985): Taxonómia druhu *Dianthus praecox* Kit. na Slovensku [Taxonomie der Art *Dianthus praecox* Kit. in der Slowakei]. – Biol. Práce 31/5: 3–85.
- Kmeťová E. (2012): *Dianthus* L. – In: Goliašová K. & Michalková E. (eds), Flóra Slovenska VI/3, p. 589–655. Veda, Bratislava.
- Kobiv J. (2010): Ekologické osoblivosti oselyšč rídkisnych vydív roslyn Ukraïns'kich Karpat [Ecological characteristics of habitats of rare plant species in the Ukrainian Carpathians]. – Ukraïns'k. Bot. Zhurn. 67: 355–372.

- Kobiv J. (2012a): Roľ prýdatných mikrooselyš u samovidovlenni populácií rідkisnich vydiv roslyn Ukraїns'kych Karpat [Role of suitable microhabitats in population recruitment of rare plant species in the Ukrainian Carpathians]. – Ukraїns'k. Bot. Zhurn. 69: 178–189.
- Kobiv J. (2012b): Does *Aquilegia transsilvanica* Schur (Ranunculaceae) occur in the Ukrainian Carpathians? – Ukraїns'k. Bot. Zhurn. 69: 493–501.
- Kobiv J., Prokopiv A., Heleš M., Borsukevyč L. & Nadraha M. (2007a): Pošyrennja i stan populácií rідkisnych, zahroženych ta endemičnych vydiv roslyn u pivničnij častyni prikordonnoї diljanky Čyvčyns'kych hir (Ukraїns'ki Karpaty) [Distribution and state of populations of rare, endangered and endemic species of plants in the northern part of frontier area in the Chyvchyny Mountains (Ukrainian Carpathians)]. – Visnyk L'viv. Univ., Ser. Biol. 45: 71–84.
- Kobiv J., Heleš M. & Borsukevich L. (2007b): *Saussurea porcii* Degen (Asteraceae) in the Svydovets Mountains (Ukrainian Carpathians): location, coenotic conditions, population parameters and conservation. – Ukraїns'k. Bot. Zhurn. 64: 825–832.
- Kochjarová J. (1998): Poznámky k rozšíreniu, cenológii a ohrozenosti populácií zástupcov rodu *Tephroseris* (Rchb.) Rchb. na Slovensku. II: *T. longifolia* subsp. *moravica* v Západných Karpatoch [Notes on distribution, coenology and threat of populations of *Tephroseris* (Rchb.) Rchb. in Slovakia. II: *T. longifolia* subsp. *moravica* in the West Carpathians]. – Bull. Slov. Bot. Spoločn. 20: 69–77.
- Kochjarová J. & Hrouda L. (2004): *Tephroseris* (Reichenb.) Reichenb. – In: Slavík B., Štěpánková J. & Štěpánek J. (eds), Květena České republiky 7, p. 300–306. Academia, Praha.
- Kochjarová J. & Hrouda L. (2006): Príspevok k rozlišovaniu a rozšíreniu západokarpatského endemického druhu *Soldanella carpatica* (Primulaceae) [Contribution to distinguishing and distribution of the Western Carpathian endemic species *Soldanella carpatica* (Primulaceae)]. – Bull. Slov. Bot. Spoločn. 28: 127–139.
- Kochjarová J. & Valachovič M. (2002): *Cochlearia* L. – In: Goliašová K. & Šipošová H. (eds), Flóra Slovenska V/4, p. 549–561. Veda, Bratislava.
- Kochjarová J., Vlčko J. & Hrivnák R. 2004. Diploidné populácie *Scilla bifolia* agg. v Západných Karpatoch a príľahlej časti Panónskej nížiny [The diploid populations of *Scilla bifolia* group in the Western Carpathians and adjacent part of the Pannonian lowland]. – Bull. Slov. Bot. Spoločn., Suppl. 10: 171–175.
- Kochjarová J., Hrivnák R. & Vlčko J. (2005): Diploidné populácie *Scilla bifolia* agg. na Slovensku [The diploid populations of *Scilla bifolia* group in Slovakia]. – Bull. Slov. Bot. Spoločn. 27: 53–62.
- Kochjarová J., Valachovič M., Bureš P. & Mráz P. (2006): The genus *Cochlearia* L. in the Eastern Carpathians and adjacent areas. – Bot. J. Linn. Soc. 151: 355–364.
- Kolář F., Štech M., Trávníček P., Rauchová J., Urfus T., Vít P., Kubešová M. & Suda J. (2009): Towards resolving the *Knautia arvensis* agg. (Dipsacaceae) puzzle: primary and secondary contact zones and ploidy segregation at landscape and microgeographic scales. – Ann. Bot. 103: 963–974.
- Kolarčík V., Zozomová-Lihová J. & Mártonfi P. (2010): Systematics and evolutionary history of the *Asterotricha* group of the genus *Onosma* (Boraginaceae) in central and southern Europe inferred from AFLP and nrDNA ITS data. – Pl. Syst. Evol. 290: 21–45.
- Kolarčík V., Zozomová-Lihová J., Ducár E. & Mártonfi P. (2014): Evolutionary significance of hybridization in *Onosma* (Boraginaceae): analyses of stabilized hemisexual odd polyploids and recent sterile hybrids. – Biol. J. Linn. Soc. 112: 89–107.
- Kolník M. & Marhold K. (2006): Distribution, chromosome numbers and nomenclatural conspect of *Arabidopsis halleri* (Brassicaceae) in the Carpathians. – Biologia 61: 41–50.
- Konětopský A. (1963): Nejdůležitější výsledky taxonomické revize československých druhů rodu *Erysimum* L. [Die wichtigsten Ergebnisse taxonomischer Revision der tschechoslowakischen Arten der Gattung *Erysimum* L.]. – Preslia 35: 135–145.
- Korzeniak U. (2008): *Erysimum pieninicum* (Zapał.) Pawł. – In: Mirek Z. & Piękoś-Mirkowa H. (eds), Czerwona księga Karpat Polskich. Rośliny naczyniowe, p. 150–151. Instytut Botaniki im. W. Szafera PAN, Kraków.
- Koutecký P. (2013): A revision of the endemic species *Centaurea carpatica*. – Acta Biol. Cracov., Bot. 55, Suppl. 1: 52.
- Kovács J. A. (2011): Coenological differentiation of *Peucedanum* species (sect. *Peucedanum*) stands in the Carpathian Basin. – Ann. Bot. 2011: 19–28.
- Kovanda M. (1985): Jarabiny Folkmárskej skaly [*Sorbus* species in Mt. Folkmárska skala]. – Pulsatilla 9: 7–9.
- Kovanda M. (1986): *Sorbus scepusiensis*, a new species of *Sorbus* (Rosaceae) from Eastern Slovakia. – Willdenowia 16: 117–119.
- Kožuharov S. & Petrova A. (1988): Atlas florae Europae notes. 12. – Ann. Bot. Fenn. 25: 389–390.
- Králik E. (1997): *Euphrasia* L. – In: Goliašová K. (ed.), Flóra Slovenska V/2, p. 329–375. Veda, Bratislava.
- Kricsfalusy V. (1999): Flora and vegetation of the Ukrainian upper Tisa Basin: Aspects of biodiversity conservation. – In: Hamar J. & Sárkány-Kiss A. (eds), The upper Tisa valley. Preparatory proposal for Ramsar site designation and an ecological background Hungarian, Romanian, Slovakian and Ukrainian co-operation. Tiscia monograph series 4, p. 273–292. Szeged.

- Kricsfalussy V. V. & Budnikov G. B. (2002): Endemic vascular plants in the Ukrainian Carpathians. – In: Hamor F. D. (ed.), Mountains and people (in the context of sustainable development). The conference proceedings dedicated to the International Year of Mountains, Rakhiv, October 14–18, 2002, p. 356–360.
- Kricsfalussy V. & Budnikov G. (2007): Threatened vascular plants in the Ukrainian Carpathians: current status, distribution and conservation. – *Thaiszia-J. Bot.* 17: 11–32.
- Kricsfalussy V. V. & Vajnagi A. V. (1994): Biologie und Ökologie von *Scilla kladnii* Schur (Hyacinthaceae) in den Ostkarpathen. – *Linzer Biol. Beitr.* 26: 1081–1111.
- Kučera J., Turis P., Zozomová-Lihová J. & Slovák M. (2013): *Cyclamen fatrense*, myth or true Western Carpathian endemic? Genetic and morphological evidence. – *Preslia* 85: 133–158.
- Kurtto A., Fröhner S. E. & Lampinen R. (eds) (2007): Atlas Florae Europaeae 14. Rosaceae (*Alchemilla* and *Aphanes*). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Kurtto A., Uotila P. & Sennikov A. (2009): *Alchemilla* in Mediterranean Europe as revealed by Atlas Florae Europaeae. – *Boccone* 23: 221–235.
- Kuzmanović N., Comanescu P., Frajman B., Lazarević M., Paun O., Schönschwetter P. & Lakušić D. (2013): Genetic, cytological and morphological differentiation within the Balkan-Carpathian *Sesleria rigida* sensu Fl. Eur. (Poaceae): A taxonomically intricate tetraploid-octoploid complex. – *Taxon* 62: 458–472.
- Kuz'mina M. L. (2004): *Dianthus* L. – In: Cvelev N. N. (ed.) Flora vostočnoj Evropy 11, p. 273–297. Товарищество научных изданий КМК, Москва & Санкт-Петербург.
- Lendvay B. & Kalapos T. (2014): Population dynamics of the climate-sensitive endangered perennial *Ferula sadleriana* Ledeb. (Apiaceae). – *Pl. Spec. Biol.* 29: 138–151.
- Lendvay B., Pedryc A. & Höhn M. (2013): Genetic characteristics of the Carpathian endemic *Syringa josikaea*: regional structuring of small and isolated populations. – *Acta Biol. Cracov., Bot.* 55, Suppl. 1: 28.
- Letz R. (1998): Vybrané problémy taxonomickej diferenciácie rodov *Sempervivum* a *Jovibarba* v Európe [Selected problems of taxonomic differentiation of the genera *Sempervivum* and *Jovibarba* in Europe]. – PhD thesis, Faculty of Sciences, Comenius University, Bratislava.
- Letz R. (2002): The nomenclature of the *Sempervivum montanum* group (Crassulaceae) in the Carpathians. – *Phyton (Horn)* 42: 109–115.
- Letz D. R. (2009): A new species of the *Sempervivum marmoreum* group in the Central Europe. – *Preslia* 81: 293–308.
- Letz R. & Marhold K. (1998): Multivariate morphometric study of the *Sempervivum montanum* group (Crassulaceae) in the West Carpathians. – *Phyton (Horn)* 38: 323–336.
- Letz D. R. & Michalková E. (2012): *Cerastium* L. – In: Goliašová K. & Michalková E. (eds), Flóra Slovenska VI/3, p. 291–388. Veda, Bratislava.
- Lysák M. A. (1996): Taxonomie a chorologie českých a slovenských druhů rodu *Sesleria* [Taxonomy and chorology of the Czech and Slovak *Sesleria* species]. – Diploma thesis, Palacký University, Olomouc.
- Maciejewska-Rutkowska I., Bednorz L. & Fujiki T. (2007): SEM observations of pollen grains, fruits and seeds of the Pieniny Mts (South Poland) endemic species *Erysimum pieninicum* (Zapał.) Pawł. (Brassicaceae). – *Acta Soc. Bot. Polon.* 76: 127–132.
- Májovský J. (1970): Neue Arten in der Flora der Slowakei. I. – *Acta Fac. Rerum Nat. Univ. Comen., Bot.* 18: 7–16.
- Májovský J. & Bernátová D. (1996): *Sorbus pekarovae* sp. nova – a new hybridogenous species from the Veľká Fatra Mts. – *Biologia* 51: 23–26.
- Májovský J., Murín A. et al. (1987): Karyotaxonomický prehľad flóry Slovenska [Karyotaxonomic survey of the flora of Slovakia]. – Veda, Bratislava.
- Májovský J., Murín A. & Hindáková M. (1991): Karyotaxonomic studies of Slovak populations of the genus *Crocus*. Part I. – *Acta Fac. Rerum Nat. Univ. Comen., Bot.* 38: 49–87.
- Májovský J., Bernátová D., Obuch J. & Topercer J. (1998): *Sorbus margittiana*, an endemic of Krivánska Fatra Mts. – *Biologia* 53: 29–35.
- Malynovs'kyj K., Caryk J., Kyjak V. & Nesteruk J. (2002): Ridkisi, endemični, reliktovi ta pohranično-areál'ni vydy roslyn Ukraïns'kykh Karpat [Rare, endemic, relict and boundary areal plant species of the Ukrainian Carpathians]. – Instytut ekolohii Karpat Nacional'noi akademii nauk Ukraïny, Lviv.
- Mardari C. (2009): Aspects of the floristic diversity in Neagra Broștenilor River Basin (Eastern Carpathians) (II). – *J. Pl. Developm.* 16: 29–38.
- Marhold K. & Kochjarová J. (2002): *Cardamine* L. – In: Goliašová K. & Šipošová H. (eds), Flóra Slovenska V/4, p. 316–382. Veda, Bratislava.
- Marinov Y., Tosheva A. & Pachedjieva K. (2014): Current status of the rare species *Lathyrus transsilvanicus* in Bulgaria. – *Bulgarian J. Agric. Sci.* 20, Suppl. 1: 160–164.
- Markgraf-Dannenbergh I. (1980): *Festuca* L. – In: Tutin T. G. et al. (eds), Flora Europaea 5, p. 125–153. Cambridge University Press, Cambridge.
- Mártonfi P. (1996): *Thymus alternans* Klokov – a new species of Slovak flora. – *Biologia* 51: 27–29.
- Mártonfi P. (1997): Nomenclatural survey of the genus *Thymus* sect. *Serpyllum* from Carpathians and Pannonia. – *Thaiszia-J. Bot.* 7: 111–181.
- Mártonfi P., Kolarčík V. & Somlyay L. (2014): Lectotypification of *Onosma viridis* and synonymization of *O. tornensis* with *O. viridis* (Boraginaceae). – *Ann. Bot. Fenn.* 51: 201–206.
- Mátis A., Szabó A. & Bartha L. (2014): Lost and found: Rediscovery of *Saussurea porcii* Degen in the Rodnei Mountains (Eastern Carpathians, Romania) after more than a century. – *Contr. Bot. Cluj* 49: 39–42.

- Mazúr E. & Lukniš M. (1980): Geomorfologické jednotky [Geomorphological units] 1: 500 000. – In: Mazúr E. (ed.), Atlas Slovenskej socialistickej republiky, p. 54–55, map IV/16. SAV & SÚGK, Bratislava.
- McNeill J., Barrie F. R., Buck W. R., Demoulin V., Greuter W., Hawksworth D. L., Herendeen P. S., Knapp S., Marhold K., Prado J., Prud'homme van Reine W. F., Smith G. F., Wiersema J. H. & Turland N. J. (eds) (2012): International Code of Nomenclature for algae, fungi, and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. – Regnum Veg. 154. Koeltz Scientific Books, Königstein.
- Melzer H. & Polatschek A. (1971): *Erysimum hungaricum* Zapal. – auch in den Ostalpen. – Ann. Naturhist. Mus. Wien 75: 103–109.
- Mered'a P., Eliáš P. jr., Dítě D. & Štrba P. (2012): *Silene* L. – In: Goliašová K. & Michalková E. (eds), Flóra Slovenska VI/3, p. 410–533. Veda, Bratislava.
- Měsíček J. (2002): *Cardaminopsis* (C. A. Mey.) Hayek. – In: Goliašová K. & Šipošová H. (eds), Flóra Slovenska V/4, p. 388–415. Veda, Bratislava.
- Michalik S. & Mitka J. (2008): *Melampyrum saxosum* Baumg. – In: Mirek Z. & Piękoś-Mirkowa H. (eds), Czerwona księga Karpat Polskich. Rośliny naczyniowe, p. 300–301. Instytut Botaniki im. W. Szafera PAN, Kraków.
- Michalková E. (1993): *Galium mollugo* agg. (Rubiaceae) in Slovakia. – Preslia 65: 201–207.
- Michalková E. (1999): *Erysimum pallidiflorum* (Brassicaceae) na Slovensku. – Ochr. Prír. (Banská Bystrica) 17: 39–47.
- Michalková E. (2002): *Erysimum* L. – In: Goliašová K. & Šipošová H. (eds), Flóra Slovenska V/4, 182–226. Veda, Bratislava.
- Mihály A. V. & Komendar V. I. (1993): State of population of *Crocus banaticus* J. Gay in the Transcarpathian region of the Tisza valley. – Tiscia 27: 61–63.
- Mihok C., Antal D. S. & Csédő C. (2005): The repartition of *Aconitum* species throughout the Romanian Carpathians. – Gyepgazdálkodási Közlem. 3: 26–28.
- Mihoková L. & Mikoláš V. (1994): A contribution to the karyology of *Euphrasia tatrae* Wettst. (Scrophulariaceae). – Thaiszia-J. Bot. 4: 77–80.
- Mikoláš V. (1996): *Sorbus dolomiticola* Mikoláš, a new hybridogenous species of the genus *Sorbus* s. l. from eastern Slovakia. – Thaiszia-J. Bot. 6: 1–12.
- Mikoláš V. (2003): *Sorbus amici-petri* Mikoláš, a new hybridogenous species of the genus *Sorbus* s. l. from eastern Slovakia. – Thaiszia-J. Bot. 13: 127–133.
- Mirek Z. & Piękoś-Mirkowa H. (2008b): *Erigeron hungaricus* (Vierh.) Pawł. – In: Mirek Z. & Piękoś-Mirkowa H. (eds), Czerwona księga Karpat Polskich. Rośliny naczyniowe, p. 380–381. Instytut Botaniki im. W. Szafera PAN, Kraków.
- Mirek Z. & Piękoś-Mirkowa H. (2008c): *Poa granitica* Braun-Blanq. – In: Mirek Z. & Piękoś-Mirkowa H. (eds), Czerwona księga Karpat Polskich. Rośliny naczyniowe, p. 544–545. Instytut Botaniki im. W. Szafera PAN, Kraków.
- Mirek Z. & Piękoś-Mirkowa H. (2009): Fitogeograficzne aspekty endemizmu w Polsce [Phytogeographic aspects of endemism in Poland]. – Wiadom. Bot. 53: 7–30.
- Mirek Z. & Piękoś-Mirkowa H. (2010): Fitogeograficzna analiza endemitów roślin naczyniowych Tatrzańskiego Parku Narodowego [Phytogeographical analysis of the endemic vascular plant taxa occurring in the Tatry Mts]. – Nauka a zarządzanie obszarem Tatr i ich otoczeniem 2: 11–15.
- Mirek Z., Piękoś-Mirkowa H., Zajac A. & Zajac M. (1995): Vascular plants of Poland. A checklist. Krytyczna lista roślin naczyniowych Polski. – Polish Bot. Stud., Guidebook Series 15: 7–303.
- Mirek Z., Piękoś-Mirkowa H., Zajac A. & Zajac M. (2002): Flowering plants and pteridophytes of Poland. A checklist. – W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.
- Mitka J. (2001). Systematyka *Aconitum* subgen. *Aconitum* w Karpatach Wschodnich – wstępne wyniki badań [Systematics of *Aconitum* subgen. *Aconitum* in the Eastern Carpathians – preliminary results]. – Roczn. Bieszcz. 9 (2000): 79–116.
- Mitka J. (2002): Phenetic and geographic pattern of *Aconitum* sect. *Napellus* (Ranunculaceae) in the Eastern Carpathians – a numerical approach. – Acta Soc. Bot. Polon. 71: 35–48.
- Mitka J. (2003): The genus *Aconitum* L. (Ranunculaceae) in Poland and adjacent countries. A phenetic-geographic study. – Institute of botany of the Jagiellonian University, Kraków.
- Mitka J. (2008): *Aconitum moldavicum* Hacq. (Ranunculaceae) and its hybrids in the Carpathians and adjacent regions. – Roczn. Bieszcz. 16: 233–252.
- Mitka J. (2012): *Aconitum* in Central Europe: from Linnaean taxonomy to molecular markers. – Modern Phytomorphol. 1: 7–9.
- Mitka J. & Starmühler W. (2000): Phenetic variability of *Aconitum lasiocarpum* (Rchb.) Gayer (Ranunculaceae): extension of taxonomic and geographic borders. – Acta Soc. Bot. Polon. 69: 145–155.
- Mitka J., Sutkowska A., Ilnicki T. & Joachimiak A. J. (2007): Reticulate evolution of high-alpine *Aconitum* (Ranunculaceae) in the Eastern Sudetes and Western Carpathians (Central Europe). – Acta Biol. Cracov., Bot. 49: 15–26.
- Mitka J., Zemanek B., Winnicki T. & Mirek Z. (2008): *Delphinium elatum* L. subsp. *nacladense* (Zapal.) Holub. – In: Mirek Z. & Piękoś-Mirkowa Z. (eds), Czerwona księga Karpat Polskich. Rośliny naczyniowe, p. 84–85. Instytut Botaniki im. W. Szafera PAN, Kraków.

- Mitka J., Boroń P. & Sutkowska A. (2013): Holocene history of *Aconitum* in the Polish Western Carpathians and adjacent regions: long-distance migrations or cryptic refugia? – *Modern Phytomorphol.* 3: 9–18.
- Morariu I. (1961): Valerianaceae Batsch. – In: Nyárády E. I. (ed.), *Flora Republicii Populare Romîne* 8, p. 615–639. Edit. Acad. Rep. Pop. Romîne, București.
- Morariu I. & Beldie A. (1976): Endemisme din Flora R.S.R. [Endemics in the Flora R. S. R.]. – In: Beldie A. & Morariu I. (eds), *Flora Republicii Socialiste România* 13, p. 97–105. Edit. Acad. Rep. Soc. România, București.
- Mosyakin S. L. & Fedoronchuk M. M. (1999): Vascular plants of Ukraine. A nomenclatural checklist. – National Academy of Sciences of Ukraine, M. G. Kholodny Institute of Botany, Kiev.
- Mráz P. (2001a): *Hieracium rohacsense*, endemit Západných Karpát, a poznámky k jeho taxonómii, chorológii a ekológii [*Hieracium rohacsense*, endemic of the West Carpathians, and notes on its taxonomy, chorology and ecology]. – *Preslia* 73: 341–358.
- Mráz P. (2001b): Chromosome numbers in selected species of *Hieracium* sect. *Alpina* (Asteraceae) from Central and Eastern Europe. – *Folia Geobot.* 36: 321–332.
- Mráz P. (2002): Contribution to the knowledge of the *Hieracium rohacsense* group in the Carpathians. – *Thaiszia-J. Bot.* 12: 109–135.
- Mráz P. (2003a): The *Hieracium pietroszense* group in the Carpathians. – *Folia Geobot.* 38: 299–318.
- Mráz P. (2003b): Mentor effects in the genus *Hieracium* s. str. (Compositae, Lactucaee). – *Folia Geobot.* 38: 345–350.
- Mráz P. (ed.) (2005a): Chromosome number and DNA ploidy level reports from Central Europe – 1. – *Biologia* 60: 99–103.
- Mráz P. (2005b): *Hieracium silesiacum* (Asteraceae) in Poland. – *Polish Bot. J.* 50: 65–68.
- Mráz P. (ed.) (2006): Chromosome number and DNA ploidy level reports from Central Europe 2. – *Biologia* 61: 115–120.
- Mráz P. & Paule J. (2006): Experimental hybridization in the genus *Hieracium* s. str.: crosses between diploid taxa. – *Preslia* 78: 1–26.
- Mráz P. & Szeląg Z. (2004): Chromosome numbers and reproductive systems in selected species of *Hieracium* and *Pilosella* (Asteraceae) from Romania. – *Ann. Bot. Fenn.* 41: 405–414.
- Mráz P., Chrtek J. jr., Fehrer J. & Plačková I. (2005): Rare recent natural hybridization in *Hieracium* s. str. – evidence from morphology, allozymes and chloroplast DNA. – *Pl. Syst. Evol.* 255: 177–192.
- Mráz P., Bouchier R. S., Treier U. A., Schaffner U. & Müller-Schärer H. (2011a): Polyploidy in phenotype space and invasions context: a morphometric study of *Centaurea stoebe* s. l. – *Int. J. Plant Sci.* 172: 386–402.
- Mráz P., Garcia-Jacas N., Gex-Fabry E., Susanna A., Barres L. & Müller-Schärer H. (2012): Allopolyploid origin of highly invasive *Centaurea stoebe* s. l. (Asteraceae). – *Molec. Phylogenet. Evol.* 62: 612–623.
- Mucher W. (1993): Systematics and chorology of *Aconitum* ser. *Toxicum* (Ranunculaceae) in Europe. – *Phyton (Horn)* 33: 51–76.
- Myhal' A. V. (2009): *Crocus banaticus* J. Gay. – In: Diduch J. P. (ed.), *Červona knyha Ukraïny. Roslynnyj svit*, p. 118. Globalconsalting, Kyïv.
- Negrea B. M. & Pricop E. (2009a): The endemic plant species *Pietrosia levitomentosa*, a real conservation challenge. – *Advances Environm. Sci.* 2009, 1: 1–11.
- Negrea B. M. & Pricop E. (2009b): Rediscovery of *Pietrosia levitomentosa* E. I. Nyárády ex Sennik., an endemic and endangered plant species from Pietrosul Bistriței Mountain, Romania. – *Rom. J. Biol., Plant Biol.* 54: 101–114.
- Negrean G. (2004): Genul *Pietrosia* a fost reabilitat. – *Bul. Grăd. Bot. Iași* 12: 11–13.
- Negrean G. (2011): Addenda to „Flora Romaniaae“ volumes 1–12. Newly published plants, nomenclature, taxonomy, chorology and commentaries (Part 1). – *Kanitzia* 18: 89–194.
- Negrean G. & Oltean M. (1989): Endemite și zone endemoconservatoare din Carpații Sud-Estici [Endemics and endemoconservation areas in the South-Eastern Carpathians]. – *Ocrot. Nat.* 33: 15–25.
- Novikov A. V. (2013a): Vstup do taxonomii rodu *Aconitum* L. (Ranunculaceae) u flori Slovaččyny [Introduction to taxonomy of genus *Aconitum* L. (Ranunculaceae) in the flora of Slovakia]. – In: Aktual'ni problemy doslidžennja dovkillja. Zbirnyk naukovych prac' (za materialamy V Mižnarodnoi konferencii, 23–25 travnja 2013, Sumy), p. 303–306. Sumy.
- Novikov A. V. (2013b): Ranunculaceae Zachidnoi Ukraïny. I. Ključ dlja vyznačennja vydiv [Ranunculaceae of the Western Ukraine. Identification key]. – *Modern Phytomorphol.* 3: 297–319.
- Novikoff A. V. & Mitka J. (2011a): Taxonomy and ecology of the genus *Aconitum* L. in the Ukrainian Carpathians. – *Wulfenia* 18: 37–61.
- Novikov A. V. & Mitka J. (2011b). Rid *Aconitum* v Ukraïns'kyh Karpatach [The genus *Aconitum* in the Ukrainian Carpathians]. – *Biol. Stud.* 5: 153–172.
- Nyárády E. I. (1955): Cruciferae B. Juss. – In: Săvulescu T. (ed.), *Flora Republicii Populare Romîne* 3, p. 102–509. Edit. Acad. Rep. Pop. Romîne, București.

- Nyárády E. I. 1965: Compositae Giseke. – In: Nyárády E. I. (ed.), Flora Republicii Populare Romîne 10, p. 17–736. Edit. Acad. Rep. Pop. Romîne, București.
- Ochsmann J. (2000): Morphologische und molekularsystematische Untersuchungen an der *Centaurea stoebe* L.-Gruppe (Asteraceae-Cardueae) in Europa. – J. Cramer, Berlin & Stuttgart.
- Olšovská K. & Löser C. J. (2013): Mating system and hybridization of the *Cyanus triumfetti* and *C. montanus* groups (Asteraceae). – Folia Geobot. 48: 537–554.
- Olšovská K., Perný M., Kučera J. & Hodálová I. (2011): Biosystematic study of the *Cyanus triumfetti* group in Central Europe. – Preslia 83: 59–98.
- Olšovská K., Perný M., Španiel S. & Šingliarová B. (2012): Nuclear DNA content variation among perennial taxa of the genus *Cyanus* (Asteraceae) in Central Europe and adjacent areas. – Pl. Syst. Evol. 298: 1463–1482.
- Olšovská K., Šingliarová B., Kochjarová J., Labdík Z., Škodová I., Hegedúšová K. & Janišová M. (2015): Exploring patterns of variation within the central-European *Tephrosieris longifolia* agg.: karyological and morphological study. – Preslia 87: 163–194.
- Oltean M., Negrean G., Popescu A., Roman N., Dihoru Gh., Sanda V. & Mihăilescu S. (1994): Lista roșie a plantelor superioare din România [Red list of vascular plants of Romania]. – Studii, sinteze, documentații de ecologie 1: 1–52.
- Oprea A. (2005): Lista critică a plantelor vasculare din România [Critical list of vascular plants of Romania]. – Editura Universității „Alexandru Ioan Cuza”, Iași.
- Oprea A. (2007): Flora and vegetation of the Natural Reserve „Zugreni Gorges” (Suceava County). – Rom. J. Biol., Plant Biol. 52: 89–121.
- Oprea A. & Sîrbu C. (2012): The vascular flora of Rarău massif (Eastern Carpathian, Romania). Note I. – Mem. Sci. Sect. Romanian Acad., Biol. 35: 55–84.
- Oprea A. & Sîrbu C. (2013): The vascular flora of Rarău massif (Eastern Carpathian, Romania). Note II. – Mem. Sci. Sect. Romanian Acad., Biol. 36: 18–52.
- Pachswöll C. 2013. Lectotypification of *Doronicum carpaticum* (Griseb. et Schenk) Nyman (Asteraceae), with some remarks on its distribution and taxonomy. – Contr. Bot. Cluj 48: 7–14.
- Paclová L. (1999): *Ranunculus altitatisensis* Paclová et Murín. – In: Čerovský J. et al. (eds) (1999): Červená kniha vzácných a ohrozených druhů vyšších rostlin SR a ČR 5. Vyššie rastliny, p. 308. Príroda, Bratislava.
- Paucă A. & Beldie A. (1972): *Orchis* L. – In: Nyárády E. I. et al. (eds), Flora Republicii Socialiste România 12, p. 663–707. Edit. Acad. Rep. Soc. România, București.
- Paucă-Comănescu M. & Negrean G. (1994): Studiul ecologic al unor populații periclitate aparținând speciei endemice *Peucedanum rochelianum* Heuffel [Ecological research of some endangered populations belonging to endemic species *Peucedanum rochelianum* Heuffel]. – Ocrot. Nat. 38: 57–62.
- Pawłowski B. (1939): De positione systematica nec non affinitate Minuartiae oxypetalae (Wol.) Kulcz. – Acta Soc. Bot. Polon. 16: 153–167.
- Pawłowski B. (1952): *Alchemilla* carpatice et balcanice novae. – Bull. Acad. Polon. Sci. Lettr., Cl. Math.-Nat., Ser. Nat. (1951): 307–359.
- Pawłowski B. (1962): *Armeria pocutica* B. Pawł. – Fragm. Florist. Geobot. 8: 399–403.
- Pawłowski B. (1970): Remarques sur l'endémisme dans la flore des Alpes et des Carpates. – Vegetatio 21: 181–243.
- Pawlus M. (1988): Investigations on the genus *Alchemilla* L. in the Eastern Carpathians. – In: Marhold K. (ed.), Carpathian Flora, p. 82–94. ÚEBE CBEV SAV, Bratislava.
- Pečinka A., Suchánková M., Lysák M. A., Trávníček B. & Doležel J. (2006): Nuclear DNA content variation among Central European *Koeleria* taxa. – Ann. Bot. 98: 117–122.
- Peniașteková M. & Kliment J. (2002): *Draba* L. – In: Goliašová K. & Šipošová H. (eds), Flóra Slovenska V/4, p. 500–540. Veda, Bratislava.
- Perný M. & Michalková E. (2012): *Arenaria* L. – In: Goliašová K. & Michalková E. (eds), Flóra Slovenska VI/3, p. 219–230. Veda, Bratislava.
- Petrova A. (2011): *Linum uninerve* (Rochel) Borbás. – In: Peev D. & Vladimirov V. (eds), Red data book of the Republic of Bulgaria. 1. Plants and fungi. Digital edition, sine pag. Bulgarian Academy of sciences & Ministry of environment and water, Sofia. URL: <http://e-ecodb.bas.bg/rdb/en/vol1/Linunine.html>.
- Petrova A. & Vladimirov V. (eds) (2009): Red list of Bulgarian vascular plants. – Phytol. Balc. 15: 63–94.
- Piękoś-Mirkowa H. (2008): *Poa nobilis* Skalińska. – In: Mirek Z. & Piękoś-Mirkowa H. (eds), Czerwona księga Karpat Polskich. Rośliny naczyniowe, p. 546–547. Instytut Botaniki im. W. Szafera PAN, Kraków.
- Piękoś-Mirkowa H. & Mirek Z. (2003): Endemic taxa of vascular plants in the Polish Carpathians. – Acta Soc. Bot. Polon. 72: 235–242.
- Piękoś-Mirkowa H. & Mirek Z. (2009): Distribution patterns and habitats of endemic vascular plants in the Polish Carpathians. – Acta Soc. Bot. Polon. 78: 321–326.
- Piękoś-Mirkowa H., Mirek Z. & Miechówka A. (1996): Endemic vascular plants in the Polish Tatra Mts. Distribution and ecology. – Polish Bot. Stud. 12: 1–107.
- Plocek A. (1973): *Alchemilla suavis*, a new species from the Western Carpathians. – Oesterr. Bot. Z. 122: 195–198.
- Plocek A. (1983): Fifteen new species and varieties of *Alchemilla* (Rosaceae). – Folia Geobot. Phytotax. 18: 415–432.
- Plocek A. (1985): *Alchemilla connivens* Buser a príbuzné druhy v Karpatech a okolním území [*Alchemilla connivens* Buser and related species in the Carpathians and adjacent territories]. – Preslia 57: 9–30.

- Plocek A. (1986): New taxa and combinations in *Alchemilla* for the flora of Slovakia and the flora of Czechoslovakia (1). – Folia Geobot. Phytotax. 21: 423–427.
- Plocek A. (1990): New taxa and combinations in *Alchemilla* (Rosaceae) (2). – Folia Geobot. Phytotax. 25: 389–406.
- Plocek A. (1992): *Alchemilla* L. – In: Bertová L. (ed.), Flóra Slovenska IV/3, p. 250–373. Veda, Bratislava.
- Počynok T. V. & Prokopiv A. I. (2010): Pošyrennja vysokohirnych ridkisnych vydiv Asteraceae v Ukraïns'kykh Karpatach [Distribution of rare high-mountain species of Asteraceae in the Ukrainian Carpathians]. – In: Roslynnij svit u Červonij knyzi Ukraïny: vprovadžennja Hlobal'noï stratehii zberežennja roslyn. Materialy mižnarodnoï konferencii, (11–15 žovtnja 2010 r., Kiïv), p. 168–171. Alterpress, Kiïv.
- Podlech D. (2008): The genus *Astragalus* L. (Fabaceae) in Europe with exclusion of the former Soviet Union. – Feddes Repert. 119: 310–387.
- Podlech P. (2011): Thesaurus Astragalorum. Index of all taxa described within the genus *Astragalus* L. and other genera but belonging to the genus *Astragalus*. Taxa of the Old World and related taxa of the New World. Last updated version december 2011. – URL: http://www.sysbot.biologie.uni-muenchen.de/de/personen/podlech/thesaurus_astragalus.pdf.
- Pogan E. (1977): Kariologia flory polskiej [Karyology of the Polish flora]. – In: Szafer W. & Zarzycki K. (eds), Szata roślinna Polski 1. Ed. 3, p. 207–236. Państw. Wydaw. Nauk., Warszawa.
- Popescu A. & Sanda V. (1998): Conspectul florei cormofitelor spontane din România [Enumeration of spontaneous vascular plants of Romania]. – Editura Universităţii din Bucureşti, Bucureşti.
- Popescu G., Costache I., Rădoşoiu D. & Boruz V. (2003): The ecology, coenology and chorology of the endemites and subendemites in the region of Oltenia (Romania). – Contr. Bot. Cluj 38: 147–156.
- Proc' P. H. & Kiš P. J. (2009): *Lathyrus transsilvanicus* (Spreng.) Rchb. – In: Diduch J. P. (ed.), Červona knyha Ukraïny, p. 472. Roslynnij svit. Globalconsalting, Kyïv.
- Procházka F. & Chrtek J. jr. (1999): *Hieracium albinum* Fries. – In: Čeřovský J. et al. (eds), Červená kniha vzácných a ohrožených druhů vyšších rostlin SR a ČR 5. Vyššie rastliny, p. 183. Příroda, Bratislava.
- Prodan I. (1961): Dipsacaceae B. Juss. – In: Nyárady E. I. (ed.), Flora Republicii Populare Romîne 8, p. 640–685. Edit. Acad. Rep. Pop. Romîne, Bucureşti.
- Prokudin J. N. (ed.) (1987): Opredelitel' vyššich rastenij Ukraïny [Identification key of vascular plants of Ukraine]. – Naukova dumka, Kiïv.
- Prokudin J. N., Vovk A. G., Petrova O. A., Ermolenko E. D. & Verničenko J. D. (1977): Zlaki Ukraïny [The grasses of Ukraine]. – Naukova dumka, Kiev.
- Răvărut M. (1960): *Euphrasia* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 7, p. 578–593. Edit. Acad. Rep. Pop. Romîne, Bucureşti.
- Rola K. (2014): Morphometry and distribution of *Senecio nemorensis* agg. species (Asteraceae) in Poland. – Polish Bot. J. 59: 1–18.
- Roman N., Roman S. & Heltmann H. (1996): Beiträge zur Verbreitung von Pflanzenarten in der Siebenbürgischen Heide und den angrenzenden Gebieten. – Stapfia 45: 135–150.
- Ronikier M. & Szeląg Z. (2008): Differentiation among disjunct populations of agamospermous species of *Hieracium* section *Cernua* (Asteraceae) in Central European subalpine habitats. – Bot. J. Linn. Soc. 158: 93–105.
- Rostański K. (1967): Rodzina Valerianaceae [Family Valerianaceae]. – In: Pawłowski B. (ed.), Flora polska 11, p. 338–357. Państw. Wydaw. Nauk., Warszawa.
- Rostański K. (1970): Rozmieszczenie gatunków rodzaju *Valeriana* L. w Polsce i na sąsiednich terytoriach ZSSR [The distribution of the species of the genus *Valeriana* L. in Poland and in adjacent territories of the U.S.S.R.]. – Fragm. Florist. Geobot. 16: 209–246.
- Rybczyński J. J., Davey M. R. & Miłkuła A. (eds) (2014): The Gentianaceae. Volume 1: Characterization and ecology. – Springer, Berlin Heidelberg.
- Sanda V., Popescu A., Doltu M. I. & Donița N. (1983): Caracterizarea ecologică și fitocenologică a speciilor spontane din flora României [Characterisation of the ecology and phytosociology for some wild species of the Romanian flora]. – Stud. Comun. Muz. Brukenthal, Ști. Nat., suppl. 25: 5–126.
- Sârbu I. & Lupu I. (1989): A new habitat with *Draba dorneri* Heuff. – Analele Ști. Univ. Iași, Ser. Biol. 35: 39–40.
- Sârbu I., Ștefan N. & Oprea A. (2013): Plante vasculare din România. Determinator ilustrat de teren [Vascular plants of Romania. An illustrated field guide]. – Victor B Victor, Bucureşti.
- Schuhwerk F. & Lippert W. (1999): Chromosomenzahlen von *Hieracium* (Compositae, Lactucaceae) Teil. 3. – Sendtnera 6: 197–214.
- Schur F. (1866): Enumeratio Plantarum Transsilvaniae. – Guilielmum Braumüller, Vindobonae.
- Sennikov A. N. (1999a): Pietrosia Nyárady – a restored genus of the subtribe Hieraciinae. – Komarovia 1: 77–78.
- Simon T. (1992): A Magyarországi edényes flóra határozója. Harasztok – virágos növények [Field guide to vascular flora of Hungary. Pteridophytes – Angiosperms]. – Tankönyvkiadó, Budapest.

- Simonkai L. (1887) [„1886”]: Erdély edényes flórájának helyesbitett foglalata [Enumeratio Florae Transsilvanicae vespulosae critica]. – Budapest.
- Skalický V. (1990): Rod *Aconitum* v Československu [Die Gattung *Aconitum* in der Tschechoslowakei]. – Zprávy Českoslov. Bot. Společn. 25, 2: 1–27.
- Skalińska M. (1955): *Poa nobilis* n. sp., a new viviparous species of the High Tatra. – Acta Soc. Bot. Polon. 24: 749–761.
- Slavkovska V., Zlatković B., Bräuchler Ch., Stojanović D., Tzakou O. & Couladis M. (2013): Variations of essential oil characteristics of *Clinopodium pulegium* (Lamiaceae) depending on phenological stage. – Bot. Serb. 37: 97–104.
- Slovák M., Kučera J., Turis P. & Zozomová-Lihová J. (2012): Multiple glacial refugia and postglacial colonization routes inferred for a woodland geophyte, *Cyclamen purpurascens*: patterns concordant with the Pleistocene history of broadleaved and coniferous tree species. – Biol. J. Linn. Soc. 105: 741–760.
- Smejkal M. (1963): Taxonomická studie československých druhů rodu *Euphrasia* L. [Taxonomic study of Czechoslovak species of *Euphrasia* L.]. – Biol. Práce 9/9: 4–82.
- Smejkal M. & Čerovský J. (1999): *Euphrasia exaristata* Smejkal. – In: Čerovský J. et al. (eds), Červená kniha vzácných a ohrozených druhů vyšších rostlin SR a ČR 5. Vyššie rastliny, p. 154. Príroda, Bratislava.
- Soják J. (1983b): Einige Bemerkungen zur Flora der UdSSR (2.). – Sborn. Nár. Muz., Ser. B 39: 53–60.
- Somogyi G., Barina Z. & Höhn M. (2012): A Magyarországi tollas szegfűvek taxonómiai megítélésének természetvédelmi vonatkozásai [The conservation biological aspects of taxonomic treatment of the Hungarian feather carnations]. – Rosalia (Budapest) 8: 293–306.
- Soó R. (1964): A magyar flóra és vegetáció rendszertani-növényföldrajzi kézikönyve 1 [Synopsis systematico-geobotanica florum vegetationisque Hungariae 1]. – Akadémiai kiadó, Budapest.
- Soó R. (1967): Die *Dactylorhiza*-Arten und -Formen in der Flora Rumäniens. – Rev. Roum. Biol., Bot. 12: 225–231.
- Soó R. (1968): A magyar flóra és vegetáció rendszertani-növényföldrajzi kézikönyve 3 [Synopsis systematico-geobotanica florum vegetationisque Hungariae 3]. – Akadémiai kiadó, Budapest.
- Soó R. (1973): A magyar flóra és vegetáció rendszertani-növényföldrajzi kézikönyve 5 [Synopsis systematico-geobotanica florum vegetationisque Hungariae 5]. – Akadémiai kiadó, Budapest.
- Soó R. (1980a): A magyar flóra és vegetáció rendszertani-növényföldrajzi kézikönyve 6 [Synopsis systematico-geobotanica florum vegetationisque Hungariae 6]. – Akadémiai kiadó, Budapest.
- Soó R. (1980b): *Dactylorhiza* Necker ex Nevski. – In: Tutin T. G. et al. (eds), Flora Europaea 5, p. 333–337. Cambridge University Press, Cambridge.
- Starmühler W. (1996a): *Aconita rarissima* selecta. – Fritschiana 7: 11–39.
- Starmühler W. (1996b): *Delphinia rarissima* selecta. – Fritschiana 7: 41–45.
- Starmühler W. (1997): Ranunculaceae subtrib. Delphiniinae exsiccatae, Fasc. 2 (Nr. 21–40). – Fritschiana 10: 1–11.
- Starmühler W. (1998a): Ranunculaceae subtrib. Delphiniinae exsiccatae, Fasc. 3 (Nr. 41–60). – Fritschiana 13: 1–10.
- Starmühler W. (1998b): *Aconita rarissima* selecta (1998). – Fritschiana 13: 11–39.
- Starmühler W. (1999): *Aconita rarissima* selecta (1999). – Fritschiana 18: 11–38.
- Starmühler W. (2000): Fortschritte in der Systematik der Gattung *Aconitum* im Siebenbürgen. – In: Heltmann H. & Killyen H. (eds), Naturwissenschaftliche Forschungen über Siebenbürgen VI. Beiträge zur Geographie, Botanik, Zoologie und Paläontologie, p. 9–24. Böhlau Verlag, Köln, Weimar & Wien.
- Starmühler W. (2004): *Aconita rarissima* selecta (2004). – Fritschiana 48: 9–39.
- Starmühler W. & Mitka J. (2001): Systematics and chorology of *Aconitum* sect. *Napellus* (Ranunculaceae) and its hybrids in the Northern Carpathians and Forest Carpathians. – Thaiszia-J. Bot. 10: 115–136.
- Staszkievicz J. (2009): *Euphrasia exaristata* (Scrophulariaceae), a Western Carpathian endemic species new to the flora of Poland. – Polish Bot. J. 54: 27–29.
- Ștefureac T. I. & Țăcină A. (1979): Cytotaxonomical and chorological investigations of the endemic taxon *Hieracium pojoritense* Wol. – Rev. Roum. Biol., Biol. Veg. 24: 109–120.
- Stevanović V. (2013): *Tulipa hungarica*. – The IUCN Red List of Threatened Species. Version 2014.2. URL: <http://www.iucnredlist.org/details/162072/0> (accessed 26 October 2014).
- Stevanović V., Niketić M. & Lakušić D. (1991): Chorological additions to the flora of eastern Yugoslavia. – Fl. Medit. 1: 121–142.
- Stojko S. M. & Tasenkewitsch L. O. (1991): Pflanzengeographische Stellung und Schutz von Flora und Vegetation der Ukrainischen Karpaten. – Verh. Zool.-Bot. Ges. Österreich 128: 165–172.

- Stoyko S. M. & Tasenkevich L. (1993): Some aspects of endemism in the Ukrainian Carpathians. – *Fragm. Florist. Geobot.*, Suppl. 2 (1): 343–353.
- Strajeru S. & Stevanović V. 2013. *Barbarea lepuznica*. – The IUCN Red List of Threatened Species. Version 2014.2. URL: <http://www.iucnredlist.org/details/176458/0> (accessed 14 October 2014).
- Strid A. & Tan K. (eds) (1991): *Mountain flora of Greece*. 2. – Edinburgh University Press, Edinburgh.
- Șuteu A., Gafta G. & Groza G. (2003): Reproductive and distribution patterns in two populations of the rare endemic *Astragalus peterfii* Jáv. (Transylvania, Romania). – *Bocconeia* 16: 455–464.
- Șuteu D. C. (2012): Molecular taxonomy studies on several endemic and rare plants species from Romania. – Summary of the PhD thesis, Babeș-Bolyai University, Cluj-Napoca.
- Șuteu D., Pușcaș M., Băcilă J., Coste A., Stoica I. A., Hurdu B. I., Ursu T. & Coldea Gh. (2011): Does *Primula intricata* Gren. et Godr. merit species rank? A taxonomic revision based on nrDNA, cpDNA and AFLP data. – *Not. Bot. Hort. Agrobot. Cluj* 39: 24–29.
- Șuteu D., Pușcaș M., Filipaș L., Băcilă I., Hurdu B.-I. & Coldea G. (2013): *Primula leucophylla* in South-Eastern Carpathians: a genetic and morphometric study within *Primula elatior* group. – *Acta Biol. Cracov., Bot.* 55, Suppl. 1: 69.
- Syčák N. M. (1992): Novyj vyd rodu *Alchemilla* L. (Rosaceae) z Ukraïnskych Karpat [New species of *Alchemilla* L. (Rosaceae) from the Ukrainian Carpathians]. – *Ukrains'k. Bot. Zhurn.* 49: 99–101.
- Syčák N. M. (2002): Ridkisi ta endemični vydy rodu *Alchemilla* L. (Rosaceae) u flori Ukraïns'kych Karpat [Rare and endemic species of genus *Alchemilla* L. (Rosaceae) in the flora of the Ukrainian Carpathians]. – In: Hamor F. D. (ed.), *Mountains and people (in the context of sustainable development)*. Proceedings of the International conference, October 14th–18th, 2002, Vol. 2, p. 473–477. Rakhiv.
- Syčák N. M. (2011): Ključ dlja vyznačennja vydiv rodu *Alchemilla* L. (Rosaceae) flory Ukraïnskych Karpat [Key for determination of the species of genus *Alchemilla* L. (Rosaceae) in the flora of the Ukrainian Carpathians]. – *Naukovi osnovy zberežennja biotyčnoï riznomanitnosti* 2 (9): 215–228.
- Szeląg Z. (2003a): Validation of Nyárády's names in *Hieracium* sect. *Cernua* (Asteraceae). – *Polish Bot. J.* 48: 11–12.
- Szeląg Z. (2003b): A synopsis of *Hieracium* sect. *Cernua* (Asteraceae). – *Polish Bot. J.* 48: 89–97.
- Szeląg Z. (2004a): Taxonomic and nomenclatural notes on *Hieracium pawlowskianum* (Asteraceae) and its relatives. – *Polish Bot. J.* 49: 11–14.
- Szeląg Z. (2004b): Taxonomic and nomenclatural notes on *Hieracium silesiacum* (Asteraceae). – *Polish Bot. J.* 49: 15–20.
- Szeląg Z. (2006a): Taxonomic and nomenclatural notes on *Hieracium tubulare* (Asteraceae) with description of a new species from the Eastern Carpathians. – *Ann. Bot. Fenn.* 43: 310–314.
- Szeląg Z. (2006b): Taxonomic revision of *Hieracium* sect. *Cernua* (Asteraceae) in the Carpathians, Sudetes and Alps. – *Polish Bot. J.* 51: 97–153.
- Szeląg Z. (2007): Typification of the *Hieracium* (Asteraceae) names described by E. Wołoszczak from the Eastern Carpathians. – *Polish Bot. J.* 52: 99–118.
- Szeląg Z. (2008): Taxonomic and nomenclatural notes on *Pilosella alpicola* agg. (Asteraceae) in the Balkans and Carpathians. – *Ann. Bot. Fenn.* 45: 301–306.
- Szeląg Z. (2010): A new species in the *Hieracium lycopifolium* agg. (Asteraceae) from the Western Carpathians. – *Acta Soc. Bot. Polon.* 79: 245–248.
- Szeląg Z. (2011): *Hieracia balcanica* VII. Identity and typification of *Hieracium bohatschianum* (Asteraceae). – *Polish Bot. J.* 56: 155–157.
- Szeląg Z. (2012): New chorological data on *Hieracium coldei* (Asteraceae). – *Polish Bot. J.* 57: 487–489.
- Szeląg Z. (2013): Taxonomic and nomenclatural notes on *Hieracium tomosense* (Asteraceae), a neglected Carpathian endemic. – *Polish Bot. J.* 58: 359–362.
- Šafářová L., Duchoslav M., Jandová M. & Krahulec F. (2011): *Allium oleraceum* in Slovakia: cytotype distribution and ecology. – *Preslia* 83: 513–527.
- Šeffer J., Lasák R., Šefferová-Stanová V., Janák M. & Guttová A. (2010): Towards an Ecological Network for the Carpathians II. – *Carpathian Ecoregion Initiative*, Bratislava.
- Šingliarová B. & Mráz P. (2009): A taxonomic revision of the *Pilosella alpicola* group in the Carpathians. – *Preslia* 81: 23–41.
- Šingliarová B., Chrtek J. jr. & Mráz P. (2008): Loss of genetic diversity in isolated populations of an alpine endemic *Pilosella alpicola* subsp. *ullepitschii*: effect of long-term vicariance or long-distance dispersal? – *Pl. Syst. Evol.* 275: 181–191.
- Šingliarová B., Chrtek J., Plačková I. & Mráz P. (2011a): Allozyme variation in diploid, polyploid and mixed-ploidy populations of the *Pilosella alpicola* group (Asteraceae): relation to morphology, origin of polyploids and breeding system. – *Folia Geobot.* 46: 387–410.
- Šingliarová B., Hodálová I. & Mráz P. (2011b): Biosystematic study of the diploid-polyploid *Pilosella alpicola* group with variation in breeding system: Patterns and processes. – *Taxon* 60: 450–470.
- Šingliarová B., Šuvada R. & Mráz P. (2013): Allopatric distribution, ecology and conservation status of the *Pilosella alpicola* group (Asteraceae). – *Nord. J. Bot.* 31: 122–128.

- Šípošová H., Bernátová D., Mráz P., Kliment J., Goliašová K. & Uhlířová J. (2004a): Taxóny cievnatých rastlín, ktoré boli opísané z Veľkej Fatry. 1. časť [Vascular plant taxa described from the Veľká Fatra Mts. Part 1]. – In: Kadlečík J. (ed.), Turiec a Fatra 2004, p. 77–85. Správa Národného parku Veľká Fatra, Vrútky.
- Šípošová H., Bernátová D., Mráz P., Kliment J., Goliašová K. & Uhlířová J. (2004b): Taxóny cievnatých rastlín, ktoré boli opísané z Nízkych Tatier. 1. časť [Vascular plant taxa described from the Nízke Tatry Mts. Part 1]. – *Príroda Nízkych Tatier* 1: 129–136.
- Šljakov P. N. (1989): *Hieracium* L. – In: Cvelev N. N. (ed.), Flora evropejskoj časti SSSR 8, p. 140–300. Nauka, Leningrad.
- Štech M. & Drábková L. (2005): Morphometric and RAPD study of *Melampyrum sylvaticum* group in the Sudeten, the Alps and Carpathians. – *Folia Geobot.* 40: 177–193.
- Štěpánek J. (1983): Eine neue Art der Gattung *Knautia* (Dipsacaceae) aus Westkarpaten. – *Preslia* 55: 1–8.
- Štěpánek J. (1985): *Knautia* L. – In: Bertová L. (ed.), Flóra Slovenska IV/2, p. 154–177. Veda, Bratislava.
- Štěpánek J. (1997): *Knautia* L. – In: Slavík B. (ed.), Květena České republiky 5, p. 543–554. Academia, Praha.
- Štěpánek J., Kirschner J., Jarolímová V. & Kirschnerová L. (2011): *Taraxacum nigricans*, *T. alpestre* and allies in the *Taraxacum* sect. *Alpestris*: taxonomy, geography and conservation status. – *Preslia* 83: 537–564.
- Štorchová J., Chrtěk J. jr., Bartish V., Tetera M., Kirschner J. & Štěpánek J. (2002): Genetic variation in agamospermous taxa of *Hieracium* sect. *Alpina* (Compositae) in the Tatry Mts. (Slovakia). – *Pl. Syst. Evol.* 235: 1–17.
- Tacik T. (1980): *Taraxacum* Wiggers. – In: Jasiewicz A. (ed.), Flora Polska 14, p. 7–199. Państw. Wydaw. Nauk., Warszawa.
- Tan K. & Vladimirov V. (2001): *Swertia punctata* (Gentianaceae) in Bulgaria. – *Bocconeia* 13: 461–466.
- Tasenevich L. (2002): Red list of vascular plants of the Carpathian mountains. – State museum of natural history NAS of Ukraine, Lviv.
- Tasenevych L. (2003b): Rozmaïtja flory sudynnych roslyn v Ukraïns'kych Karpatach [Vascular plants flora diversity of the Ukrainian Carpathians]. – *Pratsi Nauk. Tovar. Schevchenka* 12: 147–157.
- Tasenevich L. (2011): Endemism in the Carpathian flora – a chorological aspect. – In: Zemanek B. (ed.), Geobotanist and Taxonomist. A volume dedicated to Professor Adam Zajac on the 70th anniversary of his birth, p. 157–168. Institute of Botany Jagellonian University, Cracow.
- Tasenevych L. (2014): Endemizm flory Karpat – ekoloïiïni ta choroloïiïni osoblyvosti [Endemism of the Carpathians' flora – ecological and chorological peculiarities]. – *Pratsi Nauk. Tovar. Schevchenka* 39: 22–36.
- Teppner H. (2004): A review of new results in *Nigritella* (Orchidaceae). – *Spraw. Posiedz. Komis. Nauk.* 46: 111–116.
- Teppner H. & Klein E. (1998): Etiam atque etiam – *Nigritella* versus *Gymnadenia*: Neukombinationen und *Gymnadenia dolomitensis* spec. nova (Orchidaceae-Orchideae). – *Phyton* (Horn) 38: 220–224.
- Teppner H., Klein E., Drescher A. & Zagulskij M. (1994): *Nigritella carpatica* (Orchidaceae-Orchideae) – ein Reliktenendemit der Ost-Karpaten. – *Phyton* (Horn) 34: 169–187.
- Těšitel J. & Štech M. (2007): Morphological variation in the *Melampyrum sylvaticum* group within the transitional zone between *M. sylvaticum* s. str. and *M. herbichii*. – *Preslia* 79: 83–99.
- Těšitel J., Malinová T., Štech M. & Herbstová M. (2009): Variation in the *Melampyrum sylvaticum* group in the Carpathian and Hercynian region: two lineages with different evolutionary histories. – *Preslia* 81: 1–22.
- Tomšovic P. (1988): Carpathian populations of *Erysimum odoratum* agg. – In: Marhold K. (ed.), Carpathian Flora, p. 127–130. ÚEBE CBEV SAV, Bratislava.
- Tosheva A., Assyov B & Denchev C. M. (2011): *Lathyrus transsilvanicus* (Spreng.) Fritsch. – In: Peev D. & Vladimirov V. (eds), Red data book of the Republic of Bulgaria. 1. Plants and fungi. Digital edition. Bulgarian Academy of sciences & Ministry of environment and water, Sofia. URL: <http://e-ecodb.bas.bg/rdb/en/vol1/Lattrans.html>.
- Trávníček B., Duchoslav M., Šarhanová P. & Šafařová L. (2010): Squills (*Scilla* s. lat., Hyacinthaceae) in the flora of the Czech Republic, with taxonomical notes on Central-European squill populations. – *Acta Mus. Morav., Sci. Biol.* 94 (2009): 157–205.
- Tribsch A. & Schönswetter P. (2003): Patterns of endemism and comparative phylogeography confirm palaeoenvironmental evidence for Pleistocene refugia in the Eastern Alps. – *Taxon* 52: 477–497.
- Turis P. (2009): Ekobiológia, vybrané populačné charakteristiky, rozšírenie a ochrana cyklámenu fatranského vo východnej časti areálu [Ecobiology, selected population characteristics, distribution and protection of *Cyclamen fatrense* in the eastern part of its distribution area]. – PhD thesis, Institute of Landscape Ecology, Slovak Academy of Sciences, Bratislava.

- Turis P., Eliáš P. jun., Schmotzer A., Király G., Schneider E., Kuciel H., Szewczyk M., Kozurak A., Antosyak T., Voloshchuk M., Lazarević P. & Lustyk P. (2014): Red List of vascular plants of the Carpathians, p. 44–105. – In: Kadlečík J. (ed.), Carpathian Red List of forest habitats and species. Carpathian list of invasive alien species. (Draft). The State Nature Conservancy of the Slovak Republic.
- Turlaj O. I. (2011): Osoblyvosti petioljarnoj struktury lystkiv dejakych vydiv žovteciv iz sekcii *Acris* Shur [The features of anatomical structure of leaves of some species *Ranunculus* from the section *Acris* Schur]. – Biol. Syst. 3, 1: 97–100.
- Tutin T. G. (1972): *Gentiana* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 59–63. Cambridge University Press, Cambridge.
- Tutin T. G. & Akeyrod J. R. (1993): *Ranunculus* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 269–286. Cambridge University Press, Cambridge.
- Uhlířová J. & Bernátová D. (2004): K floristicko-fytocenologickej štruktúre reliktnej kosodreviny vo Veľkej Fatre [The floristic and phytosociological structure of the relict dwarf pine scrub in the Veľká Fatra Mts]. – In: Kadlečík J. (ed.), Turiec a Fatra 2004, p. 67–75. Správa Národného parku Veľká Fatra, Vrútky.
- Váczy C. & Beldie A. (1976): Stațiuni noi de specii și subspecii [New localities of species and subspecies]. – In: Beldie A. & Morariu I. (eds), Flora Republicii Socialiste România 13, p. 54–65. Edit. Acad. Rep. Soc. România, București.
- Vasil'eva I. M. (2001): *Aquilegia* L. – In: Cvelev N. N. (ed.), Flora Vostočnoj Evropy 10, p. 183–186. Tovariščestvo naučnych izdanij KMK, Moskva & Sankt-Peterburg.
- Vasil'eva I. M. (2012): *Aquilegia* L. – In: Gel'tman D. V. (ed.), Konspekt flory Vostočnoj Evropy, p. 154. Tovariščestvo naučnych izdanij KMK, Sankt-Peterburg & Moskva.
- Vázquez F. M. & Gutiérrez M. (2011): Classification of species of *Stipa* with awns having plumose distal segments. – Telopea 13: 155–176.
- Velev N. I. & Apostolova I. I. (2009): A review of *Potentillo ternatae-Nardion strictae* alliance. – Hacquetia 8: 49–66.
- Vladimirov V., Dane F., Matevski V., Stevanović V. & Tan K. (eds) (2011): New floristic records in the Balkan: 15. – Phytol. Balc. 17: 129–156.
- Vojtkó A., Sass-Gyarmati A., Pócs T. & Dulai S. (2012): Critical assessment of the flora of the Vargyas Gorge (Eastern Carpathians). – Acta Biol. Plant. Agriensis 2: 27–74.
- Volgin S. A. & Syčák N. N. (1989a): Manžetki (*Alchemilla* L., Rosaceae) Ukrainskich Karpat. I. Sekcija *Alchemilla*, podsekcija *Calycinae* (Buser) Camus [The lady's mantles (*Alchemilla* L., Rosaceae) from the Ukrainian Carpathians. I. Section *Alchemilla*, subsection *Calycinae* (Buser) Camus]. – Bjul. Moskov. Obšč. Ispyt. Prir., Biol. 94: 71–79.
- Volgin S. A. & Syčák N. N. (1989b): Manžetki (*Alchemilla* L., Rosaceae) Ukrainskich Karpat. II. Sekcija *Alchemilla*, podsekcija *Euvulgares* Camus (řady *Glabrae* Pawł. i *Subglabrae* Pawł.) [Lady's-mantles (*Alchemilla* L., Rosaceae) of the Ukrainian Carpathians. II. Section *Alchemilla* subsection *Euvulgares* Camus (lines *Glabrae* Pawł. and *Subglabrae* Pawł.)]. – Bjul. Moskov. Obšč. Ispyt. Prir., Biol. 94: 86–94.
- Walters S. M. & Akeyrod J. R. (1993): *Draba* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 372–377. Cambridge University Press, Cambridge.
- Walters S. M. & Pawłowski B. (1968): *Alchemilla* L. – In: Tutin T. G. et al. (eds), Flora Europaea 2, p. 48–64. Cambridge University Press, Cambridge.
- Webb D. A. (1993a): *Saxifraga* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 437–458. Cambridge University Press, Cambridge.
- Witkowski Z. J., Król W. & Solarz W. (eds) (2003): Carpathian list of endangered species. – WWF and Institute of Nature Conservation, Polish Academy of Sciences, Vienna & Krakow.
- Wróbel I. & Zarzycki K. (2008): *Taraxacum pieninicum* Pawł. – In: Mirek Z. & Piękoś-Mirkowa H. (eds), Czerwona księga Karpat Polskich. Rośliny naczyniowe, p. 406–408. Instytut Botaniki im. W. Szafera PAN, Kraków.
- Zahn K. H. (1911): Beiträge zur Kenntnis der Hieracien Ungarns, Galiziens u. der Balkanländer VI. – Magyar Bot. Lapok 10: 121–174.
- Zahn K. H. (1927): Beiträge zur Kenntnis der Hieracien Ungarns und der Balkanländer VII. – Magyar Bot. Lapok 25 (1926): 121–174.
- Zahn K. H. (1930–1935): *Hieracium*. – In: Graebner P. jr. (ed.), Synopsis der mitteleuropäischen Flora 12/2. – Gebrüder Borntraeger Verlag, Leipzig.
- Zahn K. H. (1936–1938): *Hieracium*. – In: Graebner P. jr. (ed.), Synopsis der mitteleuropäischen Flora 12/3. – Gebrüder Borntraeger Verlag, Leipzig.
- Zahradníková K., Peniašteková M. & Ľavoda O. (2002): *Hesperis* L. – In: Goliašová K. & Šipošová H. (eds), Flóra Slovenska V/4, p. 230–251. Veda, Bratislava.
- Zapałowicz H. (1913): Krytyczny przegląd roślinności Galicyi XXVII [Conspectus florae Galiciae criticus XXVII]. – Rozpr. Wydz. Mat.-Przyr. Akad. Umiejętn., Ser. 3B, 13: 29–49.
- Zarzycki K. (1986): Lista wymierających i zagrożonych roślin naczyniowych Polski. List of threatened vascular plants in Poland. – In: Zarzycki K. & Wojewoda W. (eds), Lista roślin wymierających i zagrożonych w Polsce. List of threatened plants in Poland, p. 11–27. Państw. Wydaw. Nauk., Warszawa.
- Zarzycki K. (2008): *Carduus ×lobulatus* Borbás. – In: Mirek Z. & Piękoś-Mirkowa H. (eds), Czerwona księga Karpat Polskich. Rośliny naczyniowe, p. 358–359. Instytut Botaniki im. W. Szafera PAN, Kraków.
- Zelený V. (1970): Taxonomisch-chorologische Studie über die Art *Leucanthemum rotundifolium* (W. K.) DC. – Folia Geobot. Phytotax. 5: 369–400.
- Zhang L.-B. & Kadereit J. W. (2002): The systematics of *Soldanella* (Primulaceae) based on morphological and molecular (ITS, AFLPs) evidence. – Nord. J. Bot. 22: 129–169.
- Ziman S. M. (2009): *Astragalus krajinae* Domin. – In: Diduch J. P. (ed.), Červona knyha Ukraïny. Roslynnij svit, p. 442. Globalconsalting, Kyïv.

- Ziman S. & Derbak M. (2013): About the conservation of the biodiversity in the high-mountain flora of the Ukrainian Carpathians and protection of the „hot spots“. – Nat. Montenegrina 12: 639–647.
- Ziman S. M., Bulach O. V. & Hamor A. F. (2006): Pro rідkisni vysokohirni vydy kvitkovych roslyn u flory Ukraїns'kych Karpat – vikarianty vydiv z šyrokymy arealamy ta ekolohieju [About rare high-mountain species of the flowering plants within the flora of the Ukrainian Carpathians]. – Nauk. Zap. Derzhavn. Pryr. Muz., Bot. 22: 25–34.
- Ziman S. M., Bulach O. V. & Tjuch J. J. (2009): *Aquilegia transsilvanica* Schur. – In: Diduch J. P. (ed.), Červona knyha Ukraїny. Roslynnyj svit, p. 556. Globalconsalting, Kyїв.

Electronic Appendix 2. – “Critical” (unclear) endemic taxa

Taxon	Synonyms	References
<i>Aconitum moldavicum</i> subsp. <i>simonkaianum</i> (Gáyer) Starm. ²⁰⁰	<i>Aconitum simonkaianum</i> Gáyer	Grițescu 1953; Starmühler 1998a; Mitka 2008
<i>Allium zahariadi</i> Májovský subsp. <i>zahariadi</i> ²⁰¹	<i>Allium ochroleucum</i> subsp. <i>pseudosuaveolens</i> Zahar.; <i>A. ericetorum</i> subsp. <i>pseudosuaveolens</i> (Zahar.) Ciocârlan	Zahariadi 1966; Májovský & Murín 1985; Somogyi 1999a; Ciocârlan 2009; Sârbu et al. 2013
<i>Allium zahariadi</i> subsp. <i>michalkoi</i> Májovský ²⁰²		Májovský & Murín 1985; Somogyi 1999a, b, 2002
<i>Bupleurum falcatum</i> subsp. <i>dilatatum</i> Schur ²⁰³	<i>Bupleurum dilatatum</i> (Schur) Baksay	Tutin 1968; Kliment 1999; Oprea 2005
<i>Carduus fissurae</i> Nyár. ²⁰⁴		Nyárády 1965; Beldie & Alexandrescu 1976; Ciocârlan 2009
<i>Carex sempervirens</i> subsp. <i>tatrorum</i> (Zapał.) Pawł. ²⁰⁵		Kliment 1999; Marhold et al. 2007; Luceño et al. 2008
<i>Carex sempervirens</i> subsp. <i>pseudotristis</i> (Domin) Pawł. ²⁰⁶	<i>Carex sempervirens</i> var. <i>pseudotristis</i> Domin	Domin 1931b; Luceño et al. 2008
<i>Centaurea globurensis</i> Nyár. ²⁰⁷		Dostál 1976; Dihoru & Pârvu 1987; Greuter in Euro+Med
<i>Centaurea pugioniformis</i> Nyár. ²⁰⁸		Prodan & Nyárády 1964; Dostál 1976; Ciocârlan 2009
<i>Dactylorhiza carpatica</i> (Batoušek et Kreutz) P. Delforge ²⁰⁹		Kubát 2010; Kaplan 2012
<i>Dianthus carpaticus</i> Woł. ^{210, 211}	<i>Dianthus carthusianorum</i> var. <i>carpaticus</i> (Woł.) Zapał.; <i>D. bucovinensis</i> (Zapał.) Klokov	Klokov 1952; Tutin 1964; Jalas & Suominen 1986; Fedorončuk & Diduch 2002a; Kuz'mina 2004; Fedorončuk 2009
<i>Hieracium amoenanthes</i> Nyár. et Zahn ²¹²		Zahn 1936; Nyárády 1965
<i>Hieracium borzae</i> Nyár. et Zahn ²¹³		Zahn 1938; Nyárády 1965
<i>Hieracium breazense</i> Nyár. ²¹⁴		Zahn 1938; Nyárády 1965
<i>Hieracium caesiogenum</i> Woł. et Zahn ²¹⁵		Zahn 1936; Nyárády 1965; Šljakov 1989
<i>Hieracium calcogeton</i> (Zahn) Greuter ²¹⁶		Zahn 1938; Nyárády 1965
<i>Hieracium chloribracteum</i> Degen et Zahn ²¹⁷		Zahn 1936; Nyárády 1965
<i>Hieracium dentatum</i> subsp. <i>sarmaticum</i> Zahn ²¹⁸	<i>Hieracium sarmaticum</i> (Zahn) Šljakov	Zahn 1927, 1935; Nyárády 1965; Šljakov 1989
<i>Hieracium grecescui</i> Nyár. et Zahn ²¹⁹		Zahn 1938; Nyárády 1965
<i>Hieracium klopotivae</i> Pax ²²⁰		Nyárády 1965; Szeląg 2014 in litt.
<i>Hieracium krizsnae</i> Lengyel et Zahn ²²¹		Zahn 1927; Šípošová et al. 2004a
<i>Hieracium mukaczense</i> Üksip ²²²		Juksip 1959; Šljakov 1989
<i>Hieracium nyaradyanum</i> Zahn ²²³		Zahn 1938; Nyárády 1965
<i>Hieracium pelagae</i> Degen et Zahn ²²⁴		Zahn 1938; Nyárády 1965
<i>Hieracium peterfii</i> Nyár. et Zahn ²²⁵		Zahn 1938; Nyárády 1965
<i>Hieracium phaedrocheilon</i> Zahn ²²⁶		Zahn 1938; Nyárády 1965
<i>Hieracium praebiharicum</i> Boros ²²⁷		Boros 1972
<i>Hieracium prodanianum</i> Nyár. et Zahn ²²⁸		Zahn 1938; Nyárády 1965
<i>Hieracium pseudocaesiiforme</i> Nyár. et Zahn ²²⁹		Zahn 1938; Nyárády 1965
<i>Hieracium pseudocaesium</i> Degen et Zahn ²³⁰		Zahn 1938; Nyárády 1965
<i>Hieracium pseudonigrum</i> Pax ²³¹		Zahn 1938; Nyárády 1965
<i>Hieracium pseudopaltinae</i> Nyár. et Zahn ²³²		Zahn 1938; Nyárády 1965
<i>Hieracium pseudoratezatense</i> Nyár. et Zahn ²³³		Zahn 1938; Nyárády 1965
<i>Hieracium stenodontophyllum</i> Nyár. et Zahn ²³⁴		Zahn 1936; Nyárády 1965
<i>Hieracium trischistum</i> Nyár. et Zahn ²³⁵		Zahn 1938; Nyárády 1965
<i>Hieracium vurtopicum</i> Zahn ²³⁶		Zahn 1938; Nyárády 1965
<i>Hieracium wichurae</i> Zahn ²³⁷		Zahn 1936; Nyárády 1965

Taxon	Synonyms	References
<i>Knautia dipsacifolia</i> subsp. <i>lancifolia</i> (Heuff.) Ehrend. ²³⁸	<i>Knautia dipsacifolia</i> var. <i>lancifolia</i> (Heuff.) Simonk.; <i>K. sylvatica</i> subsp. <i>lancifolia</i> (Heuff.) Soó	Ehrendorfer 1976; Oprea 2005; Papović et al. 2014; Štěpánek 2014 in litt.
<i>Knautia dipsacifolia</i> subsp. <i>pocutica</i> (Szabó) Ehrend. ²³⁹	<i>Knautia dipsacifolia</i> var. <i>pocutica</i> Szabó	Ehrendorfer 1976; Oprea 2005; Štěpánek 2014 in litt.
<i>Knautia dipsacifolia</i> subsp. <i>turocensis</i> (Borbás) Jáv. ex Kiss ²⁴⁰	<i>Knautia turocensis</i> (Borbás) Szabó	Ehrendorfer 1976; Kliment 1999
<i>Leucanthemum raciborskii</i> Popov et Chrshan. ²⁴¹		Cvelev 1961, 1994; Čopyk 1976; Heywood 1976; Zelený 1982; Ziman et al. 2006
<i>Leucojum vernum</i> var. <i>carpathicum</i> Sweet ²⁴²	<i>Leucojum vernum</i> subsp. <i>carpathicum</i> (Sweet) K. Richt.	Soják 1962; Kliment 1999; Bělohávková 2010
<i>Minuartia cataractarum</i> (Janka) Nyár. ²⁴³	<i>Alsine cataractarum</i> Janka; <i>Minuartia frutescens</i> subsp. <i>cataractarum</i> (Janka) Soó; <i>M. hirsuta</i> subsp. <i>cataractarum</i> (Janka) Soó	Jalas & Suominen 1983; Halliday 1993; Dihoru & Negrean 2009; Sârbu et al. 2013
<i>Myosotis transsylvanica</i> Porcius ²⁴⁴		Grințescu & Nyárády 1960b; Grau 1964; Grau & Merxmüller 1972; Walter & Gillett 1998
<i>Poa molinerii</i> subsp. <i>glacialis</i> Beldie ²⁴⁵		Beldie 1967b
<i>Poa nyaradyana</i> Nannf. ²⁴⁶	<i>Poa laxa</i> subsp. <i>pruinosa</i> Nyár.; <i>P. laxa</i> var. <i>caesioglauca</i> Schur ex Nyár.	Nyárády 1928; Șerbănescu 1967; Ghișa & Beldie 1972; Edmondson 1980; Ciocârlan 2009
<i>Potentilla chrysantha</i> subsp. <i>pastorum</i> Soják ²⁴⁷		Soják 1993
<i>Primula elatior</i> subsp. <i>carpathica</i> (Griseb. et Schenk) W. W. Sm. et Forrest. ²⁴⁸	<i>Primula carpathica</i> (Griseb. et Schenk) Fuss	Schwarz 1968; Valentine & Kress 1972
<i>Primula elatior</i> subsp. <i>poloninensis</i> (Domin) Dostál ²⁴⁹	<i>Primula elatior</i> f. <i>poloninensis</i> Domin; <i>P. poloninensis</i> (Domin) Fed.	Morariu et al. 1960; Valentine & Kress 1972; Prokudin 1987; Ziman et al. 2006
<i>Primula matthioli</i> subsp. <i>pubens</i> (Schott, Nyman et Kotschy) Kovt. ²⁵⁰	<i>Cortusa matthioli</i> subsp. <i>pubens</i> (Schott, Nyman et Kotschy) Jáv.	Morariu & Nyárády 1960; Kobiv 1999, 2009, 2010, 2012a; Kovtonjuk 2013
<i>Pulmonaria montana</i> subsp. <i>porciusii</i> Gusul. ²⁵¹		Gușuleac 1960; Merxmüller & Sauer 1972; Ciocârlan 2009; Valdés in Euro+Med
<i>Ranunculus binatus</i> Kit. ex Rchb. ²⁵²	<i>Ranunculus auricomus</i> subsp. <i>binatus</i> (Kit. ex Rchb.) Jasiewicz	Dunkel 2011
<i>Rubus moestus</i> Holuby ²⁵³		Holuby 1873; Nyárády 1956; Kurtto in Euro+Med
<i>Rumex carpaticus</i> (Zapał.) Zapał. ²⁵⁴	<i>Rumex arifolius</i> subsp. <i>carpaticus</i> (Zapał.) Pawł.; <i>Acetosa alpestris</i> subsp. <i>carpatica</i> (Zapał.) Dostál	Borodina 1979; Cvelev 1996; Grabovskaja-Borodina 2012; Oprea & Sîrbu 2013
<i>Scabiosa lucida</i> subsp. <i>calcicola</i> Błoński ²⁵⁵	<i>Scabiosa calcarea</i> Tocl	Chrtek 1985a, b; Štěpánek & Holub 1997; Kliment 1999
<i>Scabiosa lucida</i> subsp. <i>pseudobanatica</i> (Schur) Holub ²⁵⁵	<i>Scabiosa lucida</i> subsp. <i>pseudobanatica</i> (Schur) Chrtek, nom. illeg.; <i>S. columbaria</i> subsp. <i>pseudobanatica</i> (Schur) Jáv.; <i>S. pseudobanatica</i> (Schur) Chrtek subsp. <i>pseudobanatica</i>	Chrtek 1985a, b; Kliment 1999; Oprea 2005; Bucalo et al. 2006; Kricsfalusy & Budnikov 2007
<i>Soldanella hungarica</i> group ²⁵⁶ (<i>S. angusta</i> Zhang; <i>S. hungarica</i> Simonk.; <i>S. major</i> (Neilr.) Vierh.; <i>S. marmarossiensis</i> Klášť.; <i>S. oreodoxa</i> Zhang; <i>S. pseudomontana</i> F. K. Mey.; <i>S. rugosa</i> Zhang; <i>S. stiriaca</i> F. K. Mey.; <i>S. tatricola</i> Niederle)	<i>Soldanella hungarica</i> subsp. <i>major</i> (Neilr.) Pawłowska	Klásterský 1930; Pawłowska 1963; Kress 1984; Meyer 1985; Hrouda & Kochjarová 1997; Zhang et al. 2001; Zhang & Kadereit 2002, 2004; Niederle 2003, 2005; Fischer et al. 2008
<i>Sorbus paxiana</i> Jáv. ²⁵⁷		Jávorka 1927; Kárpáti 1960
<i>Trisetum alpestre</i> subsp. <i>glabrescens</i> (Schur) Tzvelev ²⁵⁸	<i>Trisetum alpestre</i> var. <i>glabrescens</i> Schur	Schur 1866; Cvelev 1974, 1976; Prokudin et al. 1977

Notes

²⁰⁰ Starmühler (1998a) re-classified the species *Aconitum simonkaianum* to the subspecies level (*Aconitum moldavicum* subsp. *simonkaianum* (Gáyer) Starm.) and reported it from the Ukrainian and Romanian Eastern Carpathians (cf. Starmühler 1998b, Mitka 2008, Mitka & Novikoff 2011, Novikoff & Mitka 2011a, b, Novikov 2013b). However, already Grințescu (1953) considered this taxon to be a hybrid, thus he reported it as *Aconitum* × *simonkaianum* (Gáyer) Grinț. (*A. hosteanum* × *A. velutinum*) [*Aconitum velutinum* = *A. lasiostomum*]. The hybrid formula *Aconitum lasiostomum* × *A. moldavicum* was published after the subspecies name also by Mitka (2008; cf. Mitka & Gawroński 2010). According to him, the hybrid status of this taxon should be confirmed by further biochemical and molecular analyses on representative material including the whole area of *Aconitum moldavicum*. Subspecies *A. moldavicum* subsp. *simonkaianum* is evaluated as nothotaxon (*A. moldavicum* subsp. *hosteanum* × *A. lasiostomum* subsp. *lasiostomum*) also by Novikov (2014 in litt.).

²⁰¹ *Allium zahariadi* subsp. *zahariadi* is a subspecies with a complicated taxonomic history. It was described as *Allium ochroleucum* subsp. *pseudosuaveolens* from fen meadows between the Vlăhița Village and Tolvaioș Gorge in the Eastern Carpathians by Zahariadi (1966), who recorded it also in the vicinity of the nearby village of Căpilnița. Based on his description, Májovský (in Májovský & Murín 1985) described a narrowly perceived species of meadows *Allium zahariadi* with subspecies *A. zahariadi* subsp. *zahariadi* and *A. zahariadi* subsp. *michalkoi* (see below). According to at that time valid International code of botanical nomenclature (Greuter et al. 1994), Somogyi (1999a) considered the description of subsp. *pseudosuaveolens*, as well as of other derived taxa, to be invalid (due to the absence of a type specimen in the description). He ordered it to a broadly perceived species *Allium ericetorum* Thore, involving the populations of meadow communities. The subsp. *pseudosuaveolens* was reclassified to the last mentioned taxon also by Ciocârlan (2009), who reported *A. ericetorum* subsp. *pseudosuaveolens* as a narrow endemic known from only two localities mentioned in the original diagnosis (cf. Sârbu et al. 2013). Although according to Art. 40.3 of the recent code version (McNeill et al. 2012) the description of *Allium ochroleucum* subsp. *pseudosuaveolens* can be considered as valid (Marhold 2014 in litt., Turland 2014 in litt.), the endemic status of this taxon remains unclear until the detailed knowledge on the intraspecific variability of the *Allium ericetorum* group is known within its whole distribution area.

²⁰² A tall white-flowering meadows subspecies *Allium zahariadi* subsp. *michalkoi* was described from the vicinity of the Șaca Village in the Košická kotlina Basin (Májovský & Murín 1985), from where it is recently known in a single microlocality (Feráková & Somogyi 1999, Somogyi 1999b, 2002). Its evaluation as a narrow endemic to the Western Carpathians (Tasenkevich 2002) is left open by now (see the note above).

²⁰³ The subspecies *Bupleurum falcatum* subsp. *dilatatum* includes robust, up to 2 m high branched plants, with stem leaves up to 20 (25) cm long and 3–6.5 cm broad (Šourková 1984) and tetraploid chromosome number ($2n = 32$). According to Tutin (1968), it is not clear whether there is a correlation between the morphometric characters and chromosome number; the subspecies is mentioned only in a note. The precise distribution of this subspecies is also not known up to now, it is reported from the Western (Slovakia, Hungary), Eastern and Southern Carpathians (Romania) and the Transylvanian Basin (Kliment 1999, Oprea 2005) and it was evaluated as a pan-Carpathian subendemic (Kliment 1999, Šeffler et al. 2010).

²⁰⁴ Nyárady (1965) considered *Carduus fissurae* to be a hybridogenous species (*C. acanthoides* × *C. crispus*) and a local endemic to the Cheile Turzii Gorge (Trascău Mts) in the Apuseni Carpathians, where it was described (cf. Pawłowski 1970, Heltmann 1985, Oprea 2005). As a separate species it was accepted also by Greuter (in Euro+Med). Contrastingly, Beldie & Alexandrescu (1976), Morariu & Beldie (1976) and Ciocârlan (2009) identified it with the hybrid *Carduus* × *leptocephalus* Peterm. (*C. acanthoides* × *C. crispus*). Some supranational databases (The Plant List, Catalogue of Life) report *C. fissurae* as a synonym of *Carduus crispus* L.

²⁰⁵ *Carex sempervirens* subsp. *tatrorum*, a narrow-leaved calciphilous subspecies of taxonomically complicated species *Carex sempervirens* Vill., is considered to be endemic to the Western Carpathians (cf. Kliment 1999, Piękoś-Mirkowa & Mirek 2003, Mirek & Piękoś-Mirkowa 2009, 2010). However, its taxonomic evaluation is not consistent. Marhold et al. (2007) reported it as a synonym of the West-East-South-Carpathian calcicolous subspecies *Carex sempervirens* subsp. *laxiflora* (Schur) Jáv.; Luceño et al. (2008) as a synonym of a calciphilous subspecies *C. sempervirens* subsp. *sempervirens*, distributed on alkaline substrates from the mountain ranges of the Iberian Peninsula through the Alps, Jura, Apennines and Carpathians to mountain ranges of the Balkan Peninsula (cf. Marhold 1998, Jiménez-Mejías & Luceño in Euro+Med). Some supranational databases (The Plant List, Catalogue of Life) accept it only at the species level (cf. Ciocârlan 2009, Sârbu et al. 2013).

²⁰⁶ The subspecies *Carex sempervirens* subsp. *pseudotristis* was described from the Ukrainian Carpathians (Domin 1931b ut *Carex sempervirens* var. *pseudotristis*) and sporadically considered (Tasenkevych 2003b) to be endemic to the Eastern Carpathians. Luceño et al. (2008) used the name *C. sempervirens* subsp. *pseudotristis* for both the Pyrenean and the

Carpathian silicicolous populations of *Carex sempervirens*, meantime reported as *Carex granitica* Braun-Blanq. or *Carex sempervirens* subsp. *silicicola* Holub (cf. Jiménez-Mejías & Luceño in Euro+Med).

²⁰⁷ *Centaurea globurensis* is a species with unclear taxonomic status, which was described from Mt. Arjana near the Globurău Village (Cerna Mts, the Southern Carpathians) and included within the groups *Centaurea atropurpurea* Waldst. et Kit. (Dostál 1976), *C. calocephala* Willd. (Greuter in Euro+Med) and also *C. scabiosa* L. (Dihoru & Pârvu 1987).

²⁰⁸ *Centaurea pugioniformis* is a narrowly-perceived species belonging to the group of *Centaurea macroptilon* Borbás, from which it differs only by narrower appendices not covering involucral bracts (Dostál 1976). It was reported from the Eastern, Southern and Apuseni Carpathians, the Transylvanian Basin, sporadically also from the localities outside the Carpathians – Timișoara, Iași (Prodan & Nyárády 1964, Oprea 2005). Niketić (2010) reported its occurrence also from the Serbian part of the Southern Carpathians (Đerdap: Lepenski vir, Kazan). According to Hurdu (2014 in litt.) and Vonica (2014 in litt.), it concerns a hybrid forms of the subspecies *Centaurea macroptilon* subsp. *oxylepis* (Wimmer et Grab.) Soó (cf. Ciocârlan 2009).

²⁰⁹ *Dactylorhiza carpatica* is a taxonomically unclear species originated by hybridization of *Dactylorhiza* species and known only from the calcareous fen on the Moravian side of the Bílé Karpaty Mts (Kaplan 2012). It was included neither in the current checklist of vascular plants of the Czech flora (Danihelka et al. 2012) nor in the newest Czech Red list (Grulich 2012). Kubát (2010) reported it in a note to *Dactylorhiza fuchsii* (Druce) Soó with a comment that it probably concerns a taxon of hybridogenous origin, requiring further study. Mrázek (<http://botany.cz/cs/dactylorhiza-traunsteineri-carpatica>) considers it to be a hybridogenous species, stenoendemic to the Moravian Bílé Karpaty Mts.

²¹⁰ Fedorončuk (2009) evaluates *Dianthus carpaticus* as a local endemic to the Eastern Carpathians occurring in the subalpine and alpine belts of the Ukrainian Carpathians, well differentiated from the relative species *Dianthus carthusianorum* L. and *D. tenuifolius* Schur (cf. Fedorončuk & Diduch 2002a, Fedorončuk & Čornej 2005, Čornej 2011, Tasenkevich 2011). By a set of its morphological characters it represents a transitional position between these two species (Fedorončuk & Čornej l. c.). Contrastingly, Tutin (1964), Jalas & Suominen (1986) and Kuz'mina (2004) consider it to be a synonym of *Dianthus carthusianorum* L., Malynovs'kyj et al. (2002) as a synonym of *D. carthusianorum* subsp. *subalpinus* (Rehmann) Májovský et Králik. Reversely, Klokov (1952) reported *D. carthusianorum* var. *subalpinus* Rehmann as synonym of *Dianthus carpaticus*.

²¹¹ As a synonym of *Dianthus carpaticus* Fedorončuk & Diduch (2002a) reported also *Dianthus bucovinensis* (Zapał.) Klokov, considered by Klokov (1952) as endemic to the Bukovinské Karpaty Mts; contrastingly, Jalas & Suominen (1986) considered it as synonym of *Dianthus carthusianorum*.

²¹² Zahn (1936) reported *Hieracium amoenanthes* only from Mt. Retezat (Retezat Mts), 1 800–1 900 m a.s.l. in the Southern Carpathians (cf. Nyárády 1965). It is probably endemic to the Retezat Mts; however, the current taxonomic and chorological knowledge is missing. Heltmann (1985) and Dihoru & Pârvu (1987) considered it to be endemic to the Southern Carpathians.

²¹³ Zahn (1938) reported *Hieracium borzae* from Mt. Retezat (cf. Nyárády 1965). It is probably endemic to the Retezat Mts; however, the current taxonomic and chorological knowledge is missing. Heltmann (1985) and Dihoru & Pârvu (1987) considered it to be endemic to the Southern Carpathians.

²¹⁴ Zahn (1938) reported the occurrence of *Hieracium breazense* in the Răul Bescivarei Valley (Făgăraș Mts); Nyárády (1965) also in the locality Giurcuța de Sus (Apuseni Mts). The current taxonomic and chorological data on this species are not available. Heltmann (1985) considered it to be endemic to the Southern Carpathians, Dihoru & Pârvu (1987) evaluated it as endemic to the Southern and Apuseni Carpathians.

²¹⁵ *Hieracium caesiogenum* was described from Mt. Berdo near the Hryniava Village at the border of the Ukrainian and Romanian Carpathians. Zahn (1936) reported several localities from the Făgăraș Mts, Nyárády (1965) also from the Rodna Mts and Bihor-Vlădeasa Mts (cf. Oprea 2005). The Ukrainian authors evaluate it as endemic to the Eastern and Southern Carpathians in their surveys. The current data for delimitation of its real distribution are still missing; it is probably endemic to the Eastern and Southern Carpathians (cf. Zahn 1936).

²¹⁶ Zahn (1938) reported *Hieracium calcogeton* from the Bihar Mts (Apuseni Carpathians) using an invalid name *Hieracium biharianum* Prodan et Zahn (only a German diagnosis). Neither newer taxonomic nor chorological data are available (cf. Nyárády 1965). Heltmann (1985) as well as Dihoru & Pârvu (1987) ordered it as *Hieracium biharianum* to species endemic to the Apuseni Carpathians.

²¹⁷ According to Zahn (1936), *Hieracium chloribracteum* grows only in the Retezat Mts (the Southern Carpathians) at altitudes between 1 400 and 2 300 m. Nyárády (1965) considered it to be endemic to the Retezat Mts, Dihoru & Pârvu (1987) as endemic to the Southern Carpathians. The current taxonomic and chorological data for delimitation of its real distribution are missing.

²¹⁸ Zahn (1927, 1935) reported the occurrence of *Hieracium dentatum* subsp. *sarmaticum* from several mountain ranges in the Slovak part of the Western Carpathians, Šljakov (1989) from the Ukrainian Carpathians, Nyárády (1965) from the Apuseni Carpathians (Colții Trascău). Stoyko & Tasenkevich (1993) and Čornej (2011) evaluated it as endemic to the Western and Eastern Carpathians.

²¹⁹ According to Zahn (1938), *Hieracium grecescui* grows only in the Retezat Mts (cf. Nyárády 1965). Recent data on this species are missing. Heltmann (1985) and Dihoru & Pârvu (1987) considered it endemic to the Southern Carpathians.

²²⁰ *Hieracium klopotivae* was described from the Retezat Mts (Clopotiva in the Râului Mare Valley). Nyárády (1965) reported it also from the Maramureș Mts (Stâna lui Vartic). This record was adopted by Oprea (2005) and Sârbu et al. (2013), who *H. klopotivae* evaluated as endemic to Romania or to the Carpathians. According to Szeląg (2014 in litt.), *Hieracium klopotivae* s. str. grows probably only in the Retezat Mts.

²²¹ *Hieracium krizsnae* was described by Zahn (1927) from Mt. Krížna in the Veľká Fatra Mts (the Western Carpathians). Its occurrence in this locality has not been confirmed since that time, however, it was documented by Lengyel's original herbarium specimen in the collection of BP (Šípošová et al. 2004a). It was evaluated as endemic to the Veľká Fatra Mts by Mráz (in Eliáš jr. et al. 2015).

²²² Juksip (1959) described *Hieracium mukaczense* from an oak forest nearby the city Mukačevo. As endemic to the Eastern or the Ukrainian Carpathians it was evaluated e.g. by Stojko & Tasenkewitsch (1991), Stoyko & Tasenkevich (1993), Malynovs'kyj et al. (2002) and Tasenkevych (2003b). Up to now, it is known only from the type specimen and no newer data are available since its description (Sennikov 2014 in litt.). Also later floristic studies (e.g. Prokudin 1987, Šljakov 1989) contain only basic information on *locus classicus* of this narrowly-perceived species.

²²³ Zahn (1938) reported *Hieracium nyaradyanum* from the Făgăraș Mts and Țarcu Mts, Nyárády (1965) from the Făgăraș, Godeanu and Țarcu Mts. It is probably endemic to the Southern Carpathians (cf. Heltmann 1985, Dihoru & Pârvu 1987).

²²⁴ Zahn (1938) reported the occurrence of *Hieracium pelagae* from Mt. Peleaga (Retezat Mts) in the Southern Carpathians and the Țibleș Mts in the Eastern Carpathians. More recent data are not available (cf. Nyárády 1965). According to Heltmann (1985) and Dihoru & Pârvu (1987) it is endemic to the Eastern and Southern Carpathians.

²²⁵ Zahn (1938) reported *Hieracium peterfii* from two localities in the Retezat Mts (the Southern Carpathians). More recent data are not known (cf. Nyárády 1965). The Romanian authors (Heltmann 1985, Dihoru & Pârvu 1987) considered it endemic to the Southern Carpathians.

²²⁶ Zahn (1938) reported *Hieracium phaedrocheilon* from the localities Csikszentdomokos (Sândominic) and Marosfő (Izvoru Mureșului) in the Romanian Eastern Carpathians, Nyárády (1965) only mentioned its occurrence in the Southern Carpathians. More recent records are missing. Heltmann (1985) ordered it to species endemic to the Transylvanian Basin, Dihoru & Pârvu (1987) to species endemic to the Eastern Carpathians.

²²⁷ *Hieracium praebiharicum* was described from a stand of the *Seslerietum rigidae* in the vicinity of the Poșaga de Sus Village, 500 m a.s.l., in the Gilău Mts (Apuseni Carpathians, Boros 1972). Although it is considered endemic to the Apuseni Carpathians or to Romania, in the Romanian lists of endemic species as well as in the national floras (Morariu & Beldie 1976, Beldie 1979, Heltmann 1985, Dihoru & Pârvu 1987, Oprea 2005, Ciocârlan 2009, Sârbu et al. 2013), it is reported only from the *locus classicus*. Since its description, it has not been subject of a taxonomic study.

²²⁸ Zahn (1938) reported *Hieracium prodanianum* from the Retezat Mts (the Southern Carpathians) and the Bihar Mts (Apuseni Carpathians). More recent taxonomic and chorological data are not available (cf. Nyárády 1965). Heltmann (1985), Dihoru & Pârvu (1987) ordered it to species endemics to the Southern Carpathians.

²²⁹ Zahn (1938) reported *Hieracium pseudocaesiiforme* from the surroundings of the glacial lake Gemenea in the Retezat Mts; more recent data are not available (cf. Nyárády 1965). According to Heltmann (1985) and Dihoru & Pârvu (1987), it is endemic to the Southern Carpathians.

- ²³⁰ Zahn (1938) reported *Hieracium pseudocaesium* from several localities in the Retezat and Făgăraș Mts (cf. Nyárády 1965). More recent data are not available. Heltmann (1985) and Dihoru & Pârvu (1987) consider it endemic to the Southern Carpathians.
- ²³¹ *Hieracium pseudonigrum* was described and is recently known only from Mt. Barnarului in the Bistrița Mts (cf. Zahn 1938, Nyárády 1965). Heltmann (1985) ordered it to species endemic to the Eastern Carpathians.
- ²³² Zahn (1938) reported *Hieracium pseudopaltinae* from the Bârsei (Piatra Mare) and Retezat Mts; his data were adopted by Nyárády (1965). In studies of Heltmann (1985) and Dihoru & Pârvu (1987) the species is evaluated as endemic to the Southern Carpathians.
- ²³³ Zahn (1938) and Nyárády (1965) reported *Hieracium pseudoratezatense* from two localities in the Retezat Mts (the Southern Carpathians). More recent data are not available. Heltmann (1985) and Dihoru & Pârvu (1987) considered it endemic to the Southern Carpathians.
- ²³⁴ *Hieracium stenodontophyllum* was described from alpine surroundings of the glacial lake of Gemenea (Retezat Mts) in the Southern Carpathians (cf. Zahn 1936, Nyárády 1965); more recent data on its chorology are not available. Heltmann (1985) and Dihoru & Pârvu (1987) considered it endemic to the Southern Carpathians.
- ²³⁵ Zahn (1938) reported *Hieracium trischistum* from the Retezat Mts (the Southern Carpathians); Nyárády (1965) reported it as var. *dealunegri* Nyár. et Zahn also from the Godeanu Mts. According to Heltmann (1985) and Dihoru & Pârvu (1987), it is endemic to the Southern Carpathians.
- ²³⁶ *Hieracium vurtopicum* was described and is recently known only from Mt. Vîrtopu (Făgăraș Mts) in the Southern Carpathians (cf. Zahn 1938, Nyárády 1965). According to Heltmann (1985) and Dihoru & Pârvu (1987), it is endemic to the Southern Carpathians.
- ²³⁷ Zahn (1936) reported *Hieracium wichurae* from alpine regions of the Piatra Mare Massif (Bârsei Mts) in the Southern Carpathians; more recent data are missing also in the study of Nyárády (1965). According to Heltmann (1985) and Dihoru & Pârvu (1987), it is endemic to the Southern Carpathians.
- ²³⁸ The subspecies *Knautia dipsacifolia* subsp. *lancifolia* is accepted in Flora Europaea (Ehrendorfer 1976), supranational databases (The Plant List, Catalogue of Life, wikipedia, and others) as well as in more recent taxonomic publications (Oprea 2005, Ciocârlan 2009, Sârbu et al. 2013). In the included surveys of endemic species it is reported as endemic to the South-Eastern Carpathians or subendemic to the Carpathians. Besides the Romanian Eastern, Southern and Apuseni Carpathians (cf. Oprea 2005) it has been recently recorded also in the south-eastern Serbia (Papović et al. 2014). According to Štěpánek (2014 in litt.), individual subspecies were distinguished based on habitat-dependent characters; more recent taxonomic data on variability of *Knautia maxima* (Opiz) Ortmann (valid species name) are still missing. Regarding unclear taxonomic status and chorological data, individual subspecies were not considered in current lists of species endemic to the South-Eastern Carpathians (Hurdu et al. 2012a, b).
- ²³⁹ Similarly to previous subspecies, *Knautia dipsacifolia* subsp. *pocutica* is also accepted by Ehrendorfer (1976) and other above-mentioned studies. It is reported from the Slovak (Bukovské vrchy Mts) and Polish (Bieszczady Mts) Carpathians, Ukrainian as well as Romanian part of the Eastern Carpathians, and currently (e.g. Oprea 2005, Sârbu et al. 2013) also from the Apuseni Carpathians (Bihor-Vlădeasa Mts). In the included surveys of endemic species it is reported as endemic to the Eastern Carpathians, endemic to the Western and Eastern Carpathians or endemic to the Eastern, Southern and Apuseni Carpathians. Popescu & Sanda (1998) report it as a synonym of subsp. *lancifolia*.
- ²⁴⁰ The distribution of subspecies *Knautia dipsacifolia* subsp. *turocensis* varies in different published sources from the Western Carpathians (including the Északi-középhegység Mts) through the Ukrainian (Mt. Pikuj) and Romanian Eastern Carpathians (Rodna Mts), Southern Carpathians up to the Stara planina Mts (Mt. Midzor) in Bulgaria (see Kliment 1999 for details). Ehrendorfer (1976) delimited its distribution area to the Western Carpathians.
- ²⁴¹ The species *Leucanthemum raciborskii* was described from the Svydovec Mts and in several reviews of endemic species it is evaluated as endemic to the Eastern (Ukrainian) Carpathians (Čopyk 1976, Tasenkevich 2011, CBIS 2008: <http://www.carpates.org/cbisec/bot.php?id=6015>). Based on original herbarium specimens collected by the authors of species diagnosis, Ziman et al. (2006) evaluated *L. raciborskii* as an alpine endemic vicariant of *Leucanthemum vulgare* (1 600–1 800 m a.s.l., CBIS 2008), from which it differs by several morphological characters. Cvelev (1961, 1994) ordered it as a synonym of *Leucanthemum subalpinum* (Schur) Tzvelev described from the Southern Carpathians, which in this concept is considered endemic to the Eastern and Southern Carpathians (Kricsfalussy 1999, Antosyak & Kozurak 2011). Both species were identified also by Čopyk (1976), who estimated a diploid chromosome number ($2n = 18$) for *L. raciborskii*. Contrastingly, Heywood (1976) considered *Leucanthemum raciborskii* to be synonym of a broadly-percieved species

Leucanthemum vulgare Lam. Zelený (1982, 2015 in litt.) ordered *L. raciborskii* as well as *L. subalpinum* to synonyms of the Alpine-Carpathian alpine diploid subspecies *Leucanthemum vulgare* subsp. *alpicola* Á. Löve et D. Löve. All three taxa were evaluated as synonyms of *Leucanthemum gaudinii* Dalla Torre (cf. The Plant List, Catalogue of Life) by Greuter (in Euro+Med).

²⁴² *Leucojum vernal* var. *carpathicum* is a taxon with unclear taxonomy, nomenclature (different authors citations) and distribution, reported mainly as a subspecies *Leucojum vernal* subsp. *carpathicum* (Sweet) K. Richt. Soják (1962) reports it from the Vihorlat Mts in the Eastern Slovakia mentioning also its basic morphological characters (conspicuously larger flowers with yellow spots, often more robust growth and increased number of individuals with two flowers in the populations). He considered it to be geographically and morphologically not very well-defined, in its distribution restricted to the Eastern Carpathians and adjacent regions (Ukraine, Transylvania). He emphasized that the morphological criterium should not be overestimated as two-flowered individuals erroneously determined as *Leucojum carpathicum* are scattered in other regions of the species distribution (the Sudeten, the Alps) and vice versa, single-flowered individuals from Transylvania were wrongly ordered to nominate subspecies. Besides the Eastern Slovakia the taxon is reported, mainly in the older literature, also from Poland (Bieszczady Mts), Ukraine (the Carpathians, Opillja, Pravoberežnyj Lisostep) as well as from several localities at the foothill of Transylvanian mountain ranges in Romania (cf. Kliment 1999). With regard to its distribution, it was evaluated as subendemic to Eastern Carpathians in several overviews. A scattered occurrence of individuals corresponding in the abovementioned characters to subsp. *carpathicum* was recently reported by Bělohávková (2010) from the Czech Republic (Ještědský hřbet), Poland, Bavaria and the Alps with a note that the taxon requires thorough taxonomic and phytogeographical analysis within the whole area of the species distribution (cf. Dostál et al. 1999).

²⁴³ *Minuartia cataractarum* is a species with unclear taxonomic evaluation. In the Romanian surveys of endemic species as well as in taxonomic studies (e.g. Dihoru & Pârvu 1987, Negrean & Oltean 1989, Oprea 2005, Ciocârlan 2009, Hurdu et al. 2012a, b) it is evaluated as endemic to the Southern Carpathians, known only from the Iron Gate Gorge and the southern part of the Mehedinți Mts (cf. Dihoru & Negrean 2009). In other taxonomic studies (Jalas & Suominen 1983, Halliday 1993, Sârbu et al. 2013) it is included within the species *Minuartia frutescens* (Kit. ex Schult.) Tuzson ex Degen. Beldie & Alexandrescu (1976), and also Morariu & Beldie (1976) order it to synonyms of *Minuartia hirsuta* subsp. *falcata* (Griseb.) Mattf., which, however, grows only in the Balkan according to Atlas Florae Europaeae (Jalas & Suominen 1983).

²⁴⁴ *Myosotis transsylvanica* was described and is known only from the Rodna Mts in the Eastern Carpathians (cf. Grințescu & Nyárády 1960b ut *Myosotis variabilis* f. *transsylvanica* (Porcius) Jáv.). Grau (1964) identified it with the subspecies *Myosotis decumbens* subsp. *variabilis* (Angelis) Grau. However, she noted that plants from Transylvania, reported as *M. transsylvanica*, differ from this subspecies by the absence of long calyx hairs and after inspection of sufficient material it can be evaluated as a separate subspecies (cf. Grau & Merxmüller 1972). *Myosotis transsylvanica* was evaluated as a separate subspecies by Walter & Gillett (1998); preliminarily it was accepted also by Valdés (in Euro+Med).

²⁴⁵ The subspecies *Poa molinerii* subsp. *glacialis* was described by Beldie (1967b) from alpine regions of the Bucegi Mts (the Southern Carpathians), at about 2 020–2 350 m a.s.l. From the nominate subspecies it differs by a significantly shorter stalk (3–12 cm), shorter and more compact inflorescence (0.8–2 cm) and only 2–3-flowered spikelets (Ciocârlan 2009). It is possibly endemic to the Bucegi Mts (cf. Beldie 1979, Oprea 2005, Ciocârlan 2009, Sârbu et al. 2013); however, the current floristic data from this mountain range as well as the knowledge on its overall distribution are missing (Hurdu 2014 in litt.).

²⁴⁶ *Poa nyaradyana* is a species with an unclear taxonomic status and ambiguous distribution. It was described by Nyárády (1928, ut *Poa laxa* subsp. *pruinosa*) from the Făgăraș Mts (the Southern Carpathians). Ghișa & Beldie (1972, ut *Poa laxa* var. *caesioglauca*) reported it also from other mountain ranges of the Southern Carpathians, Oprea (2005) and Sârbu et al. (2013) also from some mountain ranges of the Romanian Eastern Carpathians. The morphometric comparison of *Poa laxa* Haenke and *Poa nyaradyana* (Șerbănescu 1967) detected significant overlap in all measured characters except the average of longer anthers in *Poa nyaradyana*. Therefore, Șerbănescu (l. c.) considered *Poa nyaradyana* to be an ecotype of *Poa laxa* not exceeding the variability of this species (cf. Edmondson 1980, Ciocârlan 2009); Tassenkevych (2010) considered it to be an ecological (calciphilous) subspecies of *Poa laxa*.

In the current Romanian taxonomic handbooks (Popescu & Sanda 1998, Ciocârlan 2009, Sârbu et al. 2013), *Poa tremula* Schur (non Lam., nom. illeg.) is also reported as synonym of *Poa laxa*. This species, in most surveys evaluated as endemic to the Southern Carpathians, was reported by Schur (1866) also from Mt. Ineu (Rodna Mts) in the Eastern Carpathians. Șerbănescu (l. c.) ordered populations from Mt. Ineu to *Poa laxa*. Tassenkevych (2010) considers *Poa tremula* as synonym of *Poa laxa* subsp. *pruinosa* (i.e. *Poa nyaradyana*).

²⁴⁷ Soják (1993) described the subspecies *Potentilla chrysantha* subsp. *pastorum* from the Bucegi Mts in the Southern Carpathians (subalpine meadows in the vicinity of Valea Jepilor below the top of Mt. Caraiman, above the Busteni Village); however, he has not mentioned its overall distribution. In the current Romanian taxonomic handbooks (Oprea 2005, Ciocârlan 2009, Sârbu et al. 2013) only the species is mentioned.

²⁴⁸ *Primula elatior* subsp. *carpathica* is a subspecies with an unclear taxonomic status and distribution. Schwarz (1968) delimited its distribution only to the Carpathian mountain ranges. Its relation to other subspecies reported as endemic from individual Carpathian subunits, *Primula elatior* subsp. *poloninensis* (Domin) Dostál (see below) and *P. elatior* subsp. *tatrensis* (Domin) Soó, is not clear. At the level of subspecies or species of *Primula carpathica* (Griseb. et Schenk) Fuss, it is accepted also by some supranational databases (The Plant List, Tropicos, Catalogue of Life). On the other hand, Valentine & Kress (1972) evaluated it as synonym of subsp. *elatior* (cf. Marhold in Euro+Med). The variability of *Primula elatior* (L.) Hill. requires a thorough molecular-taxonomic study.

²⁴⁹ Similarly to subsp. *carpathica*, also *Primula elatior* subsp. *poloninensis* is a subspecies with problematic taxonomic status and insufficiently known distribution. More recent chorological data are available from the Polish (Bieszczady Mts) and Slovak (Bukovské vrchy Mts) Carpathians, and Ukrainian part of the Eastern Carpathians. From the Romanian Carpathians only older data exist from the Bârsei Mts (Postăvarul Massif) and Piatra Craiului Mts in the Southern Carpathians (Morariu et al. 1960, ut *Primula elatior* f. *poloninensis*); in more recent Romanian taxonomic handbooks this subspecies is not distinguished. In most overviews of endemic species it is reported as endemic to the South-Eastern or, more sparsely, to the Eastern Carpathians. Ambiguous is also its taxonomic evaluation – from the level of a species through the one of a subspecies, variety, form to classification as a synonym of *Primula elatior* subsp. *elatior* (Valentine & Kress 1972).

²⁵⁰ *Primula matthioli* subsp. *pubens* is a taxonomically and chorologically unclear subspecies, up to now known mainly as *Cortusa matthioli* subsp. *pubens*. Most of the published studies from the Carpathian region contains only data on the species *Cortusa matthioli*, therefore the precise distribution of the subspecies is not known. More recent actual data on its distribution are available from the Ukrainian Carpathians. Kobiv (1999, 2009, 2010, 2012a) found an abundant population on the east-facing slopes of Mt. Hoverla (Čornohora Mts, the Eastern Carpathians), at about 1 630–1 650 m a.s.l. Only older records are available from the Romanian Carpathians (Morariu & Nyárády 1960), namely from the Rodna (*locus classicus*: Mt. Corongiș) and the Făgăraș Mts (Valea Bîlii; ut f. *subpubens* Nyár.). Probably with regard to the last-mentioned record, *Cortusa pubens* is in some overviews evaluated as endemic to the Eastern and Southern Carpathians (Čopyk 1976, Kobiv 1999, Tassenkevich 2011). Contrastingly, Kovtonjuk (2013), already within the genus *Primula* L., where the genus *Cortusa* was reclassified according to the results of current molecular-phylogenetic studies (Mast et al. 2001, Yan et al. 2010), evaluates it as endemic to the Eastern Carpathians (cf. Kobiv 2012a). Within the distribution area of *Primula matthioli* (L.) V. A. Richt., Kovtonjuk (l. c.) distinguished together 11 subspecies including subsp. *pubens*. As a morphologically and to some extent also ecologically separate subspecies, *Cortusa matthioli* subsp. *pubens* was evaluated also by Kobiv (1999, 2010).

²⁵¹ The subspecies *Pulmonaria montana* subsp. *porciusii* was described from the subalpine belt of the Rodna Mts (Gușuleac 1960); later it was evaluated as endemic to this mountain range (Ciocârlan 2009) or as endemic to the Eastern Carpathians (Sârbu et al. 2013). Opinions to its taxonomic evaluation differ: Merxmüller & Sauer (1972) reported it in a note to *Pulmonaria mollis* Wulfen ex Hornem. (see also The Plant List); Valdés (in Euro+Med) considers it as synonym of *Pulmonaria dacica* (Simonk.) Simonk.

²⁵² *Ranunculus binatus* represents a microspecies within the *Ranunculus auricomus* group (Dunkel 2011) with insufficiently known distribution. Based on the original diagnosis and additional typification, it surely grows in Transylvania (Romania) and in the Slovak part of the Western Carpathians (cf. Dunkel l. c.). It is reported as endemic to the Western, Eastern & Southern Carpathians and Transylvanian Basin by Tassenkevych (2003b).

²⁵³ *Rubus moestus* was described from the vicinity of the Zemianske Podhradie Village (Biele Karpaty Mts) in the Western Slovakia (Holuby 1873). Nyárády (1956) reported its occurrence from two localities (Nicolae Bălcescu, Orșova) in the western Romania (cf. Ciocârlan 2009). Tassenkevich (2011, 2014) considers it as endemic to the Western Carpathians and Transylvanian Basin. According to Kurtto (in Euro+Med), the Romanian data are erroneous. More accurate distribution of *Rubus moestus* is still not known.

²⁵⁴ *Rumex carpaticus* (Zapał.) Zapał. was described from the Ukrainian Carpathians (Čornohora Mts). It is reported also from the Polish (Bieszczady Mts) and Slovak (Bukovské vrchy Mts) parts of the Eastern Carpathians, Romanian Eastern and Southern Carpathians. Usually it is considered endemic to the Eastern and Southern Carpathians, less frequently also endemic to the Eastern Carpathians. Opinions to its taxonomic status differ. Beside the separate position at the subspecies level (see the synonyms), it is usually ordered as a synonym of *Rumex alpestris* Jacq. (e.g. Oprea & Sîrbu 2013, The Plant List, Goliašová 2014 in litt.), or *Rumex rugosus* Campd. (Borodina 1979, Cvelev 1996, Catalogue of Life). Recently, Grabovskaja-Borodina (2012) included *Rumex rugosus* as well as *R. carpaticus* (Zapał.) Zapał. as synonyms of the subspecies *Rumex arifolius* subsp. *amplexicaulis* (Lapeyr.) Nyman. On the other hand, several older sources (e.g. Jalas & Suominen 1979, Rechinger & Akeroyd 1993, Oprea 2005) report *Rumex carpaticus* Rech. f. as a synonym of *R. alpestris*, but no more accurate data are available on this taxon.

²⁵⁵ For a long time, in overviews of the Carpathian endemic species only subspecies *Scabiosa lucida* subsp. *pseudobanatica* has been reported, evaluated usually as a West-East-South-Carpathian or pan-Carpathian endemic. Chrtek (1985a) noticed the existence of the subspecies *Scabiosa lucida* subsp. *calcicola*, distributed in the promontories and at lower altitudes of the Slovak and Moravian parts of the Western Carpathians (cf. Chrtek 1985b ut *S. lucida* subsp. *pseudobanatica*; Štěpánek & Holub 1997), which is probably endemic to the Western Carpathians (Štěpánek & Holub l. c., Kliment 1999). According to Chrtek (1985a), plants identical in all characters with the Slovak ones reach towards the east up to western part of the Ukrainian Carpathians (Mt. Pikuj), towards the south up to the Északi-középhegység Mts (Chrtek sec. Kliment 1999). *Scabiosa lucida* subsp. *pseudobanatica* is further reported from the Ukrainian and Romanian Carpathians (including the Apuseni Carpathians and Transylvanian Basin; cf. Oprea 2005), the accurate distribution of both subspecies is not known. Bucalo et al. (2006) reported the occurrence of *S. lucida* subsp. *pseudobanatica* also from the National Park Kozara (Bosnia and Hercegovina).

²⁵⁶ *Soldanella hungarica* group represents a complicated taxonomic complex with an ambiguous evaluation of its inner classification, on which several considerably distinct concepts were published:

Pawłowska (1963) distinguished within the complex the species *Soldanella montana* Mikan (the Austrian Alps; Český masív; the Western and Eastern Carpathians) and *Soldanella hungarica* Simonk. with two subspecies, subsp. *major* (Neilr.) Pawłowska (the Western, Eastern and Southern Carpathians; Rila, Rodopi; the Austrian Alps) and subsp. *hungarica* (the Eastern and Southern Carpathians). The Carpathian populations, reported as *S. montana*, were later included to new-described *Soldanella pseudomontana* F. K. Mey. by Meyer (1985). Populations from the Stara Planina, Vitoša, Pirin and Rodopi Mts, reported as *S. hungarica*, were ordered to the species *Soldanella chrysosticta* Kress (Kress 1984) and *Soldanella rhodopea* F. K. Mey. (Meyer 1985). The distribution area of *Soldanella hungarica* sensu Pawłowska became thus restricted to the Carpathians and the Austrian Alps. Although Meyer (l. c.) ordered the Austrian populations to the new-described species *Soldanella stiriaca* F. K. Mey., which is however not considered even in the current Austrian identification key (Fischer et al. 2008), according to Zhang & Kadereit (2002) this represents a superfluous name for *Soldanella major* (Neilr.) Vierh.

This for a long time accepted concept was markedly discredited by a serie of molecular-taxonomic and nomenclatoric studies (Zhang et al. 2001, Zhang & Kadereit 2002, 2004). *Soldanella hungarica* (s. l.), the species of the Eastern Alps and Carpathians, was divided to several, mainly new-described and narrowly-perceived endemic species: *Soldanella angusta* Zhang (the Ukrainian and Romanian Eastern Carpathians); *Soldanella marmarossiensis* Klášterský (the Slovak and Polish Western Carpathians; the Polish, Ukrainian and Romanian Eastern Carpathians); *Soldanella rugosa* Zhang (the Eastern Carpathians: Rodna Mts); *Soldanella oreodoxa* Zhang (the Apuseni Carpathians); *S. hungarica* Simonk. s. str. (syn. *S. pseudomontana* F. K. Mey.; the Southern Carpathians); *S. major* (Neilr.) Vierh. (syn. *S. stiriaca* F. K. Mey.; the Southern Carpathians and the Eastern Alps). However, the credibility of the distinguished species is very low considering a limited number of analysed individuals as well as a low-level separation in the obtained phylogenetic analyses (cf. Zhang et al. 2001); moreover, the area of Carpathians was insufficiently covered by samples.

Further ambiguity to the concept was introduced by Niederle (2003) by description of a new West-Carpathian species *Soldanella tatricola*, to which also plants from the Vysoké Tatry and Nízke Tatry Mts were ordered, by other authors reported as *S. montana* Willd., *S. hungarica* Vierh., *S. major* sensu Vierh. and *S. marmarossiensis* Klášť. Populations from the Polish Tatry foothills (Podtatrze, Wzniesienie Gubałowskie), until then reported as *Soldanella montana* or *S. pseudomontana* (cf. Pawłowska 1963, Kliment 1999), he described as a new subspecies *Soldanella montana* subsp. *gubalowkae*. A narrowly-perceived species *S. rugosa* (see above) was evaluated (Niederle 2003, 2005) as a superfluous name of *S. marmarossiensis* described from the Ukrainian Carpathians (Klášterský 1930); from the new-described narrow species (Zhang & Kadereit 2002) he only accepted the Apuseni endemic *S. oreodoxa*.

²⁵⁷ A hybridogenous species *Sorbus paxiana* is up to now known only from three localities in the Southern Carpathians: Băile Herculane: Valea Cernei; Petroșani; Piatra Rosia (Jávorka 1927, Kárpáti 1960). More current data on its distribution are missing. Already Kárpáti (1960) evaluated *S. paxiana* only according to the literature data and herbarium specimens. Oprea (2005) reports it from the Cerna and Mehedinți Mts.

²⁵⁸ An alpine taxon *Trisetum alpestre* subsp. *glabrescens* was described by Schur (1866) ranked as a variety from the Piatra Craiului Mts in the Southern Carpathians (cf. Buia & Morariu 1972). Cvelev (1974, 1976) and Prokudin et al. (1977) reported it from the montane belt of the Ukrainian Carpathians (Čyvčyny Mts: Čornyj dil) based on herbarium material of I. V. Artemčuk from 1961 wrongly determined as *Trisetum macrotrichum*. Čornej (2011) evaluated *Trisetum *glabrescens* as endemic to the Eastern and Southern Carpathians with a comment that its taxonomic rank and distribution require further study.

References

- Antosyak T. & Kozurak A. (2011): Distribution of endemic and rare vascular plant species in the upper Tysa Basin within the Carpathian biosphere reserve (Ukraine). – Transylv. Rev. Syst. Ecol. Res. 11: 85–94.
- Beldie A. (1967b): Flora și vegetația Munților Bucegi [Flora and vegetation of the Bucegi Mts]. – Edit. Acad. Rep. Soc. România, București.
- Beldie A. (1979): Flora României. Determinator ilustrat al plantelor vasculare II [Flora of Romania. Illustrated identification key of vascular plants II]. – Edit. Acad. Rep. Soc. România, București.
- Beldie A. & Alexandrescu L. (1976): Modificări de taxonomie sau nomenclatură și sinonimizări la principalii taxonii din vol. I–XII Flora R. S. R. [Changes in taxonomy or nomenclature and synonymy of the main taxa in volumes I–XII of the Flora R. S. R.]. – In: Beldie A. & Morariu I. (eds), Flora Republicii Socialiste România 13, p. 72–96. Edit. Acad. Rep. Soc. România, București.
- Bělohávková R. (2010): Amaryllidaceae J. St.-Hil. – In: Chrtek J. jr., Kaplan Z. & Štěpánková J. (eds), Květena České republiky 8, p. 677–687. Academia, Praha.
- Borodina A. E. (1979): O vidach roda *Rumex* L. evropejskoj časti SSSR. 2. Podrody *Acetosa* (Mill.) Rech. f. i *Acetosella* (Meissn.) Rech. f. [De generis *Rumex* L. speciebus in parte Europaea URSS crescentibus. 2. Subgenera *Acetosa* (Mill.) Rech. f. et *Acetosella* (Meissn.) Rech. f.]. – Novosti Sist. Vyssh. Rast. 15 (1978): 99–112.
- Boros Á. (1972): *Hieracium praebiharicum* Boros, spec. nova. – Bot. Közlem. 59: 33–35.
- Bucalo B., Brojić J., Travar J. & Milanović B. (2006): Sistematski pregled vaskularne flore Nacionalnog parka „Kozara“ [Systematic survey of vascular flora of the National park „Kozara“]. – Šumarstvo 4: 11–24.
- Buia A. & Morariu I. (1972): *Trisetum* Pers. – In: Nyárády E. I. et al. (eds), Flora Republicii Socialiste România 12, p. 287–295. Edit. Acad. Rep. Soc. România, București.
- Chrtek J. (1985a): Poznámky k druhu *Scabiosa lucida* s. l. v Karpatech [Notes on the species *Scabiosa lucida* s. l. in the Carpathians]. – Preslia 57: 199–203.
- Chrtek J. (1985b): *Scabiosa* L. – In: Bertová L. (ed.), Flóra Slovenska IV/2, p. 178–191. Veda, Bratislava.
- Ciocârlan V. (2009): Flora ilustrată a României. Pteridophyta et Spermatophyta [Illustrated flora of Romania. Pteridophyta et Spermatophyta]. – Editura Ceres, București.
- Cvelev N. N. (1961): *Leucanthemum* Mill. – In: Komarov V. L. (ed.), Flora SSSR 26, p. 140–145. Izdatel'stvo Akademii nauk SSSR, Moskva & Leningrad.
- Cvelev N. N. (1974): Poaceae Barnh. (Graminae Juss. nom. altern.). – In: Fedorov A. A. (ed.), Flora evropejskoj časti SSSR 1, p. 117–368. Nauka, Leningrad.
- Cvelev N. N. (1976): Zlaki SSSR [Gramineae of USSR]. – Nauka, Leningrad.
- Cvelev N. N. (1994): *Leucanthemum* Mill. – In: Cvelev N. N. (ed.), Flora evropejskoj časti SSSR 7, p. 136–139. Nauka, Leningrad.
- Cvelev N. N. (ed.) (1996): Flora Vostočnoj Evropy [Flora of Eastern Europe]. – Mir i Sem'ja 95, Sankt-Peterburg.
- Čopyk V. I. (1976): Vysokohirna flora Ukraïns'kykh Karpat [High-mountain flora of the Ukrainian Carpathians]. – Naukova dumka, Kyïv.
- Čornej I. I. (2011): Krytyčna revizija taksoniv, ščo navodylis' jak endemični dlja flory Ukraïns'kykh Karpat [Critical revision of taxa reported as endemic to flora of the Ukrainian Carpathians]. – Nauk. Zap. Bukovins'k. Tovar. Pryr. 1: 23–59.
- Danihelka J., Chrtek J. jr. & Kaplan Z. (2012): Checklist of vascular plants of the Czech Republic. – Preslia 84: 647–811.
- Dihoru G. & Negrean G. (2009): Cartea roșie a plantelor vasculare din România [Red data book of vascular plants of Romania]. – Edit. Acad. Române, București.
- Dihoru G. & Pârnu C. (1987): Plante endemice în flora Românie [Endemic plants in the flora of Romania]. – Editura Ceres, București.
- Domín K. (1931b): Rasy a formy ostřice vřdyzelené (*Carex sempervirens* Vill.) [Varieties and forms of *Carex sempervirens* Vill.]. – Rozpr. České Akad. Věd, Tř. 2, Vědy Mat.-Přír. 41/9: 1–11.
- Dostál J. (1976): *Centaurea* L. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 254–301. Cambridge University Press, Cambridge.
- Dostál L., Feráková V. & Tulenková M. (1999): *Leucojum vernum* L. subsp. *carpaticum* (Spring.) O. Schwarz. – In: Čerňovský J. et al. (eds), Červená kniha ohrožených a vzácných druhov rastlín a živočíchov SR a ČR 5, p. 217. Vyššie rastliny. Príroda, Bratislava.
- Dunkel F. G. (2011): Typification of *Ranunculus binatus* Kit. ex Rchb. (Ranunculaceae). – Willdenowia 41: 239–243.
- Edmondson J. R. (1980): *Poa* L. – In: Tutin T. G. et al. (eds), Flora Europaea 5, p. 159–167. Cambridge University Press, Cambridge.
- Ehrendorfer F. (1976): *Knautia* L. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 60–67. Cambridge University Press, Cambridge.
- Eliáš P. jr., Dítě D., Kliment J., Hrivnák R. & Feráková V. (2015): Red list of ferns and flowering plants of Slovakia, 5th edition (October 2014). – Biologia 70: 218–228 & Electronic Appendix.
- Fedorončuk M. M. (2009): Analiz endemizmu Caryophyllaceae Juss. flory Ukraïny [Analysis of endemism of Caryophyllaceae Juss. in the flora of Ukraine]. – Ukraïns'k. Bot. Zhurn. 66: 541–549.

- Fedorončuk M. M. & Čornej I. I. (2005): Rid *Dianthus* L. (Caryophyllaceae Juss.) flory Ukraïny: taksonomičnyj i sozologičnyj analiz [Genus *Dianthus* L. (Caryophyllaceae Juss.) in the flora of Ukraine – taxonomic and sozological analysis]. – Zapovidna sprava v Ukraïni 11, 2: 9–18.
- Fedorončuk M. M. & Diduch J. P. (2002a): *Dianthus carpaticus* Woł. – In: Diduch J. P. (ed.), Ekoflora Ukraïny 3, p. 386–387. Fitosociocentr, Kyïv.
- Feráková V. & Somogyi J. (1999): *Allium zahariadi* Májovský subsp. *michalkoi* Májovský. – In: Čerovský J. et al. (eds), Červená kniha vzácnych a ohrozených druhov vyšších rastlín SR a ČR 5. Vyššie rastliny, p. 25. Príroda, Bratislava.
- Fischer M. A., Oswald K. & Adler W. (2008): Exkursionsflora für Österreich, Liechtenstein und Südtirol. Ed. 3. – Biologiezentrum der Oberösterreichischen Landesmuseen, Linz.
- Ghișa E. & Beldie A. (1972): *Poa* L. – In: Nyárády E. I. et al. (eds), Flora Republicii Socialiste România 12, p. 364–429. Edit. Acad. Rep. Soc. România, București.
- Grabovskaja-Borodina A. E. (2012): *Rumex* L. – In: Gel'tman D. V. (ed.), Konspekt flory Vostočnoj Evropy, p. 306–316. Tovariščestvo naučnych izdanij KMK, Sankt-Peterburg & Moskva.
- Grau J. (1964): Zur Zytotaxonomie der *Myosotis alpestris*- und der *Myosotis silvatica*-Gruppe in Europa. – Oesterr. Bot. Z. 111: 561–617.
- Grau J. & Merxmüller H. (1972): *Myosotis* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 111–117. Cambridge University Press, Cambridge.
- Greuter W., Barrie F. R., Burdet H. M., Chaloner W. G., Demoulin V., Hawksworth D. L., Jørgensen P. M., Nicolson D. H., Silva P. C., Trehane P. & McNeill J. (eds) (1994): International Code of Botanical Nomenclature (Tokyo Code), adopted by the Fifteenth International Botanical Congress, Yokohama, August – September 1993. – Regnum Veg. 131. Koeltz Scientific Books, Königstein.
- Grințescu G. (1953): *Aconitum* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 3, p. 460–511. Edit. Acad. Rep. Pop. Romîne, București.
- Grințescu G. & Nyárády E. I. (1960b): *Myosotis* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 7, p. 237–254. Edit. Acad. Rep. Pop. Romîne, București.
- Gruľich V. (2012): Red list of vascular plants of the Czech Republic: 3rd edition. – Preslia 84: 631–645 & supplementary information.
- Gușuleac M. (1960): *Pulmonaria* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 7, p. 254–271, 644. Edit. Acad. Rep. Pop. Romîne, București.
- Halliday G. (1993): *Minuartia* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 152–160. Cambridge University Press, Cambridge.
- Heltmann H. (1985): Zum Endemismus des Südost-Karpatenraumes. – In: Heltmann H. & Wendelberger G. (eds), Naturwissenschaftliche Forschungen über Siebenbürgen. III. Beiträge zur Pflanzengeographie des Südost-Karpatenraumes. Siebenbürg. Arch., III. Folge, 20, p. 157–216.
- Heywood V. H. (1976): *Leucanthemum* Miller. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 174–177. Cambridge University Press, Cambridge.
- Holuby J. L. (1873): Die Brombeeren der Flora von Ns. Podhrad in Ungarn. – Oesterr. Bot. Z. 23: 373–384.
- Hrouda L. & Kochjarová J. (1997): K taxonómii a cenológii rodu *Soldanella* L. v Západných Karpatoch. Taxonomické hodnotenie pieninských populácií [On the taxonomy and coenology of the genus *Soldanella* L. in the West Carpathians. Taxonomical evaluation of the populations from the Pieniny Mts]. – Monografické štúdie o národných parkoch 1: 38–50.
- Hurdu B. I., Pușcaș M., Turtureanu P. D., Niketić M., Coldea G. & Zimmermann N. E. (2012a): Patterns of plant endemism in the Romanian Carpathians (South-Eastern Carpathians). – Contr. Bot. Cluj 47: 25–38.
- Hurdu B. I., Pușcaș M., Turtureanu P. D., Niketić M., Vonica G., Coldea G. (2012b): A critical evaluation of the Carpathian endemic plant taxa list from the Romanian Carpathians. – Contr. Bot. Cluj 47: 39–47.
- Jalas J. & Suominen J. (eds) (1979): Atlas Florae Europaeae 4. Polygonaceae. – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J. & Suominen J. (eds) (1983): Atlas Florae Europaeae 6. Caryophyllaceae (Alsinoideae and Paronychioideae). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J. & Suominen J. (eds) (1986): Atlas Florae Europaeae 7. Caryophyllaceae (Silenoideae). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jávorka S. (1927): A *Sorbus torminalis* (L.) Cr. magyar keverékfajai [Die *Sorbus torminalis*-Bastarde in Ungarn]. – Magyar Bot. Lapok 25 (1926): 83–90.
- Juksip A. J. (1959): Species novae generis *Hieracium* L. ex URSS. – Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 19: 467–530.
- Kaplan Z. (2012): Flora and phytogeography of the Czech Republic. – Preslia 84: 505–573.
- Kárpáti Z. (1960): Die *Sorbus*-Arten Ungarns und der angrenzender Gebiete. – Feddes Repert. 62: 71–331.
- Klásterský I. (1930): Ad floram Carpatorossicam aditamenta critica. Pars II. – Preslia 9: 19–20.

- Kliment J. (1999): Komentovaný prehľad vyšších rastlín flóry Slovenska, uvádzaných v literatúre ako endemické taxóny [Annotated survey of the vascular plants of Slovak flora recorded in literature as endemic taxa]. – Bull. Slov. Bot. Spoločn. 21, Suppl. 4: 1–434.
- Klovov M. V. (1952): Caryophyllaceae Juss. – In: Kotov M. I. (ed.), Flora URSS 4, p. 421–646. Vydavnyctvo Akademii nauk Ukraïns'koï RSR, Kyïv.
- Kobiv J. (1999): *Cortusa matthioli* (Primulaceae) in the Chornohora Mts (Ukrainian Carpathians): distribution, ecology, taxonomy and conservation. – Fragm. Florist. Geobot. 44: 355–374.
- Kobiv J. (2009): *Cortusa matthioli* L. subsp. *pubens* (Schott, Nyman et Kotschy) Jáv. – In: Diduch J. P. (ed.), Červona knyha Ukraïny. Roslynnij svit, p. 543. Globalconsalting, Kyïv.
- Kobiv J. (2010): Ekolohični osoblyvosti oselyšč ridkisnych vydiv roslyn Ukraïns'kich Karpat [Ecological characteristics of habitats of rare plant species in the Ukrainian Carpathians]. – Ukraïns'k. Bot. Zhurn. 67: 355–372.
- Kobiv J. (2012a): Roľ prydatnych mikrooselyšč u samovidovlenni populacij ridkisnych vydiv roslyn Ukraïns'skych Karpat [Role of suitable microhabitats in population recruitment of rare plant species in the Ukrainian Carpathians]. – Ukraïns'k. Bot. Zhurn. 69: 178–189.
- Kovtonjuk N. K. (2013): Zametki o podsekcii *Cortusa* (L.) Kovt. sekcii *Cortusoides* Balf. f. roda *Primula* L. (Primulaceae) [Notes on the genus *Primula* L. section *Cortusoides* Balf. f. subsection *Cortusa* (L.) Kovt. (Primulaceae)]. – Novosti Sist. Vyssh. Rast. 44: 152–166.
- Kress A. (1984): Chromosomenzählungen an verschiedenen Primulaceen. B. *Soldanella*. – Primulaceen-Studien 7, Gröbenzell bei München.
- Kricsfalusy V. (1999): Flora and vegetation of the Ukrainian upper Tisa Basin: Aspects of biodiversity conservation. – In: Hamar J. & Sárkány-Kiss A. (eds), The upper Tisa valley. Preparatory proposal for Ramsar site designation and an ecological background Hungarian, Romanian, Slovakian and Ukrainian co-operation. Tiscia monograph series 4, p. 273–292. Szeged.
- Kricsfalusy V. & Budnikov G. (2007): Threatened vascular plants in the Ukrainian Carpathians: current status, distribution and conservation. – Thaiszia-J. Bot. 17: 11–32.
- Kubát K. (2010): *Dactylorhiza* Nevski. – In: Chrtěk J. jr., Kaplan Z. & Štěpánková J. (eds), Květena České republiky 8, p. 502–523. Academia, Praha.
- Kuz'mina M. L. (2004): *Dianthus* L. – In: Cvelev N. N. (ed.) Flora vostočnoj Evropy 11, p. 273–297. Tovarišestvo naučnych izdanij KMK, Moskva & Sankt-Peterburg.
- Luceño M., Escudero M. & Jiménez-Mejías P. (2008): *Carex* L. – In: Castroviejo S. (ed.), Flora Iberica 18, p. 109–250. C. S. I. C., Madrid.
- Májovský J. & Murin A. (1985): Karyotaxonomisches Studium des *Allium ericetorum* aggr. – Acta Fac. Rerum Nat. Univ. Comen., Bot. 32: 3–20.
- Malynovs'kyj K., Caryk J., Kyjak V. & Nesteruk J. (2002): Ridkysni, endemični, reliktovi ta pohranyčno-areaľni vydy roslyn Ukraïns'skych Karpat [Rare, endemic, relict and boundary areal plant species of the Ukrainian Carpathians]. – Instytut ekologii Karpat Nacional'noï akademiï nauk Ukraïny, Lviv.
- Marhold K. (ed.) (1998): Paprad'orasty a semenné rastliny [Ferns and flowering plants]. – In: Marhold K. & Hindák F. (eds), Zoznam nižších a vyšších rastlín Slovenska, p. 297–331. Veda, Bratislava.
- Marhold K., Mártonfi P., Mered'a P. & Mráz P. (eds) (2007): Chromosome number survey of the ferns and flowering plants of Slovakia. – Veda, Bratislava.
- Mast A. R., Kelso S., Richards A. J., Lang D. J., Feller D. M. S. & Conti E. (2001): Phylogenetic relationships in *Primula* L. and related genera (Primulaceae) based on noncoding chloroplast DNA. – Int. J. Plant Sci. 162: 1381–1400.
- McNeill J., Barrie F. R., Buck W. R., Demoulin V., Greuter W., Hawksworth D. L., Herendeen P. S., Knapp S., Marhold K., Prado J., Prud'home van Reine W. F., Smith G. F., Wiersema J. H. & Turland N. J. (eds) (2012): International Code of Nomenclature for algae, fungi, and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. – Regnum Veg. 154. Koeltz Scientific Books, Königstein.
- Merxmüller B. & Sauer W. (1972): *Pulmonaria* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 100–102. Cambridge University Press, Cambridge.
- Meyer F. K. (1985): Beitrag zur Kenntniss ost und südosteuropäischer *Soldanella*-Arten. – Haussknechtia 2: 7–41.
- Mirek Z. & Piękoś-Mirkowa H. (2009): Fitogeograficzne aspekty endemizmu w Polsce [Phytogeographical aspects of endemism in Poland]. – Wiadom. Bot. 53: 7–30.
- Mirek Z. & Piękoś-Mirkowa H. (2010): Fitogeograficzna analiza endemitów roślin naczyniowych Tatrzańskiego Parku Narodowego [Phytogeographical analysis of the endemic vascular plant taxa occurring in the Tatry Mts]. – Nauka a zarządzanie obszarem Tatr i ich otoczeniem 2: 11–15.
- Mitka J. (2008): *Aconitum moldavicum* Hacq. (Ranunculaceae) and its hybrids in the Carpathians and adjacent regions. – Roczn. Bieszcz. 16: 233–252.
- Mitka J. & Gawroński S. (2010): Genus *Aconitum* in Transcarpathia. – In: Dva storiččja doslidžennja roslynnoho pokryvu Karpat. Materialy mižnarodnoï naukovoï konferencii, prysvjačenoï 130-riččju narodžennja Antonija Margitaja (16–18 veresnja 2010 r., Mukačevo-Berehovo, Ukraïna), p. 129–130. Karpaty, Užhorod.
- Mitka J. & Novikoff A. V. (2011): The genus *Aconitum* in Ukraine. – In: Materialy XIII Z'їzdu Ukraïns'koho botaničnogo tovarystva (Lviv, 19–23 veresnja 2011 r.), p. 65. Lviv.

- Morariu I. & Beldie A. (1976): Endemismele din Flora R.S.R. [Endemics in the Flora R. S. R.]. – In: Beldie A. & Morariu I. (eds), Flora Republicii Socialiste România 13, p. 97–105. Edit. Acad. Rep. Soc. România, București.
- Morariu I. & Nyárády E. I. (1960): *Cortusa* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 7, p. 58–61. Edit. Acad. Rep. Pop. Romîne, București.
- Morariu I., Nyárády E. I. & Gușuleac M. (1960): *Primula* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 7, p. 77–100. Edit. Acad. Rep. Pop. Romîne, București.
- Negrean G. & Oltean M. (1989): Endemite și zone endemoconservatoare din Carpații Sud-Estici [Endemics and endemoconservation areas in the South-Eastern Carpathians]. – Ocrot. Nat. 33: 15–25.
- Niederle J. (2003): Remarks on the genus *Soldanella* L. in the West Carpathians. – Acta Mus. Richnov., Sect. Nat. 10: 171–174.
- Niederle J. (2005): *Soldanella rugosa* is synonym to real *S. marmarossiensis*. – Nord. J. Bot. 24: 547–548.
- Niketić M. (2010): Novi taksoni za floru cvetnica Srbije i susednih područja [New taxa in the flora of Serbia and neighbouring regions]. – In: 10th Symposium on the flora of southeastern Serbia and neighbouring regions, Vlasina Lake 17 to 20 June 2010. Abstracts, p. 32–33. Department of Biology and Ecology, Faculty of Sciences and Mathematics, University of Niš & Biological Society „Dr Sava Petrović“, Niš.
- Novikov A. V. (2013b): Ranunculaceae Zachidnoi Ukraïny. I. Ključ dlja vyznačennja vydiv [Ranunculaceae of the Western Ukraine. Identification key]. – Modern Phytomorphol. 3: 297–319.
- Novikoff A. V. & Mitka J. (2011a): Taxonomy and ecology of the genus *Aconitum* L. in the Ukrainian Carpathians. – Wulfenia 18: 37–61.
- Novikov A. V. & Mitka J. (2011b): Rid *Aconitum* v Ukraïns'kykh Karpatach [The genus *Aconitum* in the Ukrainian Carpathians]. – Biol. Stud. 5: 153–172.
- Nyárády E. I. (1928): Über zwei neue und seltene Gräser Rumäniens. – Verh. Mitt. Naturwiss. Hermannstadt 78: 144–153.
- Nyárády E. I. (1956): *Rubus* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 4, p. 276–580. Edit. Acad. Rep. Pop. Romîne, București.
- Nyárády E. I. 1965: Compositae Giseke. – In: Nyárády E. I. (ed.), Flora Republicii Populare Romîne 10, p. 17–736. Edit. Acad. Rep. Pop. Romîne, București.
- Oprea A. (2005): Lista critică a plantelor vasculare din România [Critical list of vascular plants of Romania]. – Editura Universității „Alexandru Ioan Cuza“, Iași.
- Oprea A. & Sirbu C. (2013): The vascular flora of Rarău massif (Eastern Carpathian, Romania). Note II. – Mem. Sci. Sect. Romanian Acad., Biol. 36: 18–52.
- Papović O., Miljković M., Randelović N. & Randelović V. (2014): Analysis of the flora of Rogozna Mountain in Southwestern Serbia. – Biol. Nyssana 5: 17–30.
- Pawłowska S. (1963): De Soldanellis, quae in parte septentrionali Carpatorum crescunt. – Fragm. Florist. Geobot. 9: 3–30.
- Pawłowski B. (1970): Remarques sur l'endémisme dans la flore des Alpes et des Carpates. – Vegetatio 21: 181–243.
- Piękoś-Mirkowa H. & Mirek Z. (2003): Endemic taxa of vascular plants in the Polish Carpathians. – Acta Soc. Bot. Polon. 72: 235–242.
- Popescu A. & Sanda V. (1998): Conspectul florei cormofitelor spontane din România [Enumeration of spontaneous vascular plants of Romania]. – Editura Universității din București, București.
- Prodan I. & Nyárády E. I. (1964): *Centaurea* L. – In: Nyárády E. I. (ed.), Flora Republicii Populare Romîne 9, p. 785–951. Edit. Acad. Rep. Pop. Romîne, București.
- Prokudin J. N. (ed.) (1987): Oprelitel' vysšich rastenij Ukraïny [Identification key of vascular plants of Ukraine]. – Naukova dumka, Kiïv.
- Prokudin J. N., Vovk A. G., Petrova O. A., Ermolenko E. D. & Verničenko J. D. (1977): Zlaki Ukraïny [The grasses of Ukraine]. – Naukova dumka, Kiev.
- Rechinger K. H. & Akeyrod J. R. (1993): *Rumex* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 99–107. Cambridge University Press, Cambridge.
- Sârbru I., Ștefan N. & Oprea A. (2013): Plante vasculare din România. Determinator ilustrat de teren [Vascular plants of Romania. An illustrated field guide]. – Victor B Victor, București.
- Schur F. (1866): Enumeratio Plantarum Transsilvaniae. – Guilielmum Braumüller, Vindobonae.
- Schwarz O. (1968): Beiträge zur Kenntnis der Gattung *Primula*. – Wiss. Zeitschr. Univ. Jena, Sect. Math.-Natur. 17: 307–332.
- Șerbănescu G. (1967): Poziția taxonomică și gradul afinităților unor taxoni critici în raport cu *Poa laxa* Hke din Carpații Românești [Taxonomic position and the affinity degree of some critical taxa in regards to *Poa laxa* Hke from the Romanian Carpathians]. – Stud. Cerc. Biol., Bot. 10: 389–399.
- Soják J. (1962): Novinky československé květeny [Neue Pflanzen der tschechoslowakischen Flora]. – Preslia 34: 403–414.
- Soják J. (1993): Taxonomische Bemerkungen zu einigen mediterranen *Potentilla*- Sippen. – Preslia 65: 117–130.
- Somogyi J. (1999a): Taxonomical and nomenclatural remarks on the *Allium ericetorum* group with regards to the Western Carpathians. – Biologia 54: 391–394.
- Somogyi J. (1999b): Rozšírenie druhu *Allium ericetorum* Thore na Slovensku [Distribution of *Allium ericetorum* Thore in Slovakia]. – Nat. Carpat. 40: 201–204.

- Somogyi J. (2002): Komentovaný červený zoznam taxónov rodu *Allium* L. na Slovensku [Commented red list of taxa of the genus *Allium* L. in Slovakia]. – Bull. Slov. Bot. Spoločn. 24: 97–100.
- Starmühler W. (1998a): Ranunculaceae subtrib. Delphiniinae exsiccatae, Fasc. 3 (Nr. 41–60). – Fritschiana 13: 1–10.
- Starmühler W. (1998b): Aconita rarissima selecta (1998). – Fritschiana 13: 11–39.
- Stojko S. M. & Tasenkewitsch L. O. (1991): Pflanzengeographische Stellung und Schutz von Flora und Vegetation der Ukrainischen Karpaten. – Verh. Zool.-Bot. Ges. Österreich 128: 165–172.
- Stoyko S. M. & Tasenkevich L. (1993): Some aspects of endemism in the Ukrainian Carpathians. – Fragm. Florist. Geobot., Suppl. 2 (1): 343–353.
- Šeffler J., Lasák R., Šefflerová-Stanová V., Janák M. & Guttová A. (2010): Towards an Ecological Network for the Carpathians II. – Carpathian Ecoregion Initiative, Bratislava.
- Šípošová H., Bernátová D., Mráz P., Kliment J., Goliašová K. & Uhlířová J. (2004a): Taxóny cievnatých rastlín, ktoré boli opísané z Veľkej Fatry. 1. časť [Vascular plant taxa described from the Veľká Fatra Mts. Part 1]. – In: Kadlečík J. (ed.), Turiec a Fatra 2004, p. 77–85. Správa Národného parku Veľká Fatra, Vrútky.
- Šljakov P. N. (1989): *Hieracium* L. – In: Cvelev N. N. (ed.), Flora evropejskoj časti SSSR 8, p. 140–300. Nauka, Leningrad.
- Šourková M. (1984): *Bupleurum* L. – In: Bertová L. (ed.), Flóra Slovenska IV/1, p. 284–309. Veda, Bratislava.
- Štěpánek J. & Holub J. (1997): *Scabiosa* L. – In: Slavík B. (ed.), Květena České republiky 5, p. 536–542. Academia, Praha.
- Tasenkevich L. (2002): Red list of vascular plants of the Carpathian mountains. – State museum of natural history NAS of Ukraine, Lviv.
- Tasenkevych L. (2003b): Rozmaïtja flory sudynnych roslyn v Ukraïns'kykh Karpatach [Vascular plants flora diversity of the Ukrainian Carpathians]. – Pratsi Nauk. Tovar. Schevchenka 12: 147–157.
- Tasenkevych L. (2010): Arko-tretynnyj element flory Karpat [Arctic-Tertiary element in the Carpathians' flora]. – Nauk. Zap. Derzhavn. Pryr. Muz. 26: 95–112.
- Tasenkevich L. (2011): Endemism in the Carpathian flora – a chorological aspect. – In: Zemanek B. (ed.), Geobotanist and Taxonomist. A volume dedicated to Professor Adam Zajac on the 70th anniversary of his birth, p. 157–168. Institute of Botany Jagellonian University, Cracow.
- Tasenkevych L. (2014): Endemizm flory Karpat – ekolohični ta chorolohični osoblyvosti [Endemism of the Carpathians' flora – ecological and chorological peculiarities]. – Pratsi Nauk. Tovar. Schevchenka 39: 22–36.
- Tutin T. G. (1964): *Dianthus* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1, p. 188–204. Cambridge University Press, Cambridge.
- Tutin T. G. (1968): *Bupleurum* L. – In: Tutin T. G. et al. (eds), Flora Europaea 2, p. 345–351. Cambridge University Press, Cambridge.
- Valentine D. H. & Kress A. (1972): *Primula* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 15–20. Cambridge University Press, Cambridge.
- Walter K. S. & Gillett H. J. (eds) (1998): 1997 IUCN Red list of threatened plants. – The World Conservation Union, Gland & Cambridge.
- Yan H.-F., He Ch.-H., Peng Ch.-I., Hu Ch.-M. & Hao G. (2010): Circumscription of *Primula* subgenus *Auganthus* (Primulaceae) based on chloroplast DNA sequences. – J. Syst. Evol. 48: 123–132.
- Zahariadi C. (1966): Liliaceae A. L. Jussieu. – In: Nyárády E. I. (ed.), Flora Republicii Socialiste România 11, p. 106–404. Edit. Acad. Rep. Soc. România, București.
- Zahn K. H. (1927): Beiträge zur Kenntnis der Hieracien Ungarns und der Balkanländer VII. – Magyar Bot. Lapok 25 (1926): 121–174.
- Zahn K. H. (1930–1935): *Hieracium*. – In: Graebner P. jr. (ed.), Synopsis der mitteleuropäischen Flora 12/2. – Gebrüder Borntraeger Verlag, Leipzig.
- Zahn K. H. (1936–1938): *Hieracium*. – In: Graebner P. jr. (ed.), Synopsis der mitteleuropäischen Flora 12/3. – Gebrüder Borntraeger Verlag, Leipzig.
- Zelený V. (1982): Kopretina bílá (*Leucanthemum vulgare* Lam. complex) v Československu [*Leucanthemum vulgare* complex in Czechoslovakia]. – Stud. Českoslov. Akad. Věd 10: 1–135.
- Zhang L.-B. & Kadereit J. W. (2002): The systematics of *Soldanella* (Primulaceae) based on morphological and molecular (ITS, AFLPs) evidence. – Nord. J. Bot. 22: 129–169.
- Zhang L.-B. & Kadereit J. W. (2004): Nomenclature of *Soldanella* L. (Primulaceae). – Taxon 53: 741–752.
- Zhang L.-B., Comes H. P. & Kadereit J. W. (2001): Phylogeny and quaternary history of the European montane/alpine endemic *Soldanella* (Primulaceae) based on ITS and AFLP variation. – Amer. J. Bot. 88: 2331–2345.
- Ziman S. M., Bulach O. V. & Hamor A. F. (2006): Pro rídkišni vysokohirni vydy kvitkovych roslyn u flory Ukraïns'kykh Karpat – vikarianty vydiv z šyrokymy arealamy ta ekolohieju [About rare high-mountain species of the flowering plants within the flora of the Ukrainian Carpathians]. – Nauk. Zap. Derzhavn. Pryr. Muz., Bot. 22: 25–34.

Electronic Appendix 3. – Taxa with a wider non-endemic distribution

Taxon	Area	References
<i>Alchemilla gorcensis</i> Pawł.	Carpathians, the Balkan Peninsula	Kurtto et al. 2007
<i>Alchemilla mollis</i> (Buser) Rothm.	Turkey, Greece, Bulgaria, Romania	Kurtto et al. 2007; Kurtto in Euro+Med
<i>Alchemilla walasii</i> Pawł.	Western & Eastern Carpathians, Eastern Poland, Belarus, Podillja	Syčák 2002; Kurtto et al. 2007; Kurtto in Euro+Med
<i>Alyssum repens</i> Baumg. (Syn.: <i>A. transsilvanicum</i> Schur)	Austria, Romania, Croatia	Jalas et al. 1996; Španiel et al. 2011a, b; Španiel 2014 in litt.
<i>Androsace villosa</i> var. <i>arachnoidea</i> (Schott, Nyman et Kotschy) R. Knuth (Syn.: <i>A. villosa</i> subsp. <i>arachnoidea</i> (Schott, Nyman et Kotschy) Nyman)	South-eastern Europe, incl. Italy	Ferguson 1972a
<i>Arabidopsis arenosa</i> subsp. <i>borbasii</i> (Zapał.) O’Kane et Al-Shehbaz (Syn: <i>Cardaminopsis arenosa</i> subsp. <i>borbasii</i> (Zapał.) Pawł.)	Western, Central, Eastern & Southern Europe	Jalas & Suominen 1994
<i>Arabis hornungiana</i> Schur	Carpathians, Bulgaria, Croatia	Kliment 1999; Ciocârlan 2009
<i>Avenula pubescens</i> subsp. <i>laevigata</i> (Schur) Holub	Western & Southern Alps, Eastern & Southern Carpathians	Holub 1980; Ciocârlan 2009
<i>Betula pubescens</i> subsp. <i>carpatica</i> (Waldst. et Kit. ex Willd.) Asch. et Graebn.	Pyrenees, Alps, Hercynian massif, Carpathians	Sýkora 1983; Kliment 1999
<i>Bupleurum longifolium</i> subsp. <i>vapincense</i> (Vill.) Todor	Auvergne, Jura, Alps, Carpathians, Dinarides, Stara Planina Mts	Šourková & Hrouda 1997
<i>Campanula abietina</i> Griseb. (Syn.: <i>C. patula</i> subsp. <i>abietina</i> (Griseb.) Simonk.)	Carpathians, mountains of the Balkan Peninsula	Fedorov & Kovanda 1976; Goliašová et al. 2008
<i>Campanula glomerata</i> subsp. <i>elliptica</i> (Kit. ex Schult.) Kirschl. (Syn.: <i>C. elliptica</i> Kit. ex Schult.)	Carpathians, Serbia, Croatia, Italy	Fedorov & Kovanda 1976; Kliment 1999; Kovačić 2004; Goliašová et al. 2008
<i>Campanula transsilvanica</i> Schur ex Andrae	Eastern & Southern Carpathians; Pirin, Rila, Vitoša & Stara Planina Mts	Oprea 2005; Assyov & Petrova 2006
<i>Carduus collinus</i> Waldst. et Kit.	Western Carpathians, Ukraine, the Balkan Peninsula, Italy	Kliment 1999
<i>Centaurea degeniana</i> J. Wagner ²⁵⁹	Southern Carpathians, Western Bulgaria	Dostál 1976; Oprea 2005; Assyov & Petrova 2006
<i>Centaurea kotschyana</i> Heuff.	Eastern & Southern Carpathians, northern part of the Balkan Peninsula	Dostál 1976; Oprea 2005; Greuter in Euro+Med
<i>Chrysanthemum zawadzkii</i> Herbach ²⁶⁰ (Syn.: <i>Dendranthema zawadzkii</i> (Herbach) Tzvelev)	Western Carpathians (Pieniny Mts), Russia, Northern Mongolia, North-eastern China	Zarzycki 1976, 1982, 2000; Holub 1999; Wróbel 2008; Szeląg & Kobiv 2014
<i>Cirsium boujartii</i> (Piller et Mitterp.) Sch. Bip. subsp. <i>boujartii</i>	Hungary, Romania	Negrean & Oltean 1989; Csiky et al. 2005
<i>Cirsium decussatum</i> Janka	Poland (Pogórze Przemyskie Mts), Belarus, Ukraine, Romania (Eastern & Southern Carpathians); Moldova	Prokudin 1987; Oprea 2005; Piórecki & Zarzycki 2008; Ciocârlan 2009; Greuter in Euro+Med
<i>Cirsium grecescui</i> Rouy	Romania (Southern Carpathians), Serbia, Macedonia	Gajić 1975; Negrean & Oltean 1989; Melovski & Matevski 2008
<i>Colchicum haynaldii</i> Heuff. (Syn.: <i>C. neapolitanum</i> var. <i>haynaldii</i> (Heuff.) Asch. et Graebn.)	Romania (Southern Carpathians), the Balkan Peninsula	Oprea 2005; Ciocârlan 2009; Alexiou 2013
<i>Dactylis glomerata</i> subsp. <i>slovenica</i> Domin (Syn.: <i>D. slovenica</i> (Domin) Domin)	Sudetes, Western & Eastern Carpathians, Ukraine (outside the Carpathians), Montenegro (rare)	Kliment 1999
<i>Dactylorhiza fuchsii</i> subsp. <i>sooana</i> (Borsos) Borsos (Syn.: <i>D. longebracteata</i> subsp. <i>sooana</i> (Borsos) Dostál)	Western Carpathians, Transdanubian Mid-Mountains, Mecsek Mts, Pannonian Plain	Kliment 1999
<i>Dactylorhiza maculata</i> subsp. <i>transsilvanica</i> (Schur) Soó (Syn.: <i>D. transsilvanica</i> (Schur) Aver.)	Carpathians, mountains of the Balkan Peninsula	Soó 1980b, Kubát 2010
<i>Dianthus collinus</i> subsp. <i>glabriusculus</i> (Kit.) Thaisz (Syn.: <i>D. glabriusculus</i> (Kit.) Borbás; <i>D. collinus</i> subsp. <i>moldavicus</i> (Prodan) Soó)	Western Carpathians (HU), Eastern Carpathians, Transdanubian Mid-Mountains, Pannonian Plain; lowlands in Eastern Romania	Tutin & Walters 1993; Kliment 1999; Ciocârlan 2009

Taxon	Area	References
<i>Erigeron macrophyllus</i> Herbich (Syn.: <i>E. acris</i> subsp. <i>macrophyllus</i> (Herbich) Gutermann)	Eastern Alps, Hercynian massif, Carpathians, mountains of the Balkan Peninsula, Belarus, Russia, Caucasus	Šida 1996
<i>Erythronium dens-canis</i> var. <i>niveum</i> Baumg. (<i>E. dens-canis</i> subsp. <i>niveum</i> (Baumg.) Buia et Paun)	Apuseni Carpathians, Oltenia region (RO), Srbija, Bulgaria	Zahariadi 1966; Delipavlov 1971; Diklić & Nikolić 1986b; Borza et al. 1998
<i>Festuca xanthina</i> Roem. et Schult.	Southern Carpathians, mountains in northern part of the Balkan Peninsula	Negrean & Oltean 1989; Markgraf-Dannenberg 1980; Foggi & Müller in Euro+Med
<i>Gagea bohémica</i> (Zauschn.) Schult. et Schult. f.	Central, South-eastern & South-western Europe, Turkey, Israel	Hrouda 2010a
<i>Galium polonicum</i> Błocki (Syn.: <i>G. carpaticum</i> Klokov)	South-eastern Poland, North-eastern Ukraine, Eastern Carpathians (UA, rare)	Kucowa 1967; Ehrendorfer & Krendl 1976; Kułak & Sender 2011
<i>Galium pseudaristatum</i> Schur	Eastern & Southern Carpathians, Transylvanian Basin, mountains of the Balkan Peninsula	Ehrendorfer & Krendl 1976; Oprea 2005; Marhold in Euro+Med
<i>Geranium caeruleum</i> Schur (Syn.: <i>G. sylvaticum</i> subsp. <i>caeruleum</i> (Schur) D. A. Webb et I. K. Ferguson)	Eastern & Southern Carpathians, mountains of the Balkan Peninsula	Webb & Ferguson 1968; Oprea 2005
<i>Hieracium atrellum</i> (Zahn) Ůksip	Sudetes, Carpathians	Šljakov 1989; Chrtek jr. 2004; Oprea 2005
<i>Hieracium catenatum</i> Sennikov ²⁶¹	Eastern Carpathians (UA), Belarus	Sennikov 1995, 1999b
<i>Hieracium stygium</i> R. Uechtr.	Eastern Sudetes, Western Carpathians	Chrtek jr. 2004; Chrtek & Mráz 2007
<i>Hieracium wimmeri</i> R. Uechtr.	Sudetes, Western Carpathians (SK), Eastern Carpathians (UA)	Šljakov 1989; Szeląg 2011
<i>Hypericum richeri</i> subsp. <i>grisebachii</i> (Boiss.) Nyman (Syn.: <i>H. richeri</i> subsp. <i>transsilvanicum</i> (Čelak.) Ciocârlan)	South-eastern Alps; Eastern, Southern & Apuseni Carpathians; mountains of the Balkan Peninsula	Robson 1968, 2010; Oprea 2005; Čornej 2011
<i>Jacobaea abrotanifolia</i> subsp. <i>carpathica</i> (Herbich) B. Nord. et Greuter (Syn.: <i>Senecio abrotanifolius</i> subsp. <i>carpathicus</i> (Herbich) Nyman; <i>S. carpathicus</i> Herbich)	Western, Eastern & Southern Carpathians; mountains of the Balkan Peninsula	Kliment 1999; Oprea 2005
<i>Jovibarba globifera</i> subsp. <i>glabrescens</i> (Sabr.) Holub ²⁶² (Syn.: <i>J. hirta</i> subsp. <i>glabrescens</i> (Sabr.) Holub)	Western Carpathians, Apuseni Carpathians, Transdanubian Mid-Mountains, Mecsek Mts, Pannonian Plain (rare)	Letz 1998; Kliment 1999
<i>Larix decidua</i> subsp. <i>polonica</i> (Racib. ex Wóycicki) Domin (Syn.: <i>L. decidua</i> subsp. <i>carpatica</i> (Domin) Domin; <i>L. decidua</i> subsp. <i>carpatica</i> (Domin) Šiman, nom. illeg.)	?Sudetes, Carpathians, Poland (outside the Carpathians)	Browicz et al. 1971; Skalická & Skalický 1987; Kliment 1999
<i>Melittis melissophyllum</i> subsp. <i>carpatica</i> (Klokov) P. W. Ball (Syn.: <i>M. carpatica</i> Klokov)	eastern part of Central Europe, eastwards to Western Ukraine and South-western Belarus (incl. Western & Eastern Carpathians)	Ball 1972; Marhold 1993; Kliment 1999
<i>Minuartia frutescens</i> (Kit. ex Schult.) Tuzson ex Degen (Syn.: <i>M. hirsuta</i> subsp. <i>frutescens</i> (Kit. ex Schult.) Hand.-Mazz.)	Western, Southern & Apuseni Carpathians, Transylvanian Basin, mountains in northern part of the Balkan Peninsula	Kliment 1999; Letz 2012
<i>Muscari transsilvanicum</i> Schur (Syn.: <i>M. botryoides</i> subsp. <i>transsilvanicum</i> (Schur) Soó; <i>M. botryoides</i> subsp. <i>hungaricum</i> Priszter)	Eastern & Southern Carpathians, Pannonian Plain, Mecsek Mts, Villány Mts	Somlyay et al. 2006
<i>Noccaea kovatsii</i> (Heuff.) F. K. Mey. (Syn.: <i>Thlaspi kovatsii</i> Heuff.; <i>T. pawlowskii</i> Dvořáková)	Eastern, Southern & Apuseni Carpathians, mountains of the Balkan Peninsula	Clapham & Akeyrod 1993; Jalas et al. 1996; Oprea 2005; Kobiv et al. 2007a
<i>Ornithogalum kochii</i> Parl. (Syn.: <i>O. orthophyllum</i> subsp. <i>kochii</i> (Parl.) Zahar.)	South-eastern (incl. northern Italy) & Central Europe	Hrouda 2010b
<i>Petasites kablikianus</i> Tausch ex Bercht.	Sudetes, Carpathians, mountains of the Balkan Peninsula	Kliment 1999; Oprea 2005; Ciocârlan 2009 etc.
<i>Poa cenisia</i> subsp. <i>contracta</i> Nyár. (Syn.: <i>P. psychrophila</i> Boiss. et Heldr.)	Southern Carpathians, mountains of the Balkan Peninsula, Crete	Ghişa & Beldie 1972; Edmondson 1980; Ciocârlan 2009
<i>Polygala supina</i> subsp. <i>hospita</i> (Heuffel) McNeill	Southern Carpathians, northern part of the Balkan Peninsula	McNeill 1968; Negrean & Oltean 1989

Taxon	Area	References
<i>Primula halleri</i> subsp. <i>platyphylla</i> O. Schwarz	Carpathians, mountains of the Balkan Peninsula	Schwarz 1968; Nikolić & Diklić 1986; Kliment 1999; Assyov & Vassilev 2004; Mustafa et al. 2013; Schönschwetter & Božo 2013
<i>Pulmonaria rubra</i> Schott subsp. <i>rubra</i>	Eastern, Southern & Apuseni Carpathians, mountains of the Balkan Peninsula	Guşuleac 1960; Merxmüller & Sauer 1972
<i>Rhinanthus gracilis</i> Schur ²⁶³	Eastern & Southern Carpathians, Rila Mts	Păuca & Nyárády 1960; Assyov & Petrova 2006
<i>Rhododendron myrtifolium</i> Schott et Kotschy (Syn.: <i>R. kotschy</i> Simonk.)	Eastern & Southern Carpathians, mountains of the Balkan Peninsula	Popova 1972; Căprar et al. 2014
<i>Satureja kitaibelii</i> Wierzb. ex Heuffel (Syn.: <i>S. montana</i> subsp. <i>kitaibelii</i> (Wierzb. ex Heuffel) P. W. Ball)	Southern Carpathians (RO, SRB), north part of the Balkan Peninsula	Ball & Getliffe 1972; Ančev 1989; Negrean & Oltean 1989
<i>Saussurea discolor</i> (Willd.) DC.	Alps, Carpathians, Appeniny, Rila Mts	Lipschitz 1976; Assyov & Petrova 2006
<i>Saxifraga carpatica</i> Sternb. (Syn.: <i>S. carpathica</i> Rchb., nom. illeg.)	Eastern Alps (rare), Carpathians, northern part of the Balkan Peninsula	Webb 1993a; Schneeweiss 1998; Kliment 1999
<i>Saxifraga luteoviridis</i> Schott et Kotschy (Syn.: <i>S. corymbosa</i> Boiss.)	Eastern, Southern & Apuseni Carpathians, mountains of the Balkan Peninsula	Beldie 1977; Webb 1993a; Jalas et al. 1999; Kobiv et al. 2007a
<i>Saxifraga pedemontana</i> subsp. <i>cymosa</i> Engl.	Eastern & Southern Carpathians, mountains of the Balkan Peninsula	Jalas et al. 1999; Oprea 2005
<i>Scorzoneroideis rilaensis</i> (Hayek) Holub (Syn.: <i>Leontodon rilaensis</i> Hayek; <i>L. croceus</i> subsp. <i>rilaensis</i> (Hayek) Finch et P. D. Sell)	Eastern Carpathians (rare), Southern Carpathians, mountains of the Balkan Peninsula	Finch & Sell 1976; Ciocârlan 2009; Dakskobler et al. 2010; Oprea & Sîrbu 2013
<i>Securigera elegans</i> (Pančić) Lassen (Syn.: <i>Coronilla elegans</i> Pančić)	Western (SK, HU), Eastern & Southern Carpathians, mountains of the Balkan Peninsula	Chrtková 1988; Oprea 2005; Niketić & Tomović 2008
<i>Silene heuffelii</i> Soó	Eastern & Southern Carpathians, mountains of the Balkan Peninsula	Chater et al. 1993; Oprea 2005
<i>Swertia perennis</i> subsp. <i>alpestris</i> (Fuss) Simonk. (Syn.: <i>S. alpestris</i> Fuss; <i>S. perennis</i> var. <i>alpestris</i> (Fuss) Sagorski)	Alps, Hercynian Massif, Carpathians, Bulgaria (Vitoša Mts, Pirin Mts, Rila Mts)	Kliment 1999
<i>Tephrosieris papposa</i> (Rchb.) Schur (Syn.: <i>Senecio papposus</i> (Rchb.) Less.)	Eastern, Southern & Apuseni Carpathians, mountains of the Balkan Peninsula	Kliment 1999; Greuter in Euro+Med
<i>Tozzia carpathica</i> Woł. (Syn.: <i>T. alpina</i> subsp. <i>carpathica</i> (Woł.) Dostál)	Carpathians, mountains of the Balkan Peninsula	Kliment 1999; Sârbu et al. 2013
<i>Trifolium repens</i> var. <i>ochranthum</i> K. Malý (Syn.: <i>T. repens</i> subsp. <i>ochranthum</i> (K. Malý) Nyár.)	Eastern & Southern Carpathians, mountains in northern part of the Balkan Peninsula	Cincović 1972; Oprea 2005
<i>Trollius europaeus</i> subsp. <i>transsilvanicus</i> (Schur) Domin (Syn.: <i>T. transsilvanicus</i> Schur; <i>T. altissimus</i> Crantz)	South-eastern Alps, Carpathians	Jalas & Suominen 1989; Tutin 1993
<i>Verbascum glabratum</i> subsp. <i>brandzae</i> (Franch. ex Brândză) Murb.	Southern Carpathians (rare), mountains in northern part of the Balkan Peninsula	Ferguson 1972b; Oprea 2005; Dihoru & Negrean 2009
<i>Veronica baumgartenii</i> Roem. et Schult.	Eastern & Southern Carpathians, mountains in northern part of the Balkan Peninsula	Walters & Webb 1972; Peev 1995; Oprea 2005; Ciocârlan 2009
<i>Veronica spicata</i> subsp. <i>crassifolia</i> (Nyman) Hayek	Southern Carpathians, northern part of the Balkan Peninsula	Walters & Webb 1972; Oprea 2005
<i>Waldsteinia teppneri</i> Májovský	Eastern Alps, Western Carpathians (rare)	Kliment 1999

Notes

²⁵⁹ According to Vonica (2014 in litt.), *Centaurea degeniana* is a hybrid of *C. macroptilon* subsp. *oxylepis* and some other taxon of the *Centaurea* genus (cf. Koutecký 2008).

²⁶⁰ *Chrysanthemum zawadzki* is evaluated as Eurasian continental species distributed from the Carpathians to the eastern Asia, with a discontinuous distribution in Europe (Zarzycki 1976, Holub 1999, Wróbel 2008, Szelağ & Kobiv 2014). At the same time it is a very polymorphic species forming a serie of specific individual populations along a longitudinal gradient, which differ mainly by the colour of ligular flowers and shape of leaves (Holub 1981). Morphologic differences stimulated description of several subspecies (for details see Zarzycki 1976, Greuter in Euro+Med, and others). A relic exclave distribution in the Pieniny Mts (*locus classicus* of the species and the only occurrence in the Carpathians) is far remote (by more than 1 000 km; Zarzycki 2000) not only from the margin of its continuous Siberian ditribution but also from its distribution in the European part of Russia. The isolated population from the Pieniny Mts has been considered to be endemic (usually at the level of nominate subspecies or a variety) for a long time since the study by Pax (1898) was published (see the summary by Kliment 1999). On the other hand, Zarzycki (1982) considered it to be a late-glacial relic, reaching the Pieniny Mts in Late Pleistocene during maximum development of the pine and larch-pine forests. The species was ordered to relics also by Holub (1981, 1987, 1999), who considered it as species of late-glacial steppes expanding to Europe from Siberia along the margin of a continental glacial sheet. Domin (1934) evaluated it as an old relic species with a large disjunctive distribution, which should be distinguished from true endemics. The endemic status of the Pieniny population was questioned already by Piękoś (1971); it is also not listed in the recent overview of endemic plants of Poland (Mirek & Piękoś-Mirkowa 2009).

²⁶¹ *Hieracium catenatum* is a central-European species up to now known only from Ukraine and Belarus (Sennikov 1995, 1999b) with presumed wider distribution towards the west (Sennikov 2014 in litt.).

²⁶² In the included literature *Jovibarba globifera* subsp. *glabrescens* (Sabr.) Holub is reported using an invalid name *Jovibarba hirta* subsp. *glabrescens* (Sabr.) Soó et Jáv.

²⁶³ *Rhinanthus gracilis* is a Balkan-Carpathian species with a contradictory taxonomic evaluation. Although it is accepted e.g. in the Euro+Med database, the Romanian authors by themselves evaluate it differently. E.g. Popescu & Sanda (1998) reported it as *Rhinanthus* × *gracilis* Schur (*R. alpinus* × *R. serotinus*), Oprea (2005) as a synonym of *R. alectorolophus* (Scop.) Pollich. Soó & Webb (1972) consider it to be transitional type between *R. alpinus* Baumg. and *R. angustifolius* C. C. Gmelin.

References

- Alexiou S. (2013): The genus *Colchicum* L. (Colchicaceae) in Greece. – Parnassiana Arch. 1: 59–73.
- Ančev M. (1989): *Satureja* L. – In: Kuzmanov B. (ed.), Flora na Narodna republika Bălgarija, p. 334–343. Izdatelstvo na Bălgarskata akademija na naukite, Sofija.
- Assyov B. & Petrova A. (eds) (2006): Konspekt na visšata flora na Bălgarija. Chorologija i florni elementi / Conspectus of the Bulgarian vascular flora. Distribution maps and floristic elements. Ed. 3. – Bulgarian biodiversity foundation, Sofia.
- Assyov B. & Vassilev R. (2004): New chorological data and remarks on the distribution of some vascular plants in Bulgaria. – Phytol. Balc. 10: 191–199.
- Ball P. W. (1972): *Melittis* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 143. Cambridge University Press, Cambridge.
- Ball P. W. & Getliffe F. M. (1972): *Satureja* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 163–165. Cambridge University Press, Cambridge.
- Beldie A. (1977): Flora României. Determinator ilustrat al plantelor vasculare I [Flora of Romania. Illustrated identification key of vascular plants I]. – Edit. Acad. Rep. Soc. România, Bucureşti.
- Borza T., Groza Gh., Dragoş N. & Coman N. (1998): Morphological and genetic variation in some populations of the endemic *Erythronium dens-canis* var. *niveum* and the widespread *Erythronium dens-canis* var. *dens-canis*. – Contr. Bot. 1997–1998: 39–50.
- Browicz K. (ed.), Gostyńska-Jakuszevska M. & Kaczmarek Cz. (1971): Atlas rozmieszczenia drzew i krzewów w Polsce / Atlas of distribution of trees and shrubs in Poland / Atlas rozmieszczenia drzew i krzewów w Polsce. 10. – Zakład dendrologii i Arboretum Kórnickie PAN, Poznań.
- Căprar M., Cantor M., Szatmari P. & Sicora C. (2014): *Rhododendron ferrugineum* L. and *Rhododendron myrtifolium* Schott & Kotschy in habitats from Eastern Alps mountains and Carpathian Mountains. – J. Hort. Forest. & Biotechnol. 18: 123–130.
- Chater A. O., Walters S. M., Akeyrod J. K. & Wrigley F. (1993): *Silene* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 191–218. Cambridge University Press, Cambridge.

- Chrtěk J. jr. (2004): *Hieracium* L. – In: Štěpánek J. & Štěpánková J. (eds), Květena České republiky 7, p. 540–701. Academia, Praha.
- Chrtěk J. jr. & Mráz P. (2007): Taxonomic revision of *Hieracium nigrescens* agg. in the Western Carpathians. – *Preslia* 79: 45–62.
- Chrtěková A. (1988): *Coronilla* L. – In: Bertová L. (ed.), Flóra Slovenska IV/4, p. 358–367. Veda, Bratislava.
- Cincović T. (1972): *Trifolium* L. – In: Josifović M. (ed.), Flora SR Srbije 4, p. 424–471. Srpska Akademija nauka i umetnosti, Beograd.
- Ciocârlan V. (2009): Flora ilustrată a României. Pteridophyta et Spermatophyta [Illustrated flora of Romania. Pteridophyta et Spermatophyta]. – Editura Ceres, București.
- Clapham A. R. & Akeyrod J. R. (1993): *Thlaspi* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 384–388. Cambridge University Press, Cambridge.
- Csiký J., Farkas S., Király G., Pál R., Purger D. & Tóth I. Z. (2005): A *Cirsium boujartii* (Pill. et Mitterp.) Schultz Bip. újrafelfedezése Magyarországon [Rediscovery of *Cirsium boujartii* (Pill. et Mitterp.) Schultz Bip. in Hungary]. – *Fl. Pannon.* 3: 69–77.
- Čornej I. I. (2011): Krytyčna revizija taksoniv, ščo navodylis' jak endemični dlja flory Ukraïns'kych Karpat [Critical revision of taxa reported as endemic to flora of the Ukrainian Carpathians]. – *Nauk. Zap. Bukovins'k. Tovar. Pryr.* 1: 23–59.
- Dakskobler I., Vreš B., Seliškar A., Kobal M. & Sinjur I. (2010): *Scorzoneroïdes crocea* (Haenke) Holub, a new species in the flora of Slovenia and the Southeastern Alps. – *Wulfenia* 17: 59–75.
- Delipavlov D. (1971): Einige für die bulgarische Flora neue und seltene Arten. – *Feddes Repert.* 81: 713–719.
- Dihoru G. & Negrean G. (2009): Cartea roșie a plantelor vasculare din România [Red data book of vascular plants of Romania]. – Edit. Acad. Române, București.
- Diklić N. & Nikolić V. (1986b): *Erythronium dens-canis* f. *niveum* Baumg. – In: Sarić R. M. & Diklić N. (eds), Flora SR Srbije 10. Dodatak (2), p. 220. Srpska Akademija Nauka i Umetnosti, Beograd.
- Domin K. (1934): Vegetační poměry slovenské přírodní rezervace v Pieninách. – *Bratislava* 8: 177–185.
- Dostál J. (1976): *Centaurea* L. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 254–301. Cambridge University Press, Cambridge.
- Edmondson J. R. (1980): *Poa* L. – In: Tutin T. G. et al. (eds), Flora Europaea 5, p. 159–167. Cambridge University Press, Cambridge.
- Ehrendorfer F. & Krendl F. (1976): *Galium*. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 14–36. Cambridge University Press, Cambridge.
- Fedorov A. A. & Kovanda M. (1976): *Campanula* L. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 74–93. Cambridge University Press, Cambridge.
- Ferguson I. K. (1972a): *Androsace* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 20–23. Cambridge University Press, Cambridge.
- Ferguson I. K. (1972b): *Verbascum* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 205–216. Cambridge University Press, Cambridge.
- Finch R. A. & Sell P. D. (1976): *Leontodon* L. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 310–315. Cambridge University Press, Cambridge.
- Gajić M. (1975): Asteraceae. – In: Josifović M. (ed.), Flora SR Srbije 7, p. 1–465. Srpska akademija nauka i umetnosti, Beograd.
- Ghișa E. & Beldie A. (1972): *Poa* L. – In: Nyárády E. I. et al. (eds), Flora Republicii Socialiste România 12, p. 364–429. Edit. Acad. Rep. Soc. România, București.
- Goliašová K., Michalková E. & Mráz P. (2008): *Campanula* L. – In: Goliašová K. & Šípošová H. (eds), Flóra Slovenska VI/1, p. 239–349. Veda, Bratislava.
- Gușuleac M. (1960): *Pulmonaria* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 7, p. 254–271, 644. Edit. Acad. Rep. Pop. Romîne, București.
- Holub J. (1980): *Avenula* (Dumort.) Dumort. – In: Tutin T. G. et al. (eds), Flora Europaea 5, p. 210–216. Cambridge University Press, Cambridge.
- Holub J. (1981): Ochrana fytogenofondu z hlediska taxonomického a fytogeografického [Protection of the floristic diversity from the aspect of taxonomy and phytogeography]. – *Stud. Českoslov. Akad. Věd* 20: 27–39.
- Holub J. (1987): K fytogeografické analýze československé květeny [On the phytogeographical analysis of Czechoslovak flora]. – *Zprávy Českoslov. Bot. Společn.* 22, Mater. 6: 15–33.
- Holub J. (1999): *Dendranthema zawadzki* (Herb.) Cvelev. – In: Čerovský J. et al. (eds), Červená kniha vzácných a ohrožených druhů vyšších rostlin SR a ČR 5. Vyššie rastliny, p. 127. Příroda, Bratislava.
- Hrouda L. (2010a): *Gagea* Salisb. – In: Chrtěk J. jr., Kaplan Z. & Štěpánková J. (eds), Květena České republiky 8, p. 403–418. Academia, Praha.
- Hrouda L. (2010b): *Ornithogalum* L. – In: Chrtěk J. jr., Kaplan Z. & Štěpánková J. (eds), Květena České republiky 8, p. 600–613. Academia, Praha.
- Jalas J. & Suominen J. (eds) (1989): Atlas Florae Europaeae 8. Nymphaeaceae to Ranunculaceae. – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J. & Suominen J. (eds) (1994): Atlas Florae Europaeae 10. Cruciferae (*Sisymbrium* to *Aubrieta*). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.

- Jalas J., Suominen J. & Lampinen R. (eds) (1996): Atlas Florae Europaeae 11. Cruciferae (*Ricotia* to *Raphanus*). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J., Suominen J., Lampinen R. & Kurtto A. (eds) (1999): Atlas Florae Europaeae 12. Resedeaceae to Platanaceae. – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Kliment J. (1999): Komentovaný prehľad vyšších rastlín flóry Slovenska, uvádzaných v literatúre ako endemické taxóny [Annotated survey of the vascular plants of Slovak flora recorded in literature as endemic taxa]. – Bull. Slov. Bot. Spoločn. 21, Suppl. 4: 1–434.
- Kobiv J., Prokopiv A., Heleš M., Borsukevyč L. & Nadraha M. (2007a): Pošyrennja i stan populacij ridkisnych, zahroženych ta endemičnych vydiv roslyn u pivničnij častyni prikordonnoï diljanky Čyvčyns'kyh hir (Ukraïns'ki Karpaty) [Distribution and state of populations of rare, endangered and endemic species of plants in the northern part of frontier area in the Chyvchyny Mountains (Ukrainian Carpathians)]. – Visnyk L'viv. Univ., Ser. Biol. 45: 71–84.
- Koutecký P. (2008): Taxonomická studie středoevropských zástupců *Centaurea* sect. *Jacea* [Taxonomic study of Central European taxa of *Centaurea* sect. *Jacea*]. – PhD thesis, University of South Bohemia, České Budějovice.
- Kovačić S. (2004): The genus *Campanula* L. (Campanulaceae) in Croatia, circum-Adriatic and west Balkan region. – Acta Bot. Croat. 63: 171–202.
- Kubát K. (2010): *Dactylorhiza* Nevski. – In: Chrtěk J. jr., Kaplan Z. & Štěpánková J. (eds), Květena České republiky 8, p. 502–523. Academia, Praha.
- Kucowa I. (1967): *Galium* L. – In: Pawłowski B. (ed.), Flora Polska 11, p. 292–324. Państw. Wydaw. Nauk., Warszawa.
- Kuľak A. & Sender J. (2011): Park-and-palace complexes as biodiversity refuges on example Galezów ex-manor park. – Teka Kom. Ochr. Środ. Przyr. 8: 73–85.
- Kurtto A., Fröhner S. E. & Lampinen R. (eds) (2007): Atlas Florae Europaeae 14. Rosaceae (*Alchemilla* and *Aphanes*). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Letz R. (1998): Vybrané problémy taxonomickej diferenciácie rodov *Sempervivum* a *Jovibarba* v Európe [Selected problems of taxonomic differentiation of the genera *Sempervivum* and *Jovibarba* in Europe]. – PhD thesis, Faculty of Sciences, Comenius University, Bratislava.
- Letz D. R. (2012): *Minuartia* L. – In: Goliašová K. & Michalková E. (eds), Flóra Slovenska VI/3, p. 140–190. Veda, Bratislava.
- Lipschitz S. J. (1976): *Saussurea* DC. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 216–217. Cambridge University Press, Cambridge.
- Marhold K. (1993): *Melittis* L. – In: Bertová L. & Goliašová K. (eds), Flóra Slovenska V/1, p. 221–223. Veda, Bratislava.
- Markgraf-Dannenbergh I. (1980): *Festuca* L. – In: Tutin T. G. et al. (eds), Flora Europaea 5, p. 125–153. Cambridge University Press, Cambridge.
- McNeill J. (1968): *Polygala* L. – In: Tutin T. G. et al. (eds), Flora Europaea 2, p. 231–236. Cambridge University Press, Cambridge.
- Melovski L. & Matevski V. (2008): Development of criteria for the selection of wetland habitats for strict protection – case study of Bunec site in the National park „Mavrovo“. – In: Proceedings of the III Congress of Ecologists of the Republic of Macedonia with International Participation, 06–09. 10. 2007, Struga. Special issues of Macedonian Ecological Society, Vol. 8, p. 157–166.
- Merxmüller B. & Sauer W. (1972): *Pulmonaria* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 100–102. Cambridge University Press, Cambridge.
- Mirek Z. & Piękoś-Mirkowa H. (2009): Fitogeograficzne aspekty endemizmu w Polsce [Phytogeographic aspects of endemism in Poland]. – Wiadom. Bot. 53: 7–30.
- Mustafa B., Hajdari A., Veselaj Z., Beadini N., Ibrahim H., Mustafa N. & Morina I. (2013): Extension of the National park „Sharri“ boundaries, significant action for preservation of natural values. – Nat. Montenegro 12: 607–616.
- Negrean G. & Oltean M. (1989): Endemite și zone endemoconservatoare din Carpații Sud-Estici [Endemics and endemoconservation areas in the South-Eastern Carpathians]. – Ocrot. Nat. 33: 15–25.
- Niketić M. & Tomović G. (2008): Survey of some rare and endangered plants in Serbia with new chorological data. – Bull. Nat. Hist. Mus. 2008, 1: 113–148.
- Nikolić V. & Diklić N. (1986): *Primula halleri* Gmelin. – In: Flora SR Srbije 10. Dodatak (2), p. 76–77. Srpska akademija nauka i umetnosti, Beograd.
- Oprea A. (2005): Lista critică a plantelor vasculare din România [Critical list of vascular plants of Romania]. – Editura Universității „Alexandru Ioan Cuza“, Iași.
- Oprea A. & Sîrbu C. (2013): The vascular flora of Rarău massif (Eastern Carpathian, Romania). Note II. – Mem. Sci. Sect. Romanian Acad., Biol. 36: 18–52.
- Paucă A. & Nyárády E. I. (1960): *Rhinanthus* L. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 7, p. 609–622. Edit. Acad. Rep. Pop. Romîne, București.
- Pax F. (1898): Grundzüge der Pflanzenverbreitung in den Karpathen. 1. – Wilhelm Engelmann, Leipzig.
- Piękoś H. (1971): *Dendranthema* (DC.) Des Moul. – In: Pawłowski B. & Jasiewicz A. (eds), Flora Polska 12, p. 286–288. Państw. Wydawn. Nauk., Warszawa & Kraków.

- Piórecki J. & Zarzycki K. (2008): *Cirsium decussatum* Janka. – In: Mirek Z. & Piękoś-Mirkowa H. (eds), Czerwona księga Karpat Polskich. Rośliny naczyniowe, p. 364–365. Instytut Botaniki im. W. Szafera PAN, Kraków.
- Popescu A. & Sanda V. (1998): Conspectul florei cormofitelor spontane din România [Enumeration of spontaneous vascular plants of Romania]. – Editura Universității din București, București.
- Popova T. N. (1972): *Rhododendron* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 8–9. Cambridge University Press, Cambridge.
- Prokudin J. N. (ed.) (1987): Opredelitel' vysshich rastenij Ukrainy [Identification key of vascular plants of Ukraine]. – Naukova dumka, Kiiv.
- Robson N. K. B. (1968): *Hypericum* L. – In: Tutin T. G. et al. (eds), Flora Europaea 2, p. 261–269. Cambridge University Press, Cambridge.
- Robson N. K. B. (2010): Studies in the genus *Hypericum* L. (Hypericaceae) 5(1). Section 10. *Olympia* to 15/16. *Crossophyllum*. – Phytotaxa 4: 5–126.
- Sârbu I., Ștefan N. & Oprea A. (2013): Plante vasculare din România. Determinator ilustrat de teren [Vascular plants of Romania. An illustrated field guide]. – Victor B Victor, București.
- Schneeweiss G. M. (1998): *Saxifraga carpatica* (Saxifragaceae) – neu für die Alpen. – Phytom (Horn) 38: 301–305.
- Schönschwetter P. & Božo F. (eds) (2013): Montenegro 26. 6.–4. 7. 2013. Exkursionsprotokoll. – Universität Innsbruck, Institut für Botanik. URL: http://www.uibk.ac.at/botany/studies/downloads/vm3-exkursion-mit-uebung_montenegro_26.6.-4.7.2013.pdf.
- Schwarz O. (1968): Beiträge zur Kenntnis der Gattung *Primula*. – Wiss. Zeitschr. Univ. Jena, Sect. Math.-Natur. 17: 307–332.
- Sennikov A. N. (1995): Novye vidy roda *Hieracium* (Asteraceae) iz Vostočnoj Evropy [The new species of the genus *Hieracium* (Asteraceae) from the Eastern Europe]. – Bot. Zhurn. 80: 78–84.
- Sennikov A. N. (1999b): Pervye itogi izučeniya jastrebinok (*Hieracium*, Asteraceae) flory Belorussii [The preliminary results of study of *Hieracium* (Asteraceae) from the Byelorussian flora]. – Bot. Zhurn. 84: 119–128.
- Skalická A. & Skalický V. (1987): Taxonomic and nomenclatoric notes on the threatment of Coniferae for the Flora of Czech socialistic republic. – Novit. Bot. Univ. Carol. 3: 53–62.
- Somlyay L., Pintér I. & Csontos P. (2006): Taxonomic studies of the *Muscari botryoides* complex in Hungary. – Folia Geobot. 41: 213–228.
- Soó R. (1980b): *Dactylorhiza* Necker ex Nevski. – In: Tutin T. G. et al. (eds), Flora Europaea 5, p. 333–337. Cambridge University Press, Cambridge.
- Soó R. & Webb D. A. (1972): *Rhinanthus* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 276–281. Cambridge University Press, Cambridge.
- Syčák N. M. (2002): Ridkysni ta endemični vydy rodu *Alchemilla* L. (Rosaceae) u flori Ukraïns'kych Karpat [Rare and endemic species of genus *Alchemilla* L. (Rosaceae) in the flora of the Ukrainian Carpathians]. – In: Hamor F. D. (ed.), Mountains and people (in the context of sustainable development). Proceedings of the International conference, October 14th–18th, 2002, Vol. 2, p. 473–477. Rakhiv.
- Sýkora T. (1983): Taxonomie a rozšíření bříz okruhu *Betula alba* v Českém masívu [Die Sippen von *Betula alba* agg. und ihre Verbreitung im Böhmischem Massiv]. – Zprávy Českoslov. Bot. Společn. 18: 1–14.
- Szeląg Z. (2011): *Hieracia balcanica* VII. Identity and typification of *Hieracium bohatschianum* (Asteraceae). – Polish Bot. J. 56: 155–157.
- Szeląg Z. & Kobiv J. (2014): Typification of *Chrysanthemum zawadzki* (Asteraceae). – Polish Bot. J. 59: 27–30.
- Šída O. (1996): Taxonomická studie *Erigeron* sect. *Trimorpha* (Cass.) DC. v České a Slovenské republice a přilehlých oblastech [Taxonomic study of *Erigeron* sect. *Trimorpha* (Cass.) DC. in the Czech Republic and Slovak Republic and neighbouring territories]. – Diploma thesis, Department of Botany, Charles University, Praha.
- Šljakov P. N. (1989): *Hieracium* L. – In: Cvelev N. N. (ed.), Flora evropejskoj časti SSSR 8, p. 140–300. Nauka, Leningrad.
- Šourková M. & Hrouda L. (1997): *Bupleurum* L. – In: Chrtěk J. jr. & Tomšovic P. (eds), Květena České republiky 5, p. 322–329. Academia, Praha.
- Španiel S., Marhold K., Filová B. & Zozomová-Lihová J. (2011a): Genetic and morphological variation in diploid-polyploid *Alyssum montanum* in Central Europe: taxonomic and evolutionary considerations. – Pl. Syst. Evol. 294: 1–25.
- Španiel S., Marhold K., Passalacqua N. G., & Zozomová-Lihová J. (2011b): Intricate variation patterns in the diploid-polyploid complex of *Alyssum montanum*–*A. repens* (Brassicaceae) in the Appenine peninsula: Evidence for long-term persistence and diversification. – Amer. J. Bot. 98: 1887–1904.
- Tutin T. G. (1993): *Trollius* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 253. Cambridge University Press, Cambridge.
- Tutin T. G. & Walters S. M. (1993): *Dianthus* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 227–246. Cambridge University Press, Cambridge.
- Walters S. M. & Webb D. A. (1972): *Veronica* L. – In: Tutin T. G. et al. (eds), Flora Europaea 3, p. 242–251. Cambridge University Press, Cambridge.
- Webb D. A. (1993a): *Saxifraga* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 437–458. Cambridge University Press, Cambridge.

- Webb D. A. & Ferguson I. K. (1968): *Geranium* L. – In: Tutin T. G. et al. (eds), *Flora Europaea* 2, p. 193–199. Cambridge University Press, Cambridge.
- Wróbel I. (2008): *Dendranthema zawadzkii* (Herbich) Tzvelev. – In: Mirek Z. & Piękoś-Mirkowa H. (eds), *Czerwona księga Karpat Polskich. Rośliny naczyniowe*, p. 376–377. Instytut Botaniki im. W. Szafera PAN, Kraków.
- Zahariadi C. (1966): *Liliaceae* A. L. Jussieu. – In: Nyárády E. I. (ed.), *Flora Republicii Socialiste România* 11, p. 106–404. Edit. Acad. Rep. Soc. România, București.
- Zarzycki K. (1976): Małe populacje pienińskich roślin reliktowych i endemicznych, ich zagrożenie i problemy ochrony. – *Ochr. Przyr.* 41: 7–75.
- Zarzycki K. (1982): Rośliny wyższe (kwiatowe i paprotniki). Rośliny rodzime [Vascular plants (flowering plants and ferns). Autochthonous plants]. – In: Zarzycki K. (ed.), *Przyroda Pienin w obliczu zmian*, p. 127–142. Państw. Wydaw. Nauk., Warszawa.
- Zarzycki K. (2000): Paprotniki i rośliny kwiatowe (rośliny naczyniowe). – In: Razowski J. (ed.), *Flora i fauna Pienin. Monografie Pienińskie* 1: 75–79.

Electronic Appendix 4. – Species, subspecies and varieties included to taxa with a wider non-endemic distribution

Taxon	Included in	References
<i>Achillea carpatica</i> Blocki ex Dubovik	<i>Achillea distans</i> Waldst. et Kit. ex Willd.	Danihelka 2013 in litt.
<i>Acinos alpinus</i> subsp. <i>baumgartenii</i> (Simonk.) Pawł. (Syn.: <i>A. baumgartenii</i> (Simonk.) Klokov)	<i>Acinos alpinus</i> (L.) Moench subsp. <i>alpinus</i> (syn.: <i>Calamintha alpina</i> (L.) Lam. subsp. <i>alpina</i>)	Popescu & Sanda 1998; Oprea 2005; Ciocârlan 2009; Sârbu et al. 2013
<i>Aconitum jacquinii</i> Rchb. (Syn.: <i>A. anthora</i> subsp. <i>jacquinii</i> (Rchb.) Domin)	<i>Aconitum anthora</i> L.	Novikoff & Mitka 2011a, b
<i>Alchemilla intermedia</i> subsp. <i>sooi</i> (Palitz) Soó (Syn.: <i>A. helvetica</i> subsp. <i>sooi</i> (Palitz) Dihoru et Pârvu)	<i>Alchemilla monticola</i> Opiz	Kurtto et al. 2007
<i>Alchemilla pungentiflora</i> (Plocek) Plocek	<i>Alchemilla straminea</i> Buser	Kurtto et al. 2007
<i>Alchemilla subconnivens</i> Pawł. ²⁶⁴	<i>Alchemilla connivens</i> Buser	Fröhner 1975; Kurtto et al. 2007
<i>Alchemilla tatrica</i> Pawł.	<i>Alchemilla subcrenata</i> Buser	Plocek 1992; Kurtto et al. 2007
<i>Alyssoides utriculata</i> var. <i>micrantha</i> Ciocârlan	<i>Alyssoides utriculata</i> (L.) Medik.	Dihoru & Negrean 2009
<i>Alyssum montanum</i> subsp. <i>brymii</i> (Dostál) Soó	<i>Alyssum gmelinii</i> Jord. et Fourr.	Španiel et al. 2012; Španiel 2014 in litt.
<i>Astragalus australis</i> var. <i>bucsecsii</i> (Jáv.) Guşul. ²⁶⁵ (Syn.: <i>A. australis</i> proles <i>bucsecsii</i> Jáv.)	<i>Astragalus australis</i> (L.) Lam.	Jávorka 1916; Podlech 2011
<i>Biscutella laevigata</i> subsp. <i>hungarica</i> Soó (Syn.: <i>B. austriaca</i> subsp. <i>hungarica</i> (Soó) Peniaşteková)	<i>Biscutella laevigata</i> subsp. <i>austriaca</i> (Jord.) Mach.-Laur.	Jalas et al. 1996; Peniaşteková 2002
<i>Bupleurum subfalcatum</i> Schur ²⁶⁶ (Syn.: <i>B. exaltatum</i> Schur)	<i>Bupleurum falcatum</i> subsp. <i>cernuum</i> (Ten.) Arcang.	Todor 1958; Čopyk 1976; Malynovs'kyj et al. 2002; Oprea 2005; Abduloeva & Fedorončuk 2006; Čornej 2011
<i>Campanula patula</i> subsp. <i>peterfii</i> (Soó) Soó ²⁶⁷	<i>Campanula patula</i> L. subsp. <i>patula</i>	Ghişa et al. 1964; The Plant List
<i>Cardamine marholdii</i> Tzvelev ²⁶⁸	<i>Cardamine pratensis</i> L.	Cvelev 2003; Marhold 2014 in litt.
<i>Carduus bicolorifolius</i> Klokov	<i>Carduus personata</i> subsp. <i>albidus</i> (Adamovič) Kazmi	Oprea 2005; Čornej 2011; Greuter in Euro+Med
<i>Centaurea phrygia</i> subsp. <i>nigriceps</i> (Dobroc. z.) Dostál	<i>Centaurea phrygia</i> L.	Koutecký 2014 in litt.
<i>Centaurea coziensis</i> Nyár. (Syn.: <i>C. stoebe</i> var. <i>coziensis</i> (Nyár.) Soó)	<i>Centaurea stoebe</i> L.	Beldie & Alexandrescu 1976; Morariu & Beldie 1976; Mráz 2014 in litt.
<i>Centaurea ratezatensis</i> Prodan ²⁶⁹ (Syn.: <i>C. phrygia</i> subsp. <i>ratezatensis</i> (Prodan) Dostál; <i>C. pseudophrygia</i> subsp. <i>ratezatensis</i> (Prodan) Soó)	<i>Centaurea stenolepis</i> Kern.	Vonica et al. 2013; Koutecký 2014 in litt.
<i>Cerastium arvense</i> subsp. <i>calcicolum</i> (Schur) Borza	<i>Cerastium arvense</i> L.	Letz & Michalková 2012
<i>Cerithe glabra</i> subsp. <i>tatrica</i> Hadač	<i>Cerithe glabra</i> Mill. subsp. <i>glabra</i>	Valdés in Euro+Med; cf. Kliment 1999
<i>Chenopodium wolffii</i> Simonk.	<i>Oxybasis glauca</i> (L.) S. Fuentes et al. (Syn.: <i>Chenopodium glaucum</i> L.)	Jalas & Suominen 1980; Ciocârlan 2009; Uotila in Euro+Med
<i>Dianthus carthusianorum</i> subsp. <i>saxigenus</i> (Schur) Dostál (Syn.: <i>D. saxigenus</i> Schur)	<i>Dianthus carthusianorum</i> L.	Jalas & Suominen 1986
<i>Dianthus simonkaianus</i> Péterfi (Syn.: <i>D. petraeus</i> subsp. <i>simonkaianus</i> (Péterfi) Tutin)	<i>Dianthus petraeus</i> subsp. <i>orbeticus</i> (Velen.) Greuter et Burdet	Oprea 2005; Ciocârlan 2009; Dihoru & Negrean 2009; Sârbu et al. 2013; Marhold in Euro+Med
<i>Draba aizoides</i> subsp. <i>zmudae</i> Zapal.	<i>Draba aizoides</i> L. subsp. <i>aizoides</i>	Jalas et al. 1996; Peniaşteková & Kliment 2002
<i>Edraianthus kitaibelii</i> (A. DC.) A. DC.	<i>Edraianthus graminifolius</i> (L.) A. DC.	Beldie & Alexandrescu 1976; Beldie 1979; Popescu & Sanda 1998; Oprea 2005; Sârbu et al. 2013
<i>Festuca aglochis</i> Borbás	<i>Festuca supina</i> Schur	Futák 1971; Kliment 1999; Piękoś-Mirkowa & Mirek 2003
<i>Galium bellatulum</i> Klokov	<i>Galium anisophyllum</i> Vill. ²⁷⁰	Ehrendorfer et al. 1996; Marhold et al. 2007
<i>Galium carpaticum</i> Klokov	<i>Galium polonicum</i> Blocki ²⁷¹	Ehrendorfer & Krendl 1976; Malynovs'kyj et al. 2002

Taxon	Included in	References
<i>Galium pawlowskii</i> Kucova ²⁷²	<i>Galium saxatile</i> L.	Ehrendorfer & Krendl 1976; Mirek & Piękoś-Mirkowa 1984; Marhold in Euro+Med
<i>Geranium alpestre</i> Schur	<i>Geranium sylvaticum</i> L.	Kliment 1999; Oprea 2005; Aedo in Euro+Med
<i>Heliosperma quadrifidum</i> subsp. <i>carpaticum</i> Zapal. ²⁷³ (Syn.: <i>H. carpaticum</i> (Zapal.) Klokov; <i>Ixoca carpatica</i> (Zapal.) Ikonn.)	<i>Heliosperma pusillum</i> (Waldst. et Kit.) Rchb.	Jalas & Suominen 1986; Čornej 2011
<i>Hesperis rominacea</i> F. Dvořák	<i>Hesperis dinarica</i> Beck	Jalas & Suominen 1994; Negrean 2011
<i>Hesperis matronalis</i> subsp. <i>nivea</i> (Baumg.) Kulcz. (Syn.: <i>H. nivea</i> Baumg.)	<i>Hesperis matronalis</i> subsp. <i>candida</i> (Kit. ex Schulzer, Kanitz et Knapp) Hegi et E. Schmidt	Dvořák 1968; Jalas & Suominen 1994; Zahradníková et al. 2002
<i>Hesperis moniliformis</i> Schur (Syn.: <i>H. matronalis</i> subsp. <i>moniliformis</i> (Schur) Borza)	<i>Hesperis matronalis</i> subsp. <i>candida</i> (Kit. ex Schulzer, Kanitz et Knapp) Hegi et E. Schmidt	Ball 1993; cf. Dvořák 1968
<i>Hieracium berdoense</i> Wol.	<i>Pilosella guthnickiana</i> (Hegetschw.) Soják	Bräutigam & Greuter in Euro+Med
<i>Hieracium melananthum</i> (Nägeli et Peter) P. D. Sell et C. West (Syn.: <i>H. chondrillifolium</i> subsp. <i>melananthum</i> (Nägeli et Peter) Zahn)	<i>Hieracium chondrillifolium</i> Fr.	Mirek et al. 2002; Chrtěk jr. 2014 in litt.; Szeląg 2014 in litt.
<i>Hieracium roxolanicum</i> Rehm.	<i>Pilosella guthnickiana</i> (Hegetschw.) Soják	Bräutigam & Greuter in Euro+Med
<i>Hypericum richeri</i> subsp. <i>transsilvanicum</i> (Čelak.) Ciocârlan ²⁷⁴ (Syn.: <i>H. transsilvanicum</i> Čelak.; <i>H. alpinum</i> Kit.)	<i>Hypericum richeri</i> subsp. <i>grisebachii</i> (Boiss.) Nyman	Robson 1968, 2010
<i>Iris dacica</i> Beldie (Syn.: <i>I. hungarica</i> subsp. <i>dacica</i> (Beldie) Prodan et Nyár.)	<i>Iris aphylla</i> L.	Webb & Chater 1980; Popescu & Sanda 1998; Oprea 2005; Ciocârlan 2009; Sârbu et al. 2013
<i>Leontodon gutzlourum</i> V. N. Vassil.	<i>Scorzoneroideis autumnalis</i> subsp. <i>borealis</i> (Ball) Greuter	Greuter in Euro+Med
<i>Leontodon schischkinii</i> V. N. Vassil.	<i>Leontodon hispidus</i> subsp. <i>hastilis</i> (L.) Corb.	Hel'tman 1989; Greuter in Euro+Med
<i>Leucanthemum subalpinum</i> (Schur) Tzvelev	<i>Leucanthemum gaudinii</i> Dalla Torre	Greuter in Euro+Med; The Plant List
<i>Libanotis humilis</i> Schur (Syn.: <i>L. pyrenaica</i> subsp. <i>humilis</i> (Schur) Soó)	<i>Seseli libanotis</i> (L.) W. D. J. Koch	Čornej 2011; The Plant List
<i>Minuartia graminifolia</i> subsp. <i>hungarica</i> Jáv.	<i>Minuartia graminifolia</i> (Ard.) Jáv. subsp. <i>graminifolia</i>	Jalas & Suominen 1983; Halliday 1993; Oprea 2005; Dihoru & Negrean 2009; Sârbu et al. 2013
<i>Muscari carpaticum</i> Racib.	<i>Muscari botryoides</i> (L.) Mill.	Davis & Stuart 1980; Prokudin 1987
<i>Onobrychis transsilvanica</i> Simonk. (Syn.: <i>O. montana</i> subsp. <i>transsilvanica</i> (Simonk.) Jáv.)	<i>Onobrychis montana</i> DC. subsp. <i>montana</i>	Beldie & Alexandrescu 1976; Oprea 2005
<i>Oxytropis montana</i> subsp. <i>retezatensis</i> Pawl.	<i>Oxytropis neglecta</i> J. Gay ex Ten.	Beldie & Alexandrescu 1976; Ciocârlan 2009; Dihoru & Negrean 2009; Sârbu et al. 2013
<i>Pinus nigra</i> var. <i>banatica</i> Georgescu et Ionescu ²⁷⁵ (Syn.: <i>P. banatica</i> (Georgescu et Ionescu) Gergescu et Ionescu)	<i>Pinus nigra</i> J. F. Arnold subsp. <i>nigra</i>	Businský 2008; Raab-Straube in Euro+Med; World Checklist; Businský 2014 in litt.
<i>Ranunculus pseudobulbosus</i> Schur ²⁷⁶	<i>Ranunculus sardous</i> Crantz	Tutin & Cook 1964; The Plant List
<i>Ribes carpaticum</i> Kit.	<i>Ribes petraeum</i> Wulfen	Jalas et al. 1999; Sennikov 2001; Oprea 2005
<i>Ribes heteromorphum</i> Topa	<i>Ribes spicatum</i> E. Robson	Webb 1993b; Jalas et al. 1999; Oprea 2005; Ciocârlan 2009; Dihoru & Negrean 2009; Sârbu et al. 2013
<i>Silene carpatica</i> (Zapal.) Czopik ²⁷⁷ (Syn.: <i>Oberna carpatica</i> (Zapal.) Czerep.; <i>O. behen</i> subsp. <i>carpatica</i> (Zapal.) Tzvelev)	<i>Silene vulgaris</i> (Moench) Garcke	Klokov 1952; Jalas & Suominen 1986; Cvelev 2004b
<i>Silene donetzica</i> subsp. <i>sillingeri</i> (Hendrych) Šourková ²⁷⁸	<i>Silene donetzica</i> Kleopow	Mered'a et al. 2012; cf. Chater et al. 1993
<i>Thalictrum transsilvanicum</i> Schur	<i>Thalictrum minus</i> L.	Nyárady 1953; Diduch et al. 2004b; The Plant List
<i>Thlaspi pawlowskii</i> Dvořáková	<i>Nocca kovatsii</i> (Heuff.) F. K. Mey.	Pyšek et al. 2002; Ciocârlan 2009; Kobiv 2010
<i>Thymus clandestinus</i> Schur	<i>Thymus pulegioides</i> L.	Mártonfi 1997

Taxon	Included in	References
<i>Thymus subalpestris</i> Klokov	<i>Thymus alpestris</i> Tausch ex A. Kern.	Mártonfi 1997
<i>Tragopogon transcarpaticus</i> Klokov	<i>Tragopogon pratensis</i> subsp. <i>orientalis</i> (L.) Čelak.	Richardson 1976; Cvelev 1989; Negru & Ionița 2009

Notes

²⁶⁴ Syčák (2011) accepted *Alchemilla subconnivens* as a separate species.

²⁶⁵ In the studies of the Romanian authors *Astragalus australis* var. *bucsecsii* is usually reported as *A. australis* subsp. *bucsecsii* Jáv. and evaluated as endemic to the Romanian Carpathians occurring in the Ceahlău and Bucegi Mts. However, Jávorka (1916) described *Astragalus australis* proles *bucsecsii* without a rank from the Bucegi Mts, and this taxon was ordered to species synonyms by Podlech (2011). Ciocârlan (2009) reports from Romania only *Astragalus australis*; var. (subsp.) *bucsecsii* was not included to species endemic to the Romanian Carpathians neither by Hurdu et al. (2012a, b).

²⁶⁶ *Bupleurum subfalcatum* was described by Schur (1866) from the Rodna Mts (Mt. Ineu), with a synonym *B. exaltatum* Schur. Todor (1958) evaluated it only at the level of a form *Bupleurum falcatum* var. *cernuum* f. *subfalcatum* (Schur) Todor and reported it from the Romanian Eastern and Southern Carpathians. Oprea (2005) and Hand (in Euro+Med) ordered it to synonyms of *B. falcatum* subsp. *cernuum*. Čopyk (1976) and Malynovs'kyj et al. (2002) considered *B. subfalcatum* as endemic to the Eastern and Southern Carpathians. According to Čornej (2011) and Abduloeva & Fedorončuk (2006), the records of this species from Ukraine require a confirmation.

²⁶⁷ Soó (1926) described *Campanula patula* subsp. *peterfii* at the level of a variety from the castle hill in the Deva City in Transylvania (cf. Ghișa et al. 1964), later it was upgraded to the level of a subspecies (Soó 1977). Dihoru & Pârvu (1987) reported *C. patula* subsp. *peterfii* as endemic to the Southern Carpathians. Fedorov & Kovanda (1976) report it neither within the characteristics of the species *Campanula patula* L. nor as a synonym of by them accepted subspecies.

²⁶⁸ Cvelev (2003) described *Cardamine marholdii* based on quantitative differences (smaller flowers and leaflets in comparison with *Cardamine pratensis*) from the Ukrainian Carpathians (Horhany Mts: Mt. Perednja); outside this area he reports it also from the mountain ranges of the south-eastern Romania and south-eastern Poland. As a separate species it was accepted by Il'jins'ka & Diduch (2007), Čornej (2011) and Dorofeev (2012). Marhold (2014 in litt.) considers *C. marholdii* to be a synonym of *Cardamine pratensis* L.

²⁶⁹ According to Koutecký (2014 in litt.), *Centaurea ratezatensis* represents a morphotype of the closely related *Centaurea stenolepis*; it can be included within this species or evaluated as its variety (cf. Vonica et al. 2013).

²⁷⁰ According to the current knowledge (Marhold et al. 2007), to *Galium anisophyllum* belong also the diploid West-Carpathian populations reported by invalid names *Galium fatrense* Ehrend. et Šipošová and *G. anisophyllum* subsp. *fatrense* Ehrend. evaluated as endemic to the Western Carpathians (cf. Kliment 1999).

²⁷¹ *Galium polonicum* was evaluated as endemic to the Eastern Carpathians by some Ukrainian authors (e.g. Kricsfalussy 1999, Malynovs'kyj et al. 2002); but better part of its localities are outside the Carpathians (cf. Kucowa 1967, Čornej 2011).

²⁷² *Galium pawlowskii* was described by Kucowa (1962) based on Pawłowski collections from the Čyvčyny Mts (Mt. Stoh, ca 1 500 m) in the Ukrainian Carpathians. Chorney et al. (2008) and Čornej (2011) evaluated it as endemic to this mountain range, Tasenkevyč (2003b) as a paleoendemic. Mirek & Piękoś-Mirkowa (1984) considered *Galium pawlowskii* to be conspecific with *Galium saxatile* L. (cf. Ehrendorfer & Krendl 1976; Marhold in Euro+Med and others) and suggested its evaluation as the variety *G. saxatile* var. *pawlowskii*. As they did not report full quotation of bazionym, the suggested combinaton is invalid.

²⁷³ Zapałowicz (1911) reported *Heliosperma quadridum* subsp. *carpaticum* from the montane and alpine regions of the Tatry Mts, the Ukrainian and Romanian Eastern Carpathians, at about 700–1 950 m a.s.l. Klokov (1952), Ikonnikov (2004) and Tasenkevich (2011) ordered it to taxa endemic to the Western and Eastern Carpathians. All reported localities are part of the *Heliosperma pusillum* distribution area (cf. Frajman 2007, Frajman & Oxelman 2007, Šingliarová & Mráz 2012).

²⁷⁴ In the included literature *Hypericum richeri* subsp. *transsilvanicum* (Čelak.) Ciocârlan is reported as *H. richeri* subsp. *transsilvanicum* (Čelak.) Beldie; *H. richeri* subsp. *transsilvanicum* (Čelak.) Beldie et L. Alex. [nom. inval.].

²⁷⁵ In the Romanian literature, *Pinus nigra* var. *banatica* was reported mainly as *Pinus nigra* subsp. *banatica* (Borbás) Novák. However, Borbás (1886) published only nomen nudum; the name of the variety was validly published only by Georgescu & Ionescu in 1935 (cf. Beldie 1952). Novák (in Klika et al. 1953) published the combination *Pinus nigra* subsp. *banatica* (Georgescu et Ionescu) Novák without full quotation of the basionym, and therefore invalidly.

²⁷⁶ According to Tutin & Cook (1964), *Ranunculus pseudobulbosus* probably represents a subspecies or a variety of the species *Ranunculus sardous* Crantz. Nyárády (1953) reported it as the form *R. bulbosus* f. *pseudobulbosus* (Schur) A. Nyár.

²⁷⁷ Zapałowicz (1911) reported by him described variety *Silene venosa* var. *carpatica* from the highest regions of the Tatry Mts, the Polish, Ukrainian as well as Romanian Eastern Carpathians, hence the area falling under the distribution area of *Silene vulgaris* (cf. Mered'a et al. 2012). Although Jalas & Suominen (1986) included *Silene carpatica* to synonyms of *S. vulgaris*, several Ukrainian authors (e.g. Klovok 1952, Čopyk 1976, Cvelev 2004b), using various names, accepted a separate position of this taxon and consider it to be endemic to the Carpathians (Čopyk 1976, Kricsfalussy 1999).

²⁷⁸ In the included literature *Silene donetzica* subsp. *sillingeri* is reported also by an invalid name *S. sillingeri* (Hendrych) Hendrych.

References

- Abduloeva O. P. & Fedorončuk M. M. (2006): Morfoloho-anatomični osoblyvosti plodiv vydiv rodu *Bupleurum* L. flory Ukraïny [The morphological and anatomical fruit features of species of the genus *Bupleurum* L. within the Ukrainian flora]. – Ukrains'k. Bot. Zhurn. 63: 56–69.
- Ball P. W. (1993): *Hesperis* L. – In: Tutin T. G. et al. (eds), *Flora Europaea* 1. Ed. 2, p. 336–337. Cambridge University Press, Cambridge.
- Beldie A. (1952): Pinaceae Lindl. – In: Săvulescu T. (ed.), *Flora Republicii Populare Române* 1, p. 159–176. Edit. Acad. Rep. Pop. Române, București.
- Beldie A. (1979): *Flora României. Determinator ilustrat al plantelor vasculare II* [Flora of Romania. Illustrated identification key of vascular plants II]. – Edit. Acad. Rep. Soc. România, București.
- Beldie A. & Alexandrescu L. (1976): Modificări de taxonomie sau nomenclatură și sinonimizări la principalii taxonii din vol. I–XII *Flora R. S. R.* [Changes in taxonomy or nomenclature and synonymy of the main taxa in volumes I–XII of the *Flora R. S. R.*]. – In: Beldie A. & Morariu I. (eds), *Flora Republicii Socialiste România* 13, p. 72–96. Edit. Acad. Rep. Soc. România, București.
- Borbás V. (1886): *Pinus nigra* Arn. és var. *Banatica* Endl. [*Pinus nigra* Arn. and var. *banatica* Endl.]. – Erdész. Lapok 25: 338–341.
- Businský R. (2008): The genus *Pinus* L., pines: contribution to knowledge. – Acta Pruhon. 88: 1–126.
- Chater A. O., Walters S. M., Akeyrod J. K. & Wrigley F. (1993): *Silene* L. – In: Tutin T. G. et al. (eds), *Flora Europaea* 1. Ed. 2, p. 191–218. Cambridge University Press, Cambridge.
- Chorney I., Budzhak V. & Tokaryuk A. (2008): Rare, endangered and endemic species of plants of the Chyvchyny / Cívcin Mountains (Carpathians). – Transylv. Rev. Syst. Ecol. Res. 5: 37–44.
- Ciocârlan V. (2009): *Flora ilustrată a României. Pteridophyta et Spermatophyta* [Illustrated flora of Romania. Pteridophyta et Spermatophyta]. – Editura Ceres, București.
- Cvelev N. N. (1989): *Tragopogon* L. – In: Cvelev N. N. (ed.), *Flora evropejskoj časti SSSR* 8, p. 46–56. Nauka, Leningrad.
- Cvelev N. N. (2003): O nekotorych krestocvetnykh (Brassicaceae) Vostočnoj Evropy [De Brassicaceis nonnullis Europae Orientalis]. – Novosti Sist. Vyssh. Rast. 35: 95–108.
- Cvelev N. N. (2004b): *Oberna* Adans. – In: Cvelev N. N. (ed.), *Flora Vostočnoj Evropy* 11, p. 229–233. Tovariščestvo naučnykh izdanij KMK, Moskva & Sankt-Peterburg.
- Čopyk V. I. (1976): Vysokohirna flora Ukraïns'kykh Karpat [High-mountain flora of the Ukrainian Carpathians]. – Naukova dumka, Kyïv.
- Čornej I. I. (2011): Krytyčna revizija taksoniv, ščo navodylis' jak endemični dlja flory Ukraïns'kykh Karpat [Critical revision of taxa reported as endemic to flora of the Ukrainian Carpathians]. – Nauk. Zap. Bukovins'k. Tovar. Pryr. 1: 23–59.
- Davis P. H. & Stuart D. C. (1980): *Muscari* Miller. – In: Tutin T. G. et al. (eds), *Flora Europea* 5, p. 46–49. Cambridge University Press, Cambridge.
- Diduch J. P., Ziman S. M., Korotčenko I. A., Burda P. I., Bulach O. V., Heljuta V. P. & Tychonenko J. J. (2004b): *Thalictrum minus* L. – In: Diduch J. P. (ed.), *Ekoflora Ukraïny* 2, p. 284–285. Fitosociocentr, Kyïv.
- Dihoru G. & Negrean G. (2009): *Cartea roșie a plantelor vasculare din România* [Red data book of vascular plants of Romania]. – Edit. Acad. Române, București.
- Dihoru G. & Pârnu C. (1987): *Plante endemice în flora Românie* [Endemic plants in the flora of Romania]. – Editura Ceres, București.

- Dorofeev V. I. (2012): Brassicaceae Burnett (Cruciferae Juss.). – In: Gel'tman D. V. (ed.), Konspekt flory Vostočnoj Evropy, p. 364–437. Tovariščestvo naučnyh izdanij KMK, Sankt-Peterburg & Moskva.
- Dvořák F. (1968): Rod *Hesperis* v Československu. Taxonomická studie [The genus *Hesperis* in Czechoslovakia. Taxonomic study]. – Biol. Práce 14/3: 1–56.
- Ehrendorfer F. & Krendl F. (1976): *Galium*. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 14–36. Cambridge University Press, Cambridge.
- Ehrendorfer F., Samuel R. & Pinsker W. (1996): Enzyme analysis of genetic variation and relationships in diploid and polyploid taxa of *Galium* (Rubiaceae). – Pl. Syst. Evol. 202: 121–135.
- Fedorov A. A. & Kovanda M. (1976): *Campanula* L. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 74–93. Cambridge University Press, Cambridge.
- Frajman B. (2007): Taxonomy and reticulate phylogeny of *Heliosperma* and related genera (Sileneae, Caryophyllaceae). – Acta Universitatis Upsaliensis. Digital Comprehensive Summaries of Uppsala Dissertations from the Faculty of Science and Technology 328, Uppsala.
- Frajman B. & Oxelman B. (2007): Reticulate phylogenetics and phytogeographical structure of *Heliosperma* (Sileneae, Caryophyllaceae) inferred from chloroplast and nuclear DNA sequences. – Molec. Phylogen. Evol. 43: 140–155.
- Fröhner S. (1975): Kritik an der europäischen *Alchemilla*-Taxonomie. – Feddes Repert. 86: 119–169.
- Futák J. (1971): Endemické rastliny na Slovensku [Endemic plants in Slovakia]. – In: Magic D. (ed.), Zborník prednášok Zjazdu Slovenskej botanickej spoločnosti Tisovec 1970, p. 39–54. Bratislava.
- Ghişa E., Csűrös S., Morariu I. & Guşuleac M. (1964): Campanulaceae A. L. Juss. – In: Nyárády E. I. (ed.), Flora Republicii Populare Romîne 9, p. 52–149. Edit. Acad. Rep. Pop. Romîne, Bucureşti.
- Halliday G. (1993): *Minuartia* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 152–160. Cambridge University Press, Cambridge.
- Hel'tman D. V. (1989): *Leontodon* L. – In: Cvelev N. N. (ed.), Flora evropejskoj časti SSSR 8, p. 27–32. Nauka, Leningrad.
- Hurdu B. I., Puşcaş M., Turtureanu P. D., Niketić M., Coldea G. & Zimmermann N. E. (2012a): Patterns of plant endemism in the Romanian Carpathians (South-Eastern Carpathians). – Contr. Bot. Cluj 47: 25–38.
- Hurdu B. I., Puşcaş M., Turtureanu P. D., Niketić M., Vonica G., Coldea G. (2012b): A critical evaluation of the Carpathian endemic plant taxa list from the Romanian Carpathians. – Contr. Bot. Cluj 47: 39–47.
- Ikonnikov S. S. (2004): *Ixoca* Raf. – In: Cvelev N. N. (ed.), Flora Vostočnoj Evropy 11, p. 226–227. Tovariščestvo naučnyh izdanij KMK, Moskva & Sankt-Peterburg.
- Il'jins'ka A. P. & Diduch J. P. (2007): *Cardamine marholdii* Tzvelev. – In: Diduch J. P. (ed.), Ekoflora Ukraïny 5, p. 404–405. Fitosociocentr, Kyïv.
- Jalas J. & Suominen J. (eds) (1980): Atlas Florae Europaeae 5. Chenopodiaceae to Basellaceae. – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J. & Suominen J. (eds) (1983): Atlas Florae Europaeae 6. Caryophyllaceae (Alsinoideae and Paronychioideae). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J. & Suominen J. (eds) (1986): Atlas Florae Europaeae 7. Caryophyllaceae (Silenioideae). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J. & Suominen J. (eds) (1994): Atlas Florae Europaeae 10. Cruciferae (*Sisymbrium* to *Aubrieta*). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J., Suominen J. & Lampinen R. (eds) (1996): Atlas Florae Europaeae 11. Cruciferae (*Ricotia* to *Raphanus*). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jalas J., Suominen J., Lampinen R. & Kurtto A. (eds) (1999): Atlas Florae Europaeae 12. Resedeaceae to Platanaceae. – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Jávorka S. (1916): Kisseb megjegyzések és újabb adatok. IV. közlemény [Kleinere Bemerkungen und neue floristische Daten. IV.]. – Bot. Közlem. 15: 10–17.
- Klika J., Šiman K., Novák F. A. & Kavka B. (1953): Jehličnaté [Conifers]. – Nakladatelství ČSAV, Praha.
- Kliment J. (1999): Komentovaný prehľad vyšších rastlín flóry Slovenska, uvádzaných v literatúre ako endemické taxóny [Annotated survey of the vascular plants of Slovak flora recorded in literature as endemic taxa]. – Bull. Slov. Bot. Spoločn. 21, Suppl. 4: 1–434.
- Klokov M. V. (1952): Caryophyllaceae Juss. – In: Kotov M. I. (ed.), Flora URSS 4, p. 421–646. Vydavnyctvo Akademii nauk Ukraïns'koï RSR, Kyïv.

- Kobiv J. (2010): Ekolohični osoblyvosti oselyšč ridkisnych vydiv roslyn Ukraïns'kich Karpat [Ecological characteristics of habitats of rare plant species in the Ukrainian Carpathians]. – Ukraïns'k. Bot. Zhurn. 67: 355–372.
- Kricsfalussy V. (1999): Flora and vegetation of the Ukrainian upper Tisa Basin: Aspects of biodiversity conservation. – In: Hamar J. & Sárkány-Kiss A. (eds), The upper Tisa valley. Preparatory proposal for Ramsar site designation and an ecological background Hungarian, Romanian, Slovakian and Ukrainian co-operation. Tiscia monograph series 4, p. 273–292. Szeged.
- Kucowa I. (1962): Gatunki rodzaju *Galium* L. sekcji *Leptogalium* Lange z Polski i ziem ościennych [Species of the genus *Galium* L. section *Leptogalium* Lange found in Poland and neighbouring territories]. – Fragm. Florist. Geobot. 8: 417–442.
- Kucowa I. (1967): *Galium* L. – In: Pawłowski B. (ed.), Flora Polska 11, p. 292–324. Państw. Wydaw. Nauk., Warszawa.
- Kurtto A., Fröhner S. E. & Lampinen R. (eds) (2007): Atlas Florae Europaeae 14. Rosaceae (*Alchemilla* and *Aphanes*). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Letz D. R. & Michalková E. (2012): *Cerastium* L. – In: Goliašová K. & Michalková E. (eds), Flóra Slovenska VI/3, p. 291–388. Veda, Bratislava.
- Malynovs'kyj K., Caryk J., Kyjak V. & Nesteruk J. (2002): Ridkisi, endemični, reliktovi ta pohranyčno-areaľni vydy roslyn Ukraïns'kych Karpat [Rare, endemic, relict and boundary areal plant species of the Ukrainian Carpathians]. – Instytut ekologii Karpat Nacjonal'noï akademii nauk Ukraïny, Lviv.
- Marhold K., Mártonfi P., Mered'a P. & Mráz P. (eds) (2007): Chromosome number survey of the ferns and flowering plants of Slovakia. – Veda, Bratislava.
- Mártonfi P. (1997): Nomenclatural survey of the genus *Thymus* sect. *Serpyllum* from Carpathians and Pannonia. – Thaiszia-J. Bot. 7: 111–181.
- Mered'a P., Eliáš P. jr., Dítě D. & Štrba P. (2012): *Silene* L. – In: Goliašová K. & Michalková E. (eds), Flóra Slovenska VI/3, p. 410–533. Veda, Bratislava.
- Mirek Z. & Piękoś-Mirkowa H. (1984): Distribution and habitats of *Galium saxatile* L. in the Carpathian mountains. – Acta Soc. Bot. Polon. 53: 419–427.
- Mirek Z., Piękoś-Mirkowa H., Zajac A. & Zajac M. (2002): Flowering plants and pteridophytes of Poland. A checklist. – W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.
- Morariu I. & Beldie A. (1976): Endemismeale din Flora R.S.R. [Endemics in the Flora R. S. R.]. – In: Beldie A. & Morariu I. (eds), Flora Republicii Socialiste România 13, p. 97–105. Edit. Acad. Rep. Soc. România, București.
- Negrean G. (2011): Addenda to „Flora Romaniaae“ volumes 1–12. Newly published plants, nomenclature, taxonomy, chorology and commentaries (Part 1). – Kanitzia 18: 89–194.
- Negru A. & Ionița O. (2009): Speciile de *Tragopogon* L. (Asteraceae) din flora Basarabiei [Species of *Tragopogon* L. (Asteraceae) in the flora of Bessarabia]. – Cercet. Ști. 1 (43): 1–5.
- Novikoff A. V. & Mitka J. (2011a): Taxonomy and ecology of the genus *Aconitum* L. in the Ukrainian Carpathians. – Wulfenia 18: 37–61.
- Novikov A. V. & Mitka J. (2011b). Rid *Aconitum* v Ukraïns'kych Karpatach [The genus *Aconitum* in the Ukrainian Carpathians]. – Biol. Stud. 5: 153–172.
- Nyárády A. (1953): Ranunculaceae A. L. Juss. – In: Săvulescu T. (ed.), Flora Republicii Populare Române 2, p. 396–661. Edit. Acad. Rep. Pop. Române, București.
- Oprea A. (2005): Lista critică a plantelor vasculare din România [Critical list of vascular plants of Romania]. – Editura Universității „Alexandru Ioan Cuza“, Iași.
- Peniašteková M. (2002): *Biscutella* L. – In: Goliašová K. & Šipošová H. (eds), Flóra Slovenska V/4, p. 632–642. Veda, Bratislava.
- Peniašteková M. & Kliment J. (2002): *Draba* L. – In: Goliašová K. & Šipošová H. (eds), Flóra Slovenska V/4, p. 500–540. Veda, Bratislava.
- Piękoś-Mirkowa H. & Mirek Z. (2003): Endemic taxa of vascular plants in the Polish Carpathians. – Acta Soc. Bot. Polon. 72: 235–242.
- Plocek A. (1992): *Alchemilla* L. – In: Bertová L. (ed.), Flóra Slovenska IV/3, p. 250–373. Veda, Bratislava.
- Podlech P. (2011): Thesaurus Astragalorum. Index of all taxa described within the genus *Astragalus* L. and other genera but belonging to the genus *Astragalus*. Taxa of the Old World and related taxa of the New World. Last updated version december 2011. – URL: http://www.sysbot.biologie.uni-muenchen.de/de/personen/podlech/thesaurus_astragalus.pdf.
- Popescu A. & Sanda V. (1998): Conspectul florei cormofitelor spontane din România [Enumeration of spontaneous vascular plants of Romania]. – Editura Universității din București, București.
- Prokudin J. N. (ed.) (1987): Opredelitel' vysšich rastenij Ukraïny [Identification key of vascular plants of Ukraine]. – Naukova dumka, Kiïv.
- Pyšek P., Sádlo J. & Mandák B. (2002): Catalogue of alien plants of the Czech Republic. – Preslia 74: 97–186.
- Richardson I. B. K. (1976): *Tragopogon* L. – In: Tutin T. G. et al. (eds), Flora Europaea 4, p. 322–325. Cambridge University Press, Cambridge.
- Robson N. K. B. (1968): *Hypericum* L. – In: Tutin T. G. et al. (eds), Flora Europaea 2, p. 261–269. Cambridge University Press, Cambridge.
- Robson N. K. B. (2010): Studies in the genus *Hypericum* L. (Hypericaceae) 5(1). Section 10. *Olympia* to 15/16. *Crossophyllum*. – Phytotaxa 4: 5–126.

- Sârbu I., Ştefan N. & Oprea A. (2013): Plante vasculare din România. Determinator ilustrat de teren [Vascular plants of Romania. An illustrated field guide]. – Victor B Victor, Bucureşti.
- Sennikov A. N. (2001): Grossulariaceae DC. – In: Cvelev N. N. (ed.), Flora vostočnoj Evropy 10, p. 235–243. Tovariščestvo naučnyh izdanij KMK, Moskva & Sankt-Peterburg.
- Schur F. (1866): Enumeratio Plantarum Transsilvaniae. – Guilielmum Braumüller, Vindobonae.
- Soó R. (1926): Kritikai megjegyzések [Critical notes]. – Bot. Közlem. 23: 146–158.
- Soó R. (1977): Systematisch-nomenklatorischen Bemerkungen über kritische Taxa der mitteleuropäischen Flora. – Acta Bot. Acad. Sci. Hung. 23: 375–392.
- Syčák N. M. (2011): Ključ dlja vyznačennja vydiv rodu *Alchemilla* L. (Rosaceae) flory Ukraïnskych Karpat [Key for determination of the species of genus *Alchemilla* L. (Rosaceae) in the flora of the Ukrainian Carpathians]. – Naukovi osnovy zberežennja biotyčnoï riznomanitnosti 2 (9): 215–228.
- Šingliarová B. & Mráz P. (2012): *Heliosperma* (Rchb.) Rchb.). – In: Goliašová K. & Michalková E. (eds), Flóra Slovenska VI/3, p. 546–550. Veda, Bratislava.
- Španiel S., Marhold K., Thiv M. & Zozomová-Lihová J. (2012): A new circumscription of *Alyssum montanum* subsp. *montanum* and *A. montanum* subsp. *gmelinii* (Brassicaceae) in Central Europe: molecular and morphological evidence. – Bot. J. Linn. Soc. 169: 378–402.
- Tasenkevych L. (2003b): Rozmaïttja flory sudynnyh roslyn v Ukraïns'kyh Karpatach [Vascular plants flora diversity of the Ukrainian Carpathians]. – Pratsi Nauk. Tovar. Schevchenka 12: 147–157.
- Tasenkevich L. (2011): Endemism in the Carpathian flora – a chorological aspect. – In: Zemanek B. (ed.), Geobotanist and Taxonomist. A volume dedicated to Professor Adam Zajac on the 70th anniversary of his birth, p. 157–168. Institute of Botany Jagellonian University, Cracow.
- Todor I. (1958): Umbelliferae Juss. – In: Săvulescu T. (ed.), Flora Republicii Populare Romîne 6, p. 326–660. Edit. Acad. Rep. Pop. Romîne, Bucureşti.
- Tutin T. G. & Cook C. D. K. (1964): *Ranunculus* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1, p. 223–238. Cambridge University Press, Cambridge.
- Vonica G. D., Koutecký P., Cantor M., Hurdu B. I. & Bădăraş S. A. (2013): New consideration regarding the endemic *Centaurea ratezatensis* Prod. from Romanian Carpathians. – Acta Biol. Cracov., Bot. 55, Suppl. 1: 71.
- Webb D. A. (1993b): *Ribes* L. – In: Tutin T. G. et al. (eds), Flora Europaea 1. Ed. 2, p. 461–462. Cambridge University Press, Cambridge.
- Webb D. A. & Chater A. O. (1980): *Iris* L. – In: Tutin T. G. et al. (eds), Flora Europaea 5, p. 87–92. Cambridge University Press, Cambridge.
- Zahradníková K., Peniašteková M. & Ťavoda O. (2002): *Hesperis* L. – In: Goliašová K. & Šípošová H. (eds), Flóra Slovenska V/4, p. 230–251. Veda, Bratislava.
- Zapałowicz H. (1911): Krytyczny przegląd roślinności Galicyi III [Conspectus florae Galiciae criticus III]. – Akademia Umiejętności, Cracoviae.

Electronic Appendix 5. – Hybrids considered as endemic taxa

Taxon	a hybrid formula	References
<i>Aconitum</i> × <i>bartokianum</i> Starm.	<i>A. toxicum</i> × <i>A. variegatum</i>	Starmühler 1999, 2000; Mitka 2001; Ilnicki & Mitka 2011
<i>Aconitum</i> × <i>czarnohorensse</i> (Zapał.) Mitka	<i>A. firmum</i> × <i>A. nanum</i>	Mitka 2003; Novikoff & Mitka 2011a, b
<i>Aconitum</i> × <i>dragulescuanum</i> Mucher	<i>A. degenii</i> × <i>A. toxicum</i>	Mucher 1993; Starmühler 1996a, 1997, 2000; Mitka 2001; Ilnicki & Mitka 2011
<i>Aconitum</i> × <i>gayeri</i> Starm.	<i>A. degenii</i> × <i>A. lasiocarpum</i>	Starmühler 1998b; Mitka 2001, 2003; Ilnicki & Mitka 2011; Mitka & Novikoff 2011; Novikoff & Mitka 2011a, b
<i>Aconitum</i> × <i>nanum</i> (Baumg.) Simonk. (Syn.: <i>A. nanum</i> Baumg., <i>A. tauricum</i> Wulf. subsp. <i>nanum</i> (Baumg.) Gáyer	<i>A. bucovinense</i> × <i>A. firmum</i>	Mitka 2001, 2003; Ilnicki & Mitka 2009; Novikoff & Mitka 2011a, b
<i>Aconitum</i> × <i>pawlowskii</i> Mitka et Starm.	<i>A. lasiocarpum</i> × <i>A. variegatum</i>	Mitka & Starmühler 2000; Starmühler 2002; Mitka 2003; Ilnicki & Mitka 2011; Mitka et al. 2013; Sutkowska et al. 2013
<i>Aconitum firmum</i> nsubsp. <i>fussianum</i> Starm.	<i>A. firmum</i> subsp. <i>firmum</i> × <i>A. firmum</i> subsp. <i>fissuræ</i>	Starmühler 1999, 2000; Mitka 2001, 2002, 2003; Novikov & Mitka 2011b
<i>Aconitum firmum</i> nsubsp. <i>paxii</i> Starm.	<i>A. firmum</i> subsp. <i>maninense</i> × <i>A. firmum</i> subsp. <i>moravicum</i>	Mitka 2003, Mitka et al. 2007; Novikov & Mitka 2011b
<i>Aconitum firmum</i> nsubsp. <i>zapalowiczii</i> Starm.	<i>A. firmum</i> subsp. <i>firmum</i> × <i>A. firmum</i> nsubsp. <i>paxii</i>	Mitka 2003
<i>Aconitum moldavicum</i> nsubsp. <i>porcii</i> Starm.	<i>A. moldavicum</i> subsp. <i>moldavicum</i> × <i>A. moldavicum</i> subsp. <i>simonkaianum</i>	Starmühler 1998a; Mitka 2008; Mitka & Novikoff 2011; Novikoff & Mitka 2011a, b
<i>Aconitum toxicum</i> nsubsp. <i>nyaradyanum</i> Mucher	<i>A. toxicum</i> subsp. <i>crispulum</i> × <i>A. toxicum</i> subsp. <i>toxicum</i>	Mucher 1993; Starmühler 1997
<i>Aconitum toxicum</i> nsubsp. <i>ungarianum</i> Starm.	<i>A. toxicum</i> subsp. <i>bucegiense</i> × <i>A. toxicum</i> subsp. <i>toxicum</i>	Starmühler 2000
<i>Centaurea</i> × <i>melanocalathia</i> Borbás ex Czako (Syn.: <i>C. phrygia</i> subsp. <i>melanocalathia</i> (Borbás ex Czako) Dostál; <i>Jacea phrygia</i> subsp. <i>melanocalathia</i> (Borbás ex Czako) Soják)	<i>C. erdneri</i> × <i>C. jacea</i>	Koutecký et al. 2012
<i>Hieracium</i> × <i>grofæ</i> Woł.	<i>H. alpinum</i> × <i>H. umbellatum</i>	Chrték jr. et al. 2006
<i>Hieracium</i> × <i>krasanii</i> Woł. ²⁷⁹	<i>H. alpinum</i> × <i>H. transsilvanicum</i>	Mráz et al. 2005, 2011b
<i>Rosa</i> × <i>heterostyla</i> Chrshan. ²⁸⁰	<i>R. canina</i> × <i>R. dumalis</i>	Buzunova 2001; Kurtto et al. 2004
<i>Senecio</i> × <i>dominii</i> Hodálová	<i>S. germanicus</i> × <i>S. ucranicus</i>	Hodálová 1999a

Notes

²⁷⁹ Along with the diploid primary hybrids the occurrence of apomictic polyploid individuals of *Hieracium krasanii* was also sporadically recorded from Romania (Bistrița Mts; Chrték jr. 2014 in litt.).

²⁸⁰ According to Kurtto et al. (2004), the holotypus *Rosa heterostyla* Chrshan. was identified as the hybrid of *Rosa canina* × *R. dumalis* (cf. Buzunova 2001).

References

- Buzunova I. O. (2001): *Rosa* L. – In: Cvelev N. N. (ed.), *Flora Vostočnoj Evropy* 10, p. 329–361. Tovariščestvo naučnych izdanij KMK, Moskva & Sankt-Peterburg.
- Chrték J. jr., Mráz P. & Sennikov A. N. (2006): *Hieracium* × *grofæ* – a rediscovered diploid hybrid from the Ukrainian Carpathians. – *Biologia* 61: 365–374.
- Hodálová I. (1999a): Multivariate analysis of the *Senecio nemorensis* group (Compositae) in the Carpathians with a new species from the East Carpathians. – *Folia Geobot.* 34: 321–335.
- Ilnicki T. & Mitka J. (2009): Chromosome numbers in *Aconitum* sect. *Aconitum* (Ranunculaceae) from the Carpathians. – *Caryologia* 62: 198–203.
- Ilnicki T. & Mitka J. (2011): Chromosome numbers in *Aconitum* sect. *Cammarum* (Ranunculaceae) from the Carpathians. – *Caryologia* 64: 446–452.
- Koutecký P., Štěpánek J. & Baďurová T. (2012): Differentiation between diploid and tetraploid *Centaurea phrygia*: mating barriers, morphology and geographic distribution. – *Preslia* 84: 1–32.

- Kurtto A., Lampinen R. & Junikka L. (eds) (2004): Atlas Florae Europaeae 13. Rosaceae (*Spiraea* to *Fragaria*, excl. *Rubus*). – The Committee for mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- Mitka J. (2001). Systematyka *Aconitum* subgen. *Aconitum* w Karpatach Wschodnich – wstępne wyniki badań [Systematics of *Aconitum* subgen. *Aconitum* in the Eastern Carpathians – preliminary results]. – Roczn. Bieszcz. 9 (2000): 79–116.
- Mitka J. (2002): Phenetic and geographic pattern of *Aconitum* sect. *Napellus* (Ranunculaceae) in the Eastern Carpathians – a numerical approach. – Acta Soc. Bot. Polon. 71: 35–48.
- Mitka J. (2003): The genus *Aconitum* L. (Ranunculaceae) in Poland and adjacent countries. A phenetic-geographic study. – Institute of botany of the Jagiellonian University, Kraków.
- Mitka J. (2008): *Aconitum moldavicum* Hacq. (Ranunculaceae) and its hybrids in the Carpathians and adjacent regions. – Roczn. Bieszcz. 16: 233–252.
- Mitka J. & Novikoff A. V. (2011): The genus *Aconitum* in Ukraine. – In: Materiały XIII Z'jazdu Ukraińskoho botaničnogo tovarystva (Lviv, 19–23 veresnja 2011 r.), p. 65. Lviv.
- Mitka J. & Starmühler W. (2000): Phenetic variability of *Aconitum lasiocarpum* (Rchb.) Gayer (Ranunculaceae): extension of taxonomic and geographic borders. – Acta Soc. Bot. Polon. 69: 145–155.
- Mitka J., Sutkowska A., Ilnicki T. & Joachimiak A. J. (2007): Reticulate evolution of high-alpine *Aconitum* (Ranunculaceae) in the Eastern Sudetes and Western Carpathians (Central Europe). – Acta Biol. Cracov., Bot. 49: 15–26.
- Mitka J., Boroń P. & Sutkowska A. (2013): Holocene history of *Aconitum* in the Polish Western Carpathians and adjacent regions: long-distance migrations or cryptic refugia? – Modern Phytomorphol. 3: 9–18.
- Mráz P., Chrtek J. jr., Fehrer J. & Plačková I. (2005): Rare recent natural hybridization in *Hieracium* s. str. – evidence from morphology, allozymes and chloroplast DNA. – Pl. Syst. Evol. 255: 177–192.
- Mráz P., Chrtek J. jr. & Fehrer J. (2011b): Interspecific hybridization in the genus *Hieracium* s. str.: evidence for bidirectional gene flow and spontaneous allopolyploidization. – Pl. Syst. Evol. 293: 237–245.
- Mucher W. (1993): Systematics and chorology of *Aconitum* ser. *Toxicum* (Ranunculaceae) in Europe. – Phytomorphol. 33: 51–76.
- Novikoff A. V. & Mitka J. (2011a): Taxonomy and ecology of the genus *Aconitum* L. in the Ukrainian Carpathians. – Wulfenia 18: 37–61.
- Novikov A. V. & Mitka J. (2011b). Rid *Aconitum* v Ukraïns'kykh Karpatach [The genus *Aconitum* in the Ukrainian Carpathians]. – Biol. Stud. 5: 153–172.
- Starmühler W. (1996a): *Aconita rarissima* selecta. – Fritschiana 7: 11–39.
- Starmühler W. (1997): Ranunculaceae subtrib. Delphiniinae exsiccatae, Fasc. 2 (Nr. 21–40). – Fritschiana 10: 1–11.
- Starmühler W. (1998a): Ranunculaceae subtrib. Delphiniinae exsiccatae, Fasc. 3 (Nr. 41–60). – Fritschiana 13: 1–10.
- Starmühler W. (1998b): *Aconita rarissima* selecta (1998). – Fritschiana 13: 11–39.
- Starmühler W. (1999): *Aconita rarissima* selecta (1999). – Fritschiana 18: 11–38.
- Starmühler W. (2000): Fortschritte in der Systematik der Gattung *Aconitum* im Siebenbürgen. – In: Heltnmann H. & Killyen H. (eds), Naturwissenschaftliche Forschungen über Siebenbürgen VI. Beiträge zur Geographie, Botanik, Zoologie und Paläontologie, p. 9–24. Böhlau Verlag, Köln, Weimar & Wien.
- Starmühler W. (2002): Ranunculaceae subtrib. Delphiniinae exsiccatae, Fasc. 5 (Nr. 81–100). – Fritschiana 30: 1–12.
- Sutkowska A., Boroń P. & Mitka J. (2013): Natural hybrid zone of *Aconitum* species in the Western Carpathians: Linnaean taxonomy and ISSR fingerprinting. – Acta Biol. Crac., Bot. 55: 114–126.

Taxon	↑	W _c	W _s	W _i	R _c	R _s	R _i	G _c	G _s	G _i	D _s	S _c	S _s	S _i	F _{mc}	F _{ms}	F _{mi}	F _{pc}	F _{ps}	F _{dc}	F _{ds}	F _{di}	H _c	H _s	H _i
Taxon	↓	W _c	W _s	W _i	R _c	R _s	R _i	G _c	G _s	G _i	D _s	S _c	S _s	S _i	F _{mc}	F _{ms}	F _{mi}	F _{pc}	F _{ps}	F _{dc}	F _{ds}	F _{di}	H _c	H _s	H _i
<i>Astragalus peterfii</i>								+																	
<i>Astragalus pseudopurpureus</i>					+			+																	
<i>Astragalus roemeri</i>								+																	
<i>Athamanta turbith</i> subsp. <i>hungarica</i>					+			+										+							
<i>Aubrieta columnae</i> subsp. <i>platycarpa</i>					+																				
<i>Barbarea lepuznica</i>			+	+																					
<i>Bromus monocladus</i>					+			+										+		+					
<i>Campanula carpatica</i>		+			+	+	+	+	+	+								+		+			+		
<i>Campanula crassipes</i>					+			+										+		+					
<i>Campanula glomerata</i> subsp. <i>subcapitata</i>										+															
<i>Campanula kladniana</i>					+	+	+	+	+	+	+			+				+							+
<i>Campanula serrata</i>	+				+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Campanula tatrae</i>					+	+	+	+	+	+	+		+	+	+	+	+								
<i>Campanula xylocarpa</i>					+	+		+					+					+		+			+		
<i>Cardamine glanduligera</i>				+	+			+	+	+			+	+	+	+	+			+	+	+	+	+	+
<i>Carduus kernerii</i> subsp. <i>kernerii</i>					+	+		+	+				+												+
<i>Carduus kernerii</i> subsp. <i>lobulatifolius</i>								+																	
<i>Carduus lobulatus</i>					+			+																	
<i>Centaurea reichenbachii</i>								+																	
<i>Centaurea rodnensis</i>								+																	
<i>Centaurea simonkaiana</i>								+		+															
<i>Centaurea triniifolia</i>					+			+																	
<i>Cephalaria radiata</i>								+																	
<i>Cephalaria uralensis</i> subsp. <i>multifida</i>								+																	
<i>Cerastium arvense</i> subsp. <i>lerchenfeldianum</i>					+			+																	
<i>Cerastium tatrae</i>	+		+		+	+		+	+		+					+									
<i>Cerastium transsilvanicum</i>								+			+	+													
<i>Chrysosplenium alpinum</i>	+		+	+	+	+			+					+											
<i>Clinopodium pulegium</i>					+			+																	
<i>Cochlearia borzaeana</i>		+																							
<i>Cochlearia tatrae</i>			+			+			+																
<i>Cota tinctoria</i> subsp. <i>fussii</i>							+																		+
<i>Crocus banaticus</i>										+												+			
<i>Crocus discolor</i>								+	+	+	+	+	+	+											
<i>Cyanus dominii</i> subsp. <i>dominii</i>					+			+										+							
<i>Cyanus dominii</i> subsp. <i>slovenicus</i>					+			+										+							
<i>Cyanus dominii</i> subsp. <i>sokolensis</i>					+			+				+									+				
<i>Cyanus maramarosensis</i>										+	+			+							+	+			
<i>Cyanus mollis</i>					+			+	+	+	+	+		+	+						+				+
<i>Cyanus pinnatifidus</i> subsp. <i>pinnatifidus</i>					+	+		+	+																
<i>Cyanus pinnatifidus</i> subsp. <i>sooanus</i>								+																	

	↑	W _c	W _s		R _c	R _s		G _c	G _s	G _i	D _s	S _c	S _s	S _i												
Taxon	↓	W _c	W _s	W _i	R _c	R _s	R _i	G _c	G _s	G _i	D _s	S _c	S _s	S _i	S _i	F _{mc}	F _{ms}	F _{mi}	F _{pc}	F _{ps}	F _{dc}	F _{ds}	F _{di}	H _c	H _s	H _i
<i>Cyclamen purpurascens</i> subsp. <i>immaculatum</i>																					+			+		
<i>Dactylorhiza cordigera</i> subsp. <i>siculorum</i>			+	+																						
<i>Dactylorhiza maculata</i> subsp. <i>schurii</i>			+																							
<i>Daphne arbuscula</i>					+			+											+							
<i>Delphinium elatum</i> subsp. <i>nacladense</i>					+			+		+	+															
<i>Delphinium oxysepalum</i>					+	+		+	+	+	+		+			+	+									
<i>Delphinium simonkaianum</i>					+			+		+											+					
<i>Dianthus callizonus</i>					+			+																		
<i>Dianthus carthusianorum</i> subsp. <i>tenuifolius</i>					+	+		+	+	+												+				
<i>Dianthus giganteus</i> subsp. <i>banaticus</i>								+								+					+					
<i>Dianthus glacialis</i> subsp. <i>gelidus</i>					+	+		+	+		+															
<i>Dianthus henteri</i>						+	+			+																
<i>Dianthus nitidus</i>								+	+		+	+							+							
<i>Dianthus praecox</i> subsp. <i>praecox</i>					+	+		+	+		+								+					+		
<i>Dianthus praecox</i> subsp. <i>lumnitzeri</i>						+			+																	
<i>Dianthus praecox</i> subsp. <i>pseudopraecox</i>									+																	
<i>Dianthus spiculifolius</i>					+	+		+	+			+									+					
<i>Doronicum carpaticum</i>	+	+			+	+	+	+	+				+		+			+								
<i>Draba doreri</i>						+																				
<i>Draba haynaldii</i>					+			+																		
<i>Draba kotschyi</i>					+																					
<i>Draba lasiocarpa</i> subsp. <i>klasterskyi</i>						+			+																	
<i>Draba simonkaiana</i>							+																			
<i>Erigeron hungaricus</i>					+			+	+																	
<i>Eritrichium jankae</i>					+																					
<i>Erysimum hungaricum</i>									+																	
<i>Erysimum pieninicum</i>									+																	
<i>Erysimum wahlenbergii</i>					+	+		+	+				+			+					+			+		
<i>Erysimum witmannii</i> subsp. <i>witmannii</i>						+			+							+					+					
<i>Erysimum witmannii</i> subsp. <i>pallidiflorum</i>									+												+					
<i>Erysimum witmannii</i> subsp. <i>transsilvanicum</i>						+			+																	
<i>Euphorbia carpatica</i>				+		+			+		+												+			
<i>Euphorbia sojakii</i>				+						+													+			
<i>Euphrasia exaristata</i>		+						+																+		
<i>Euphrasia slovacae</i> subsp. <i>slovaca</i>								+	+		+	+														
<i>Euphrasia stipitata</i>									+																	
<i>Euphrasia tatrae</i>					+			+	+		+	+			+											
<i>Ferula sadlerana</i>						+			+		+		+								+					
<i>Festuca amethystina</i> subsp. <i>orientalis</i>								+																		
<i>Festuca bucegiensis</i>								+	+																	
<i>Festuca carpatica</i>		+	+			+	+	+	+			+	+	+	+	+					+					

	↑	W _c	W _s			R _c		R _s		G _c	G _s		G _i		D _s		S _c	S _s		S _i																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																	
--	---	----------------	----------------	--	--	----------------	--	----------------	--	----------------	----------------	--	----------------	--	----------------	--	----------------	----------------	--	----------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

[illegible]

[illegible]

- Dihoru G. & Pârvu C. (1987): Plante endemice în flora Românie [Endemic plants in the flora of Romania]. – Editura Ceres, București.
- Goliašová K. (ed.) (1997): Flóra Slovenska V/2 [Flora of Slovakia V/2]. – Veda, Bratislava.
- Goliašová K. & Michalková E. (eds) (2006): Flóra Slovenska V/3 [Flora of Slovakia V/3]. – Veda, Bratislava.
- Goliašová K. & Michalková E. (eds) (2012): Flóra Slovenska VI/3 [Flora of Slovakia VI/3]. – Veda, Bratislava.
- Goliašová K. & Šípošová H. (eds) (2002): Flóra Slovenska V/4 [Flora of Slovakia V/4]. – Veda, Bratislava.
- Goliašová K. & Šípošová H. (eds) (2008): Flóra Slovenska VI/1 [Flora of Slovakia VI/1]. – Veda, Bratislava.
- Gynda L. (2004): Variation of two endemic taxa of *Galium* L. (Rubiaceae) from the Eastern Carpathians. – Visnyk Lviv Univ., Ser. Biol. 36: 96–100.
- Hadač E. (1987): Plant ecological notes from the Belianske Tatry Mts. – Biol. Práce 33/3: 5–95.
- Hegedúšová Vantarová K. & Škodová I. (eds) (2014): Rastlinné spoločenstvá Slovenska 5. Travinno-bylinná vegetácia [Plant communities of Slovakia 5. Grassland vegetation]. – Veda, Bratislava.
- Jarolímek I. & Šibík A. (eds) (2008): Diagnostic, constant and dominant species of the higher vegetation units of Slovakia. – Veda, Bratislava.
- Jarolímek I., Zaliberová M., Mucina L. & Mochnacký S. (1997): Rastlinné spoločenstvá Slovenska 2. Synantropná vegetácia [Plant communities of Slovakia 2. Synanthropic vegetation]. – Veda, Bratislava.
- Kanka R. (2008): Lesy Belianskych Tatier [Forests of the Belianske Tatry Mts]. – Veda, Bratislava.
- Kliment J. (1999): Komentovaný prehľad vyšších rastlín flóry Slovenska, uvádzaných v literatúre ako endemické taxóny [Annotated summary of vascular plants of the Slovak flora reported as endemic taxa in literature]. – Bull. Slov. Bot. Spoločn. 21, Suppl. 4: 1–434.
- Kliment J. & Jarolímek I. (1995): The *Rumex alpinus* communities in Slovakia. – Biologia 50: 349–365.
- Kliment J. & Valachovič M. (eds) (2007): Rastlinné spoločenstvá Slovenska 4. Vysokohorská vegetácia [Plant communities of Slovakia 4. High-mountain vegetation]. – Veda, Bratislava.
- Kliment J., Šibík J., Šibíková I., Jarolímek I., Dúbravcová Z. & Uhlířová J. (2010): High-altitude vegetation of the Western Carpathians – a syntaxonomical review. – Biologia 65: 965–989.
- Kliment J., Šibíková I. & Šibík J. (2011): On the occurrence of the arctic-alpine and endemic species in the high-altitude vegetation of the Western Carpathians. – Thaiszia-J. Bot. 21: 45–61.
- Kobiv J. (2012a): Roľ prýdatných mikrooselyšč u samovidovlenni populacij rídkisnich vydiv roslyn Ukraíns'skych Karpat [Role of suitable microhabitats in population recruitment of rare plant species in the Ukrainian Carpathians]. – Ukraíns'k. Bot. Zhurn. 69: 178–189.
- Kochjarová J., Valachovič M., Bureš P. & Mráz P. (2006): The genus *Cochlearia* L. in the Eastern Carpathians and adjacent areas. – Bot. J. Linn. Soc. 151: 355–364.
- Kricsfalussy V. V. (2013): Mountain grasslands of high conservation value in the Eastern Carpathians: syntaxonomy, biodiversity, protection and management. – Thaiszia-J. Bot. 23: 67–112.
- Kučera P. (2012): Vegetačný stupeň smrečín v Západných Karpatoch – rozšírenie a spoločenstvá [Norway spruce altitudinal vegetation zone in the Western Carpathians – distribution and communities]. – Botanická záhrada UK, Biatnica.
- Lendvay B. & Kalapos T. (2014): Population dynamics of the climate-sensitive endangered perennial *Ferula sadleriana* Ledeb. (Apiaceae). – Pl. Spec. Biol. 29: 138–151.
- Malynov's'kyj K. A. & Kričfalušij V. V. (2002): Roslynni ugrupovannja vysokohir'ja Ukraíns'skych Karpat [Plant communities of the Ukrainian Carpathian highlands]. – Carpathian Tower Publishing, Uzhgorod.
- Marinov Y., Tosheva A. & Pachedjieva K. (2014): Current status of the rare species *Lathyrus transsilvanicus* in Bulgaria. – Bulgarian J. Agric. Sci. 20, Suppl. 1: 160–164.
- Mikoláš V. (1996): *Sorbus dolomiticola* Mikoláš, a new hybridogenous species of the genus *Sorbus* s. l. from eastern Slovakia. – Thaiszia-J. Bot. 6: 1–12.
- Mikoláš V. (2003): *Sorbus amici-petri* Mikoláš, a new hybridogenous species of the genus *Sorbus* s. l. from eastern Slovakia. – Thaiszia-J. Bot. 13: 127–133.
- Mirek Z. & Piękoś-Mirkowa H. (eds) (2008a): Czerwona księga Karpat Polskich. Rośliny naczyniowe [Red data book of the Polish Carpathians. Vascular plants]. – Instytut Botaniki im. W. Szafera PAN, Kraków.
- Mitka J. (2003): The genus *Aconitum* L. (Ranunculaceae) in Poland and adjacent countries. A phenetic-geographic study. – Institute of botany of the Jagiellonian University, Kraków.
- Mochnacký S. (2006): Cenologické zaradenie endemického druhu ferula Sadlerova (*Ferula sadleriana*) v NP a BR Slovenský kras [On the coenology of *Ferula sadleriana* in the NP and BR Slovenský kras]. – Bull. Slov. Bot. Spoločn. 28: 185–191.

- Oprea A. (2007): Flora and vegetation of the Natural Reserve „Zugreni Gorges“ (Suceava County). – Rom. J. Biol., Plant Biol. 52: 89–121.
- Petrík A., Šibík J. & Valachovič M. (2005): The class *Carici rupestris-Kobresietea bellardii* Ohba 1974 also in the Western Carpathians. – Hacquetia 4: 33–51.
- Piękoś-Mirkowa H., Mirek Z. & Miechówka A. (1996): Endemic vascular plants in the Polish Tatra Mts. Distribution and ecology. – Polish Bot. Stud. 12: 1–107.
- Popescu A. & Sanda V. (1998): Conspectul florei cormofitelor spontane din România [Enumeration of spontaneous vascular plants of Romania]. – Editura Universității din București, București.
- Sanda V., Öllner K. & Burescu P. (2008): Fitocenozele din România. Sintaxonomie, structură, dinamică și evoluție [The plant communities from Romania. Syntaxonomy, structure, dynamics and evolution]. – Ars Docendi, București.
- Slezák M., Hrivnák R. & Petrášová A. (2014): Numerical classification of alder carr and riparian alder forests in Slovakia. – Phytocoenologia 44: 283–308.
- Šibík J., Kliment J., Jarolímek I., Dúbravcová Z., Bělohávková R. & Pačlová L. (2006): Syntaxonomy and nomenclature of the alpine heaths (the class *Loiseleurio-Vaccinietea*) in the Western Carpathians. – Hacquetia 5: 37–71.
- Šibík J., Šibíková I. & Kliment J. (2010): The subalpine *Pinus mugo*-communities of the Carpathians with a European perspective. – Phytocoenologia 40: 155–188.
- Ujházyová M. (2007): Syntaxonómia bukových lesov na vápencoch bradlového pásma [Syntaxonomy of beech forests on the limestone of klippen belt]. – Technická univerzita, Zvolen.
- Valachovič M. (ed.) (1995a): Rastlinné spoločenstvá Slovenska 1. Pionierska vegetácia [Plant communities of Slovakia 1. Pioneer vegetation]. – Veda, Bratislava.
- Valachovič M. (1995b): *Papaverion tatrici*, a vicarious alliance of alpine limestone-scrub communities in the Western Carpathians. – Biologia 50: 377–390.
- Valachovič M. (ed.) (2001): Rastlinné spoločenstvá Slovenska 3. Vegetácia mokradí [Plant communities of Slovakia 3. Wetland vegetation]. – Veda, Bratislava.
- Vlčko J., Hrivnák R., Kochjarová J., Blanár D. & Sádovský M. (2007): *Scilla bifolia* agg. – fytoecologická väzba slovenských populácií [*Scilla bifolia* agg. – phytosociological affinity of Slovak populations]. – In: Hrubá V. & Štykar J. (eds), Geobiocenologie a její aplikace. Geobiocenologické spisy 11, p. 138–146.

Electronic Appendix 7. – Overview of higher syntaxa with the occurrence of vascular plants endemic and subendemic to the Carpathians and classification of phytosociological alliances to habitat groups and habitats. Habitat preferences of individual taxa is shown in Appendix 6.

Abbreviations:

Habitat groups: W – wetlands, R – rocky habitats, G – grasslands, D – dwarf shrubs, S – shrubs, F – forests, H – human-made (anthropogenic) habitats

Vertical distribution: ↓ – submontane and montane vegetation, ↑ – subalpine and alpine vegetation

Geological bedrock: c – vegetation on calcareous bedrock, s – vegetation on silicate (non-calcareous) bedrock (incl. effusive rocks, mylonites, schists etc.), i – vegetation indifferent to the substrate

Subdivision of forest habitats: Fm – montane spruce forests, Fp – pine forests, Fd – deciduous and mixed forests

a) wetlands (W)

Scheuchzeria palustris-*Caricetea fuscae* Tx. 1937

Caricetalia davallianae Br.-Bl. 1949

↓ **Wc** *Caricion davallianae* Klika 1934

Caricetalia fuscae Koch 1926

↑ **Ws** *Drepanocladion exannulati* Krajina 1933

↓ **Ws** *Caricion fuscae* Koch 1926

(Syn.: *Caricion canescenti-nigrae* Nordhagen 1936)

↓ **Ws** *Caricion lasiocarpae* Vanden Berhen in Lebrun et al. 1949

↓ **Ws** *Sphagno*-*Caricion canescentis* Passarge (1964) 1978

Scheuchzerietalia palustris Nordhagen 1936

↑ **Ws** *Sphagnion cuspidati* Krajina 1933

Oxycocco-Sphagnetalia Br.-Bl. et Tx. ex Westhoff et al. 1946

Sphagnetalia medii Kästner et Flössner 1933

↑ **Ws** *Oxycocco microcarpi*-*Empetrium hermaphroditi* Nordhagen ex Hadač et Váňa 1967

↓ **Ws** *Sphagnion medii* Kästner et Flössner 1933

(Syn.: *Sphagnion fuscii* Br.-Bl. 1926)

Montio-Cardaminetalia Br.-Bl. et Tx. ex Klika et Hadač 1944

Montio-Cardaminetalia Pawłowski in Pawłowski et al. 1928

↑ **Ws** *Cardamino-Montion* Br.-Bl. 1926

↑ **Ws** *Cratoneuro filicini-Calthion laetae* Hadač 1983

↑ **Wc** *Cratoneurion commutati* Koch 1928

↓ **Wc** *Lycopodo europaei-Cratoneurion commutati* Hadač 1983

Cardamino-Chrysosplenietalia Hinterlang 1992

↓ **Wi** *Caricion remotae* Kästner 1941

Phragmito-Magnocaricetea Klika in Klika et Novák 1941

Phragmitetalia Koch 1926

↓ **Wi** *Magnocaricion elatae* Koch 1926

b) rocky habitats (R)

Asplenietea trichomanis (Br.-Bl. in Meier et Br.-Bl. 1934) Oberd. 1977

Potentilletalia caulescentis Br.-Bl. in Br.-Bl. et Jenny 1926

↑ **Rc** *Potentillion caulescentis* Br.-Bl. in Br.-Bl. et Jenny 1926

↑ **Rc** *Gypsophilion petraeae* Borhidi et Pócs in Borhidi 1957

↓ **Rc** *Cystopteridion* Richard 197

(Syn.: *Moehringion muscosae* Horvat & Horvatić 1962 p. p.)

↓ **Rc** *Micromerion pulegii* Boşcaiu (1971) 1979

(Syn.: *Micromerion banaticum* Boşcaiu 1971; *Moehringion muscosae* Horvat & Horvatić 1962 p. p.)

↓ *Tortulo-Cymbalarietalia* Segal 1969

↓ **Rc** *Cymbalario-Asplenion* Segal 1969

Androsacetalia vandellii Br.-Bl. 1934

↑ **Rs** *Silenion lerchenfeldianae* Simon 1958

↓ **Rs** *Asplenion septentrionalis* Gams in Oberd. 1938

(Syn.: *Androsacion vandellii* sensu Coldea 1991)

↓ **Rs** *Hypno-Polypodium vulgaris* Mucina 1993

↓ **Rc** *Erysimo witmannii-Hackelion deflexae* Bernátová 1986

(Note: The alliance is ordered within the *Artemisietea vulgaris* class, but it includes natural relic communities)

Thlaspietalia rotundifolii Br.-Bl. 1948

Thlaspietalia rotundifolii Br.-Bl. in Br.-Bl. et Jenny 1926

↑ **Rc** *Papaverion tatrici* Pawłowski et al. 1928 corr. Valachovič 1995

↑**Rc** *Papavero-Thymion pulcherrimi* Pop 1968
Androsacetalia alpinae Br.-Bl. in Br.-Bl. et Jenny 1926
 ↑**Rs** *Androsacion alpinae* Br.-Bl. in Br.-Bl. et Jenny 1926
 ↑**Rs** *Veronicion baumgartenii* Coldea 1991
 (Syn.: *Androsacion alpinae* auct. roman.)
Galio-Parietalia officinalis Boşcaiu et al. 1966
 ↓**Rc** *Stipion [Achnatherion] calamagrostis* Jenny-Lips ex Br.-Bl. et al. 1952
 (Syntax. syn.: *Teucrion montani* Csűrös et Pop 1965)
 ↓**Rc** *Arabidion alpinae* Béguin 1972
 ↓**Rc** *Parietarion officinalis* Gergely et al. 1966
Epilobietalia fleischerii Moor 1958
 ↓**Ri** *Epilobion fleischerii* G. Br.-Bl. ex Br.-Bl. 1949
 (Syn.: *Salicion incanae* Aichinger 1933)

Sedo-Scleranthetea Br.-Bl. 1955
Sedo-Scleranthetalia Br.-Bl. 1955
 ↓**Rs** *Sedo-Scleranthion biennis* Br.-Bl. 1955
 ↓**Rs** *Arabidopsidion thalianae* Passarge 1964
 ↓**Rs** *Hyperico perforati-Scleranthion perennis* Moravec 1967
Alysso-Sedetalia Moravec 1967
 ↓**Rc** *Alysso alyssoidis-Sedion albi* Oberd. et Th. Müller in Th. Müller 1961

c) graslands (G)

Festuco-Brometea Br.-Bl. et Tx. ex Soó 1947
Stipo pulcherrimae-Festucetalia pallentis Pop 1968
 ↓**Gs** *Asplenio septentrionalis-Festucion pallentis* Zólyomi 1936 corr. Soó 1957
 ↓**Gc** *Bromo pannonici-Festucion pallentis* Zólyomi 1966
 (Syn. *Seslerio-Festucion pallentis* Klika 1931 corr. Zólyomi 1966; Syntax. syn.: *Thymo comosi-Festucion rupicola* Pop 1968)
 ↓**Gc** *Diantho lumnitzeri-Seslerion* (Soó 1971) Chytrý et Mucina in Mucina et Kolbek 1993
 ↓**Gc** *Chrysopogono-Festucion pseudodalmaticae* Coldea et Sârbu in Coldea 2012
 ↓**Gc** *Pimpinello-Thymion zygoidi* Dihoru et Donița 1970
Festucetalia valesiaca Soó 1947
 ↓**Gc** *Festucion valesiaca* Klika 1931
 (Syntax. syn.: *Festucion rupicola* Soó 1940 corr. 1964)
 ↓**Gc** *Stipion lessingiana* Soó 1947
 ↓**Gc** *Jurineo arachnoideae-Euphorbion stepposae* (Dobrescu 1970) Coldea et Sârbu in Coldea 2012
Brometalia erecti Br.-Bl. 1936
 ↓**Gc** *Bromion erecti* Koch 1926
 ↓**Gc** *Cirsio-Brachypodion pinnati* Hadač et Klika ex Klika 1951
 (Syn.: *Danthonio-Stipion stenophyllae* Soó 1947; Syntax. syn.: *Danthonio-Brachypodion pinnati* Boşcaiu 1970)
 ↓**Gs** *Chrysopogono-Danthonion calycinae* Kojić 1959

Festucetea vaginatae Soó ex Vicherek 1972
Festucetalia vaginatae Soó 1957
 ↓**Gi** *Festucion vaginatae* Soó 1929
 ↓**Gc** *Bassio laniflorae-Bromion tectorum* (Soó 1957) Borhidi 1996

Koelerio-Corynepherea Klika in Klika et Novák 1941
Corynepherea canescentis Klika 1934
 ↓**Gs** *Corynephorion canescentis* Klika 1931

Trifolio-Geranietea sanguinei Th. Müller 1962
Origanetalia vulgaris Th. Müller 1962
 ↓**Gi** *Geranion sanguinei* Tx. in Th. Müller 1962
 ↓**Gi** *Trifolion medii* Th. Müller 1962

Molinio-Arrhenatheretea Tx. 1937
Arrhenatheretalia elatioris Tx. 1931
 ↓**Gi** *Arrhenatherion elatioris* Luquet 1926
 ↓**Gi** *Cynosurion cristati* Tx. 1947
Poo alpinae-Trisetetalia Ellmauer et Mucina 1993
 ↓**Gc** *Polygono bistortae-Trisetion flavescentis* Br.-Bl. et Tx. ex Marshall 1947
 ↑**Gi** *Poion alpinae* Gams ex Oberd. 1950
Molinietalia caerulea Koch 1926
 ↓**Gi** *Molinion caerulea* Koch 1926
 ↓**Gi** *Calthion palustris* Tx. 1937 (incl. *Filipendulenion*)
 ↓**Gi** *Deschampsion caespitosae* Horvatić 1930

Nardetea strictae Rivas Goday in Rivas Goday et Rivas-Mart. 1963

Nardetalia strictae Oberd. ex Preising 1949

↑Gs *Nardion strictae* Br.-Bl. 1926

↑Gs *Potentillo ternatae-Nardion* Simon 1958

↓Gi *Nardo-Agrostion tenuis* Sillinger 1933

↓Gi *Violion caninae* Schwickerath 1944

Mulgedio-Aconitetea Hadač et Klika in Klika et Hadač 1944

(Syn. *Betulo-Adenostyletea* Br.-Bl. 1948)

Calamagrostietalia villosae Pawłowski et al. 1928

↑Gs *Calamagrostion villosae* Pawłowski et al. 1928

excl. *Calamagrostio villosae-Salicetum helveticae* Dúbravcová et Šeffler 1992 → ↑Ss

excl. *Trisetio fusci-Salicetum hastatae* Coldea (1986) 1990 → ↑Ss

↑Gs *Trisetion fusci* Krajina 1933

(Syntax. syn.: *Aconition firmi* Krajina 1933)

excl. *Deschampsio cespitosae-Salicetum helveticae* (Krajina 1933) Dúbravcová et Šeffler 1992 → ↑Ss

Gc *Calamagrostion arundinaceae* (Luquet 1926) Oberd. 1957

↑*Helianthemo grandiflorae-Calamagrostietum arundinaceae*

↑*Anemomo narcissiflorae-Laserpitietum latifolii*

↑*Allio victorialis-Calamagrostietum villosae*

↓other associations of alliance *Calamagrostion arundinaceae*

(see Kliment & Valachovič 2007 for details)

Gc *Calamagrostion varia* Sillinger 1932

↑*Geranio sylvatici-Calamagrostietum varia*

↓*Convallario-Calamagrostietum varia*

↑Gc *Festucion carpaticae* Bělohávková et Fišerová 1989

Adenostyletalia alliariae Br.-Bl. 1930

Adenostylium alliariae Br.-Bl. 1926

excl. *Salici silesiacae-Alnetum viridis* Colić et al. 1962 → ↑Si (*Alnion viridis*)

excl. *Pulmonario filarszkyanae-Alnetum viridis* Pawłowski et Walas 1949 → ↑Ss (*Alnion viridis*)

↑Gs *Adenostylenion alliariae* Klika in Klika et Hadač 1944

↑Gc *Delphinienion elati* (Hadač ex Hadač et al. 1969) Boşcaiu et Mihăilescu 1997

Petasito-Chaerophylletalia Morariu ex Kopecký 1969

↓Gi *Petasition officinalis* Sillinger 1933

Elyno-Seslerietea Br.-Bl. 1948

Seslerietalia coerulae Br.-Bl. in Br.-Bl. et Jenny 1926

Gc *Astero alpini-Seslerion calcariae* Hadač ex Hadač et al. 1969

↑*Astero alpini-Seslerienion calcariae* Kliment et al. 2005

↓*Pulsatillo slavicae-Caricion humilis* Uhlířová in Kliment et al. 2010

↑Gc *Seslerion tatrae* Pawłowski 1935 corr. Klika 1955

↑Gc *Caricion firmae* Gams 1936

↑Gc *Festuco saxatilis-Seslerion bielzii* (Pawłowski et Walas 1949) Coldea 1984

↑Gc *Bellardiochloion violaceae* Sanda et al. 2001

↓Gc *Seslerion rigidae* Zólyomi 1939

Carici rupestris-Kobresietea bellardii Ohba 1974

Oxytropido-Elynetalia Oberd. ex Albrecht 1969

↑Gc *Oxytropido-Elynion myosuroidis* Br.-Bl. (1948) 1949

↑Gs *Festucion versicoloris* Krajina 1933

Caricetea curvulae Br.-Bl. 1948 nom. cons. propos.

(Syntax. syn.: *Juncetea trifidi* Hadač in Klika et Hadač 1944 p. p.)

Caricetalia curvulae Br.-Bl. in Br.-Bl. et Jenny 1926

↑Gs *Juncion trifidi* Krajina 1933

↑Gs *Caricion curvulae* Br.-Bl. 1925

Salicetea herbaceae Br.-Bl. in Br.-Bl. et Jenny 1926

Salicetalia herbaceae Br.-Bl. in Br.-Bl. et Jenny 1926

↑Gs *Salicion herbaceae* Br.-Bl. in Br.-Bl. et Jenny 1926

↑Gs *Festucion picturatae* Krajina 1933 corr. Dúbravcová 2007

Arabidetalia caerulae Br.-Bl. 1948

↑Gc *Arabidion caerulae* Br.-Bl. in Br.-Bl. et Jenny 1926

↑Gc *Salicion retusae* Horvat 1949

d) dwarf shrubs (D)

Loiseleurio procumbentis-Vaccinietea Eggler ex Schubert 1960

Rhododendro ferruginei-Vaccinietalia Br.-Bl. in Br.-Bl. et Jenny 1926

↑Ds *Loiseleurio procumbentis-Vaccinion* Br.-Bl. in Br.-Bl. et Jenny 1926

(syn.: *Cetrario-Loiseleurion* (Br.-Bl. et al. 1939) Schubert 1960)
 ↑**Ds** *Vaccinion myrtilli* Krajina 1933
 ↑**Ds** *Rhododendro-Vaccinion* Br.-Bl. ex Schnyder 1930
 ↑**Ds** *Junipero-Bruckenthalion spiculifoliae* (Horvat 1949) Boşcaiu 1971
 ↑**Ds** *Juniperion nanae* Br.-Bl. in Br.-Bl. et al. 1939

Calluno-Ulicetea Br.-Bl. et Tx. ex Klika et Hadač 1944
Vaccinio myrtilli-Genistetalia pilosae Schubert ex Passarge 1964
 ↓**Ds** *Genisto pilosae-Vaccinion* Br.-Bl. 1926

e) scrubs (S)

Crataego-Prunetea Tx. 1956
 (Syn.: *Rhamno-Prunetea* Rivas Goday et Borja Carbonell ex Tx. 1962)
Prunetalia spinosae Tx. 1952
 ↓**Si** *Prunion fruticosae* Tx. 1952
 ↓**Si** *Berberidion vulgaris* Br.-Bl. 1950
 (Syn.: *Prunion spinosae* Soó 1931)
 ↓**Si** *Corylo avellanae-Populion tremulae* Br.-Bl. ex Jurko 1964

Betulo carpaticae-Alnetea viridis Rejmánek in Huml et al. 1979
Alnetalia viridis Rübel ex Huml et al. 1979
 ↑**Si** *Alnion viridis* Schnyder 1930
Salici silesiacae-Alnetum viridis Colić et al. 1962
Pulmonario filarszkyanae-Alnetum viridis Pawłowski et Walas 1949
 ↑**Sc** *Salicion silesiacae* Rejmánek et al. 1971
 Note: The following associations belong to subalpine shrubs, too:
 ↑**Ss** *Calamagrostio villosae-Salicetum helveticae* Dúbravcová et Šeffler 1992
 ↑**Ss** *Deschampsio cespitosae-Salicetum helveticae* (Krajina 1933) Dúbravcová et Šeffler 1992
 ↑**Ss** *Trisetio fuscii-Salicetum hastatae* Coldea (1986) 1990

Roso pendulinae-Pinetea mugo Theurillat in Theurillat et al. 1995
Junipero-Pinetalia mugo Boşcaiu 1971
 ↑**Si** *Pinion mugo* Pawłowski in Pawłowski et al. 1928
 (RO: only the association *Rhododendro myrtifolii-Pinetum mugo* Coldea 1991)
 excl.: *Campanulo abietinae-Juniperetum nanae* → ↑**Ds**, *Campanulo abietinae-Vaccinietum myrtilli* → ↑**Ds**,
Rhododendro myrtifolii-Vaccinietum gaultherioidis → ↑**Ds**
 excl.: *Rhododendro myrtifolii-Piceetum* → **Fms**, *Bruckenthalio-Piceetum* → **Fms**
 excl. *Saxifrago cuneifoliae-Laricetum* → **Fmi**

f) forests (F)

Quercetea pubescentis Doing Kraft ex Scamoni et Passarge 1959
Quercetalia pubescenti-petraeae Klika 1933
 ↓**Fdc** *Quercion pubescenti-petraeae* Br.-Bl. 1932
 ↓**Fdi** *Aceri tatarici-Quercion Zólyomi* et Jakucs 1957
 ↓**Fds** *Quercion petraeae* Issler 1931
 ↓**Fdi** *Quercion confertae-cerris* Horvat 1954
 ↓**Fdc** *Syringo-Carpinion orientalis* Jakucs (1959) 1960
 ↓**Fdi** *Querco-Carpinion orientalis* Csűrös et al. 1968

Quercetea robori-petraeae Br.-Bl. et Tx. ex Oberd. 1957
 ↓**Fds** *Quercetalia roboris* Tx. 1931
 ↓**Fds** *Pino-Quercion* Medwecka-Kornaś et al. 1959

Alnetea glutinosae Br.-Bl. et Tx. ex Westhoff et al. 1946
Alnetalia glutinosae Tx. 1937
 ↓**Fdi** *Alnion glutinosae* Malcuit 1929

Carpino-Fagetea sylvaticae Jakucs ex Passarge 1968
 (Syn.: *Querco-Fagetea* Br.-Bl. et Vlieger in Vlieger 1937 p. p.)
Fagetalia sylvaticae Pawłowski in Pawłowski et al. 1928
 ↓**Fdi** *Alnion incanae* Pawłowski in Pawłowski et al. 1928
 (Syn.: *Alno-Ulmion* Br.-Bl. et Tx. ex Tschou 1948)
Alnenion glutinoso-incanae Oberd. 1953
 excl. *Arunco vulgaris-Salicetum capreae* Hadač et al. ex Petrášová et Jarolímek 2014 → ↓**Sc**, ↓**Ss**
 ↓**Fdi** *Carpinion betuli* Issler 1931
 ↓**Fdi** *Lathyro hallersteinii-Carpinenion* (Boşcaiu 1974) Boşcaiu et al. 1982
 (Syn.: *Lathyro hallersteinii-Carpinion* Boşcaiu 1974)
 ↓*Fagion sylvaticae* Luquet 1926
Fdc *Cephalanthero-Fagenion* Tx. in Tx. et Oberd. 1958

- Fdi** *Eu-Fagenion* Oberd. 1957
- Fdi** *Acerenion* Oberd. 1957
- Fdi** *Galio-Abietenion* Oberd. 1962
- ↓**Fdi** *Symphyto cordati-Fagion* (Vida 1963) Täuber 1982
- Fdi** *Symphyto cordati-Fagenion* Vida 1963
- Fdc** *Moehringio muscosae-Acerenion* Boşcaiu et al. 1982
- Fdc** *Epipactido-Fagenion* Boşcaiu et al. 1982
- Fds** *Calamagrostio-Fagenion* Boşcaiu et al. 1982
- ↓**Fds** *Luzulo-Fagion sylvaticae* Lohmeyer et Tx. in Tx. 1954
- ↓**Fdi** *Tilio platyphylli-Acerion* Klika 1955

Erico-Pinetea Horvat 1959

Erico-Pinetalia Horvat 1959

- ↓**Fpc** *Pulsatillo slavicae-Pinion* Fajmonová 1978
- ↓**Fpc** *Seslerio rigidae-Pinion* Coldea 1991

Piceetea excelsae Klika 1948

(Syn.: *Vaccinio-Piceetea* Br.-Bl. in Br.-Bl. et al. 1939)

Piceetalia excelsae Pawłowski in Pawłowski et al. 1928

- ↓**Fms** *Piceion excelsae* Pawłowski in Pawłowski et al. 1928

(Syntax. syn.: *Athyrio alpestris-Piceion* Sýkora 1971)

incl.: *Chrysanthemo rotundifolii-Piceetum* Krajina 1933 (sensu orig.), *Hieracio rotundati-Piceetum* Pawłowski et Br.-Bl. 1939

Athyrio filicis-feminae-Piceetalia Hadač ex Hadač et al. 1969

- ↓**Fmc** *Oxalido-Piceion* Hadač et al. 1969

(see Kučera 2012 for details)

Vaccinio-Pinetalia Scamoni et Passarge 1959

- ↓**Fps** *Dicrano-Pinion sylvestris* (Libbert 1932) Matuszkiewicz 1962

(Syn.: *Vaccinio-Pinion* (Libbert 1933) Passarge 1968)

Robinietaea Jurko ex Hadač et Sofron 1980

- ↓**Fdi** *Chelidonio-Robinietaea* Jurko ex Hadač et Sofron 1980
- Robinetum pseudacaciae* auct.

h) human-made habitats: anthropogenic vegetation (H)

Epilobietea angustifolii Tx. et Preising ex von Rochow 1951

Atropetalia Vlieger 1937

- ↓**Hi** *Carici piluliferae-Epilobion angustifolii* Tx. 1950
- ↓**Hi** *Atropion* Br.-Bl. ex Aichinger 1933

Galio-Urticetea Passarge ex Kopecký 1969

Lamio albi-Chenopodietalia boni-henrici Kopecký 1969

- ↓**Hi** *Impatienti noli-tangere-Stachyion sylvaticae* Görs ex Mucina in Mucina et al. 1993
- ↓**Hi** *Carduo-Urticion dioicae* Hadač ex Hadač et al. 1969
- ↓**Hi** *Rumicion alpini* Scharfetter 1938

Artemisietea vulgaris Lohmeyer et al. ex von Rochow 1951

Agropyretalia repentis Görs 1966

- ↓**Hi** *Convolvulo arvensis-Agropyron repentis* Görs 1966